

Invariant Representations of $\mathrm{GSp}(2)$ under Tensor Product with a Quadratic Character

Ping Shun Chan

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS,
OHIO 43210

E-mail address: pschan@math.ohio-state.edu

2000 *Mathematics Subject Classification.* Primary 11F70, 11F72, 11F85

Key words and phrases. $\mathrm{GSp}(2)$, automorphic representations, p -adic harmonic analysis

ABSTRACT. Let F be a number field or a p -adic field. We introduce in Chapter 2 of this work two reductive rank one F -groups, $\mathbf{H}_1, \mathbf{H}_2$, which are twisted endoscopic groups of $\mathrm{GSp}(2)$ with respect to a fixed quadratic character ε of the idèle class group of F if F is global, $F^\times$ if F is local. When F is global, Langlands functoriality predicts that there exists a canonical lifting of the automorphic representations of $\mathbf{H}_1, \mathbf{H}_2$ to those of $\mathrm{GSp}(2)$. In Chapter 4, we establish this lifting in terms of the Satake parameters which parametrize the automorphic representations. By means of this lifting we provide a classification of the discrete spectrum automorphic representations of $\mathrm{GSp}(2)$ which are invariant under tensor product with ε .

The techniques through which we arrive at our results are inspired by those of Kazhdan's in [K], which introduced the trace formula twisted by a character. In particular, our techniques involve comparing the spectral sides of the trace formulas for the groups under consideration. We make use of the twisted extension of Arthur's trace formula, and Kottwitz-Shelstad's stabilization of the elliptic part of the geometric side of the twisted trace formula.

When F is local, in Chapter 5 we provide a classification of the irreducible admissible representations of $\mathrm{GSp}(2, F)$ which are invariant under tensor product with the quadratic character ε of $F^\times$. More precisely, we use the global results from Chapter 4 to express the twisted characters of these invariant representations in terms of the characters of the admissible representations of $\mathbf{H}_i(F)$ ($i = 1, 2$). These (twisted) character identities provide candidates for the liftings predicted by the conjectural local Langlands functoriality. The proofs are inspired by [K] and rely on Sally-Tadić's classification of the irreducible admissible representations of $\mathrm{GSp}(2, F)$, and on Flicker's results on the lifting from $\mathrm{PGSp}(2)$ to $\mathrm{PGL}(4)$.

In both the global and local cases, our results depend on the conjectural statement that there exist smooth functions on the groups which have matching (twisted) orbital integrals. In particular, these results depend on the validity of the (twisted) Fundamental Lemma in the context of $\mathrm{GSp}(2)$ and $\mathbf{H}_1, \mathbf{H}_2$. We prove in Appendix B the Fundamental Lemma in our case, making use of Flicker's work on the twisted Fundamental Lemma for $\mathrm{GL}(4)$ and $\mathrm{GSp}(2)$, and Hales' reduction of the Fundamental Lemma to unit elements.

Contents

Technical Preface	vii
ε -Endoscopic Groups	vii
Global Lifting	vii
Local Lifting	viii
Chapter 1. Introduction	1
1.1. An Overview	1
1.2. ε -Invariant Automorphic Representations	3
1.3. Local Character Identities	6
1.4. Statement of Main Results	7
1.5. Assumptions and Restrictions	13
Chapter 2. ε -Endoscopy for $\mathrm{GSp}(2)$	15
2.1. Endoscopic Data	15
2.2. Endoscopic group $\mathbf{H}_1$	17
2.3. Endoscopic group $\mathbf{H}_2$	19
2.4. Norm Correspondence	21
2.5. Matching Functions	25
Chapter 3. The Trace Formula	27
3.1. The fine χ -expansion	27
3.2. Comparison of Geometric Sides of Trace Formulas	38
3.3. Application of Kottwitz-Shelstad's Formula	41
Chapter 4. Global Lifting	43
4.1. The ε -Trace Identity	43
4.2. Frobenius-Hecke Classes	50
4.3. Packets	55
4.4. Contributions	64
4.5. Some Global Lifting Results	73
4.6. Final Words	78
Chapter 5. The Local Picture	83
5.1. Induced Representations	85
5.2. Induced Representations—Split Case	94
5.3. ε -Invariant Packets	96
5.4. Character Identities for Unstable Packets	97
5.5. Character Identities for Stable Packets	110
Appendix A. Summary of Global Lifting	121
A.1. Unstable (quasi-)packets of G	121

A.2. Stable (quasi-)packets	121
A.3. Induced representations	122
Appendix B. Fundamental Lemma	125
B.1. Norm Correspondence—Elliptic Elements	126
B.2. Comparison of Orbital Integrals	131
Bibliography	151
List of Symbols	153
Index	156

Technical Preface

ε -Endoscopic Groups

Let F be a number field. Let ε be a quadratic character of the idèle class group C_F of F . Let E be the quadratic extension of F which corresponds to ε via global class field theory. The quasi-split ε -endoscopic groups of $\mathbf{G}$ over F are as follows:

- (1) $\mathbf{H}_1 = (\mathrm{GL}(2) \times \mathrm{R}_{E/F}\mathbb{G}_m)'$, where $\mathrm{R}_{E/F}$ is the restriction of scalars (from E to F) functor. The prime denotes the condition that the determinant of the $\mathrm{GL}(2)$ -factor is equal to the norm of the $\mathrm{R}_{E/F}\mathbb{G}_m$ -factor in $\mathbb{G}_m$. In particular, the group of F -points of $\mathbf{H}_1$ is

$$\mathbf{H}_1(F) = \{(g, x) \in \mathrm{GL}(2, F) \times E^\times : \det g = \mathrm{N}_{E/F}x\}.$$

- (2) $\mathbf{H}_2$ is the unique quasi-split reductive F -group whose group of F -points is

$$\mathbf{H}_2(F) = (\mathrm{GL}(2, E) \times F^\times) / \{(\mathrm{diag}(z, z), \mathrm{N}_{E/F}z^{-1}) : z \in E^\times\}.$$

Global Lifting

We say that an automorphic representation π of $\mathrm{GL}(2, \mathbb{A}_F)$ is E -**monomial** if it is the monomial representation $\pi(\theta)$ associated with a character θ of C_E . For a representation π , let ω_π denote its central character. Let $B_{E/F}\pi$ denote the $\mathrm{GL}(2, \mathbb{A}_E)$ -module which is the base change of a $\mathrm{GL}(2, \mathbb{A}_F)$ -module π .

1. PROPOSITION. *Let π be an irreducible automorphic $\mathrm{GL}(2, \mathbb{A}_F)$ -module which is either cuspidal non- E -monomial or one dimensional. The packets $\pi \otimes_1 1$ of $\mathbf{H}_1(\mathbb{A}_F)$ and $B_{E/F}\pi \otimes_2 \omega_\pi$ of $\mathbf{H}_2(\mathbb{A}_F)$ lift to an unstable (quasi-)packet $\{\Pi\}$ of $\mathrm{GSp}(2, \mathbb{A}_F)$, which in turn lifts to the induced representation $I_{(2,2)}(\pi, \varepsilon\pi)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.*

REMARK. If π is one dimensional, $\{\Pi\}$ belongs to a class of quasi-packets constructed by Howe, Piatetski-Shapiro.

For an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$, let $\pi(\tau)$ denote the $\mathrm{GL}(4, \mathbb{A}_F)$ -module which is obtained from τ via the twisted endoscopic lifting from $\mathrm{R}_{E/F}\mathrm{GL}(2)$ to $\mathrm{GL}(4)$. let ${}^\sigma\tau$ denote the representation

$${}^\sigma\tau : (g_{ij}) \mapsto \tau(\sigma g_{ij}), \quad \forall (g_{ij}) \in \mathrm{GL}(2, \mathbb{A}_E).$$

1. PROPOSITION. *Let π be an irreducible, cuspidal, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$, μ a character of C_F , such that $\pi \neq {}^\sigma\pi$ and $\omega_\pi = \mu \circ \mathrm{N}_{E/F}$. The representation $\pi \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ lifts to a stable packet $\{\Pi\}$, which in turn lifts to the cuspidal automorphic representation $\pi(\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$. No discrete spectrum packet of $\mathbf{H}_1(\mathbb{A}_F)$ lifts to the packet $\{\Pi\}$.*

2. PROPOSITION. *Suppose $\{\Pi\}$ is a global packet of $\mathrm{GSp}(2, \mathbb{A}_F)$ which is the lift of some automorphic representation/packet of one of the ε -endoscopic groups. Then, $\{\Pi\}$*

contains at least one representation which is ε -invariant. If $\{\Pi\}$ is a discrete spectrum global packet of $\mathrm{GSp}(2, \mathbb{A}_F)$ which is not the lift of any discrete spectrum representation/packet of the ε -endoscopic groups, then $\{\Pi\}$ does not contain any discrete spectrum automorphic representation which is ε -invariant.

Local Lifting

Let F, E now be local nonarchimedean fields, with $[E : F] = 2$. Let ε be the nontrivial quadratic character of $F^\times$ corresponding to $\mathcal{K}/F$. Let $G = \mathrm{GSp}(2, F)$, $H_1 = \mathbf{H}_1(F)$, $H_2 = \mathbf{H}_2(F)$.

Let Z be the center of G . For an irreducible ε -invariant admissible representation π of G , and a smooth function f on G , put

$$\langle \pi, f \rangle_A := \mathrm{tr} \int_{Z \backslash G} \pi(g) f(g) A \, dg,$$

where A is an intertwining operator in $\mathrm{Hom}_G(\pi, \varepsilon\pi)$ such that $A^2 = 1$.

For $i = 1, 2$, let $Z_0(H_i)$ denote the maximal F -split component of the center of H_i . For an admissible representation π_i of H_i , and a smooth function f_i on H_i , put

$$\langle \pi_i, f_i \rangle := \mathrm{tr} \int_{Z_0(H_i) \backslash H_i} \pi_i(h_i) f_i(h_i) \, dh_i.$$

2. PROPOSITION. *Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial, admissible $\mathrm{GL}(2, F)$ -module whose central character ω_τ is trivial. Let $\{\pi^+, \pi^-\}$ be the local packet of G which lifts to the induced representation $I_{(2,2)}(\tau, \varepsilon\tau)$ of $\mathrm{PGL}(4, F)$ according to [F4]. There exist intertwining operators A^+ in $\mathrm{Hom}_G(\pi^+, \varepsilon\pi^+)$, and A^- in $\mathrm{Hom}_G(\pi^-, \varepsilon\pi^-)$, such that the following holds for matching functions:*

$$\begin{aligned} \langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} &= \langle \tau \otimes 1, f_1 \rangle, \\ \langle \pi^+, f \rangle_{A^+} - \langle \pi^-, f \rangle_{A^-} &= \langle B_{\mathcal{K}/F} \tau \otimes_2 \omega_\tau, f_2 \rangle. \end{aligned}$$

3. PROPOSITION. *Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial, admissible representation of $\mathrm{GL}(2, F)$, χ be a character of $\mathcal{K}^\times$, such that: There does not exist a character ε' of $F^\times$ such that ${}^\sigma\chi/\chi = \varepsilon' \circ \mathrm{N}_{\mathcal{K}/F}$ and $\tau \cong \varepsilon'\tau$. Then, there exist a collection $[\pi]$ of distinct ε -invariant, irreducible, cuspidal, admissible G -modules, and an operator $A(\pi') \in \mathrm{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:*

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes 1, \chi, f_1 \rangle.$$

4. PROPOSITION. *Let τ be an irreducible, cuspidal, admissible representation of $\mathrm{GL}(2, \mathcal{K})$, not fixed by the action of $\mathrm{Gal}(\mathcal{K}/F)$, such that its central character ω_τ is equal to $\mu \circ \mathrm{N}_{\mathcal{K}/F}$ for some character μ of $F^\times$. Then, there exist a collection $[\pi]$ of distinct ε -invariant, irreducible, cuspidal, admissible G -modules, and an operator $A(\pi') \in \mathrm{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:*

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes_2 \mu, f_2 \rangle.$$

Ping Shun Chan
February 2, 2008
Columbus, Ohio, USA

CHAPTER 1

Introduction

1.1. An Overview

1.1.1. Langlands Functoriality. In the 1960's, Robert Langlands conjectured that there exists a canonical (not necessarily one-to-one) correspondence between Galois representations and automorphic representations of reductive algebraic groups, and that this correspondence coincides with the Artin reciprocity law when the underlying group is $\mathrm{GL}(1)$. More precisely, for every reductive group $\mathbf{H}$ over a number field F with Langlands dual ${}^L\mathbf{H}$, it is believed that each homomorphism from the Langlands group L_F —a conjectural group generalizing the Galois and Weil groups—to ${}^L\mathbf{H}$ parametrizes an L -packet of automorphic representations of $\mathbf{H}$.

Suppose there are two reductive groups $\mathbf{H}, \mathbf{G}$ with L -groups ${}^L\mathbf{H}, {}^L\mathbf{G}$, respectively, such that there is a homomorphism from ${}^L\mathbf{H}$ to ${}^L\mathbf{G}$. Then, each homomorphism from the Langlands group to ${}^L\mathbf{H}$ induces one from the Langlands group to ${}^L\mathbf{G}$. Consequently, one should expect a canonical lifting of the automorphic representations of $\mathbf{H}$ to those of $\mathbf{G}$. A more precise formulation of this principle is as follows:

CONJECTURE (Langlands Functoriality). *Let $\mathbf{H}, \mathbf{G}$ be reductive groups over a number field such that there is an L -homomorphism ξ of the L -group of ${}^L\mathbf{H}$ of $\mathbf{H}$ into that of $\mathbf{G}$. Then, for every automorphic representation π of $\mathbf{H}$, there is a packet $\{\Pi\}$ of automorphic representations of $\mathbf{G}$, parametrized by the same Frobenius-Hecke classes at all but finitely many places, such that the Frobenius-Hecke classes in ${}^L\mathbf{H}$ parametrizing π are mapped by ξ to those parametrizing $\{\Pi\}$ in ${}^L\mathbf{G}$.*

Over the years, various examples of Langlands functorial lifting have been proved via the comparison of trace formulas, a technique introduced by Langlands and his collaborators ([JL], [L], [LL]). We list here a few notable cases which are of particular interest in relation to our work. They are all examples of (twisted) endoscopic liftings (see [KS]).

- The lifting from $\mathrm{R}_{E/F}\mathrm{GL}(1)$ to $\mathrm{GL}(2)$, where $[E : F] = 2$, proved by Jacquet and Langlands [JL].
- Cyclic base change lifting for $\mathrm{GL}(2)$, proved by Langlands [L], and for $\mathrm{GL}(n)$, by Arthur and Clozel [AC].
- The lifting from elliptic algebraic tori to $\mathrm{SL}(2)$, by Labesse and Langlands [LL].
- The lifting from $\mathrm{R}_{E/F}\mathrm{GL}(1)$ to $\mathrm{GL}(n)$, E/F cyclic of order n , by Kazhdan [K].
- The lifting from $\mathrm{R}_{E/F}\mathrm{GL}(m)$ to $\mathrm{GL}(n)$, where $m[E : F] = n$, by Waldspurger [Wa].
- The lifting from $\mathrm{PGSp}(2)$ to $\mathrm{PGL}(4)$, by Flicker [F4].
- Endoscopic lifting for $\mathrm{GSp}(2)$, by Arthur [A3].

The first half of this work uses the trace formula technique to establish the lifting of automorphic representations of two rank one twisted endoscopic groups $\mathbf{H}_1$ and $\mathbf{H}_2$ to the

symplectic group of similitudes $\mathrm{GSp}(2)$ (4×4 matrices). At the same time, we show that the lifting provides a classification of the automorphic representations of $\mathrm{GSp}(2)$ which are invariant under tensor product with a fixed quadratic character.

To a great extent, the conception of this project and the methodology employed are inspired by the work of Kazhdan's [K], which uses the trace formula technique both to establish Langlands functorial lifting from $\mathrm{R}_{E/F}\mathbf{G}_m$ to $\mathrm{GL}(n)$, where E/F is cyclic of degree n , and to classify the automorphic representations of $\mathrm{GL}(n, \mathbb{A}_F)$ which are invariant under tensor product with a fixed character of order n . In particular, he showed that each idèle class character θ of E lifts to an automorphic representation $\pi(\theta)$ of $\mathrm{GL}(n, \mathbb{A}_F)$. Moreover, an automorphic representation of $\mathrm{GL}(n, \mathbb{A}_F)$ is invariant under tensor product with an order n character ε if and only if it is equal to $\pi(\theta)$ for some character θ of the cyclic field extension E of F defined by ε via class field theory. That Kazhdan was able to obtain these spectacular results is a testament to the power of the trace formula technique.

In this work, we have an analogous situation for $\mathrm{GSp}(2)$, except that in this case there are nonsingleton (quasi-) L -packets, and there are two twisted endoscopic groups instead of one. Modulo the conjectural existence of matching functions and certain technical conditions on the local components of the automorphic representations, we show that the automorphic representations of the two twisted endoscopic groups ($[E : F] = 2$)

$$\mathbf{H}_1 = (\mathrm{GL}(2) \times \mathrm{R}_{E/F}\mathbf{G}_m)' \quad \text{and} \quad \mathbf{H}_2 = (\mathrm{R}_{E/F}\mathrm{GL}(2) \times \mathbf{G}_m) / \mathrm{R}_{E/F}\mathbf{G}_m$$

lift to the (quasi-)packets of $\mathrm{GSp}(2)$ containing at least one automorphic representation which is invariant under tensoring with the quadratic character ε associated with E/F . Moreover, all such packets are the lifts of the automorphic representations of the twisted endoscopic groups. Naturally, an important question is whether every representation in such a packet is invariant under tensoring with ε , but that remains unanswered in this work.

The presence of nonsingleton packets for $\mathrm{GSp}(2)$ also raises interesting questions regarding the stability of the packets. Thanks to the classification results of [A3] and [F4], we show that a packet containing an invariant representation is stable if and only if it is the lift of exactly one of the twisted endoscopic groups.

Another interesting phenomenon is that each discrete spectrum automorphic representation π of $\mathrm{GSp}(2)$ which is invariant under tensoring with ε in turn lifts to an automorphic representation Π of $\mathrm{GL}(4)$ which is invariant under tensor product with the same character. Moreover, Π is induced if and only if π belongs to an unstable (quasi-)packet.

It is also worth mentioning that the lifting from $\mathbf{H}_2$ to $\mathrm{GSp}(2)$ may be viewed as follows: Let E/F be the quadratic field extension which corresponds to the character ε . Each automorphic representation π of $\mathbf{H}_2(\mathbb{A}_F)$ may be constructed from an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$ whose central character is invariant under the action of $\mathrm{Gal}(E/F)$. Let $\varphi : L_E \rightarrow \mathrm{GL}(2, \mathbb{C})$ be the representation of the Langlands group of E which corresponds to τ . Then, the representation of L_F (the Langlands group of F) induced from φ factors through $\mathrm{GSp}(2)$:

$$\begin{array}{ccc} \mathrm{Ind}_{L_E}^{L_F} \varphi : L_F & \xrightarrow{\quad} & \mathrm{GL}(4, \mathbb{C}) , \\ & \searrow & \nearrow \\ & \mathrm{GSp}(2) & \end{array}$$

and the automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$ corresponding to $\mathrm{Ind}_{L_E}^{L_F} \varphi$ is the lift of π . This is beautifully analogous to the construction of the monomial representations $\pi(\theta)$ of $\mathrm{GL}(2)$ in [JL] and [K].

It should be noted here that automorphic representations of $\mathrm{GSp}(2)$ have received the attentions of mathematicians for quite some time. In addition to those cited above, it suffices to mention Siegel, Shimura, Weissauer, Howe, Piatetski-Shapiro, Vigneras, D. Prasad, B. Roberts, and R. Schmidt. Note that many mathematicians prefer to use $\mathrm{GSp}(4)$ rather than $\mathrm{GSp}(2)$ to denote the same group.

1.1.2. Local Lifting. Langlands Functoriality as stated above has a local analogue. Namely, it is conjectured that, for any reductive group $\mathbf{H}$ over a local p -adic field F , the homomorphisms from the Weil group W_F (or L_F) to ${}^L H$ correspond canonically to packets of admissible representations of $\mathbf{H}(F)$. Consequently, analogously to the global case, a homomorphism between L -groups should parallel the lifting of admissible representations from one p -adic group to another.

Since each square integrable admissible representation is a local component of an automorphic representation, it is a general philosophy that one can derive a large class of local liftings from global liftings. Works in the past which establish global lifting using trace formulas typically also derive local lifting. More precisely, they express the local lifts in the form of trace identities among admissible representations. For example, for each admissible representation π of $\mathrm{GL}(n, F)$ which is invariant under tensor product with a character ε of $F^\times$ of order n , Kazhdan establishes in [K] an expression of the ε -twisted Harish-Chandra character of π in terms of the Galois orbit of a character of $E^\times$, where E is a degree n cyclic extension of F determined by ε via local class field theory.

The second half of this work is occupied with deriving local twisted trace identities among admissible representations of $\mathrm{GSp}(2)$ and those of its twisted endoscopic groups. We arrive at these identities using our global lifting results.

1.2. ε -Invariant Automorphic Representations

Let F be a number field. Let $\mathbb{A}_F$ denote the ring of adèles of F and C_F the idèle class group $F^\times \backslash \mathbb{A}_F^\times$. Let V denote the set of places of F . For each finite place v of F , let $\mathcal{O}_v$ denote the ring of integers of F_v . Let $\mathbf{H}$ be a reductive algebraic group over F . Put $H := \mathbf{H}(F)$, $H_v := \mathbf{H}(F_v)$ for any place v of F .

For any element γ of any group, let subscript γ denote the centralizer of γ in that group. For example, H_γ denotes the centralizer in H of γ . Let $\mathbf{Z}$ be the center of $\mathbf{H}$. Let $\mathbf{Z}_0$ be the maximal F -split component of $\mathbf{Z}$. Put $Z := \mathbf{Z}(F)$, $Z_0 := \mathbf{Z}_0(F)$. For the groups $\mathrm{GSp}(2)$ and $\mathrm{GL}(4)$, which are F -split, there is no distinction between $\mathbf{Z}$ and $\mathbf{Z}_0$.

Let ω be a character of $Z_0 \backslash \mathbf{Z}_0(\mathbb{A}_F)$. Let $L(\mathbf{H}(\mathbb{A}_F), \omega)$ be the space of measurable functions ϕ on $H \backslash \mathbf{H}(\mathbb{A}_F)$ such that $\phi(zg) = \omega(z)\phi(g)$ for all $z \in \mathbf{Z}_0(\mathbb{A}_F)$ and

$$\int_{\mathbf{Z}_0(\mathbb{A}_F)H \backslash \mathbf{H}(\mathbb{A}_F)} |\phi(h)|^2 dh < \infty.$$

Let ρ , or ρ_ω to emphasize the dependence on ω , be the right regular representation of $\mathbf{H}(\mathbb{A}_F)$ on $L(\mathbf{H}(\mathbb{A}_F), \omega)$, that is:

$$(\rho_\omega(h)\phi)(g) = \phi(gh), \quad \forall h, g \in \mathbf{H}(\mathbb{A}_F), \phi \in L(\mathbf{H}(\mathbb{A}_F), \omega).$$

In particular, the restriction of the central character of ρ_ω to $\mathbf{Z}_0(\mathbb{A}_F)$ is equal to ω . We say that a representation of $\mathbf{H}(\mathbb{A}_F)$ is **automorphic** if it occurs in ρ_ω for some character ω of $Z_0 \backslash \mathbf{Z}_0(\mathbb{A}_F)$. Let $\mathcal{A}(\mathbf{H}, \omega)$ denote the set of equivalence classes of irreducible automorphic

representations π of $\mathbf{H}(\mathbb{A}_F)$ with the property that their central characters restrict to ω on $\mathbf{Z}_0(\mathbb{A}_F)$.

Write $\pi \in \mathcal{A}(\mathbf{H}, \omega)$ as a tensor product $\otimes_{v \in V} \pi_v$ of representations π_v of H_v . Since ρ_ω is continuous, π too is continuous with respect to the topology on $\mathbf{H}(\mathbb{A}_F)$ (see [PR]). Consequently, at almost every finite place v , the local representation π_v is unramified, i.e. it contains a nonzero vector fixed by the hyperspecial maximal compact subgroup $\mathbf{H}(\mathcal{O}_v)$.

As an $\mathbf{H}(\mathbb{A}_F)$ -module, $L(\mathbf{H}(\mathbb{A}_F), \omega)$ decomposes into a direct sum

$$L_d(\mathbf{H}(\mathbb{A}_F), \omega) \oplus L_c(\mathbf{H}(\mathbb{A}_F), \omega).$$

Here, $L_d(\mathbf{H}(\mathbb{A}_F), \omega)$ is the closed span of the irreducible, closed, invariant subspaces of $L(\mathbf{H}(\mathbb{A}_F), \omega)$ and $L_c(\mathbf{H}(\mathbb{A}_F), \omega)$ is its orthogonal complement. Let ρ_d, ρ_c denote the restrictions of $\rho = \rho_\omega$ to $L_d(\mathbf{H}(\mathbb{A}_F), \omega), L_c(\mathbf{H}(\mathbb{A}_F), \omega)$, respectively. We call the $\mathbf{H}(\mathbb{A}_F)$ -module $L_d(\mathbf{H}(\mathbb{A}_F), \omega)$ (resp. $L_c(\mathbf{H}(\mathbb{A}_F), \omega)$) the **discrete** (resp. **continuous**) **spectrum** of $\mathbf{H}(\mathbb{A}_F)$.

We say that the algebraic group $\mathbf{H}$ has the **multiplicity one** property if each automorphic representation of $\mathbf{H}(\mathbb{A}_F)$ with central character ω on $\mathbf{Z}_0(\mathbb{A}_F)$ occurs *at most once* in the discrete spectrum of ρ_ω . By the multiplicity one theorem for $\mathrm{GL}(2)$ and a result in [LL], the multiplicity one property holds for the twisted endoscopic groups studied in this work. In the case of $\mathrm{GSp}(2)$, the multiplicity one property for automorphic representations with trivial central characters is established in [F4]. For arbitrary central characters, Arthur has announced in [A3] that multiplicity one holds at least for those representations which are associated with semisimple Arthur parameters.

REMARK. An automorphic representation is really a representation which occurs in the space of slowly increasing functions called **automorphic forms**. This space is not the same as $L^2(\mathbf{H}(\mathbb{A}_F), \omega)$. However, the discrete spectrum of this space (as an $\mathbf{H}(\mathbb{A}_F)$ -module) consists of L^2 functions; thus, it coincides with the discrete spectrum of $L^2(\mathbf{H}(\mathbb{A}_F), \omega)$; see [BJ].

For any place $v \in V$, let $C(H_v, \omega_v)$ be the space of smooth functions f_v on H_v such that f_v is compactly supported modulo $Z_{0,v}$ and $f_v(zh) = \omega_v^{-1}(z)f_v(h)$ for all z in $Z_{0,v}$ and h in H_v .

If v is finite, and $\mathbf{H}$ is defined over $\mathcal{O}_v$, let K_v denote the hyperspecial maximal compact subgroup $\mathbf{H}(\mathcal{O}_v)$ of H_v . Let $\mathcal{H}(H_v, \omega_v)$ denote the Hecke algebra of K_v -biinvariant functions in $C(H_v, \omega_v)$. The algebra $\mathcal{H}(H_v, \omega_v)$ is nonzero if and only if ω_v is trivial on $Z_{0,v} \cap K_v$. If v is archimedean, we fix a maximal compact subgroup K_v of H_v . Let $\mathcal{H}(H_v, \omega_v)$ be the set of K_v -finite functions in $C(H_v, \omega_v)$.

Let $C(\mathbf{H}(\mathbb{A}_F), \omega)$ be the span of the smooth functions $f = \otimes_{v \in V} f_v$ on $\mathbf{H}(\mathbb{A}_F)$ which are compactly supported modulo $\mathbf{Z}_0(\mathbb{A}_F)$ such that $f_v \in C(H_v, \omega_v)$ for all $v \in V$ and f_v is the unit element in the Hecke algebra $\mathcal{H}(H_v, \omega_v)$ for almost all finite v .

Fix a Tamagawa measure (see [PR]) on $\mathbf{H}(\mathbb{A}_F)$. For any $f \in C(\mathbf{H}(\mathbb{A}_F), \omega)$ and $\pi \in \mathcal{A}(\mathbf{H}, \omega)$, let $\pi(f)$ denote the following convolution operator on the space of π :

$$\begin{aligned} \pi(f) &:= \int_{\mathbf{Z}_0(\mathbb{A}_F) \backslash \mathbf{H}(\mathbb{A}_F)} \pi(h) f(h) dh \\ &= \int_{\mathbf{Z}(\mathbb{A}_F) \backslash \mathbf{H}(\mathbb{A}_F)} \pi(h) \left(\int_{\mathbf{Z}_0(\mathbb{A}_F) \backslash \mathbf{Z}(\mathbb{A}_F)} \omega(z) f(zh) dz \right) dh. \end{aligned}$$

It has finite rank. Let $\langle \pi, f \rangle$ denote the trace of $\pi(f)$.

1.2.1. The ε -Twisted Trace Formula. Recall that ρ_ω is the right regular representation of $\mathbf{H}(\mathbb{A}_F)$ in $L(\mathbf{H}(\mathbb{A}_F), \omega)$. Let ε be a homomorphism in $\text{Hom}(\mathbf{H}(\mathbb{A}_F), \mathbb{C}^\times)$ which is trivial on $\mathbf{Z}(\mathbb{A}_F)H$. Let π be an automorphic representation in $\mathcal{A}(\mathbf{H}(\mathbb{A}_F), \omega)$. Let V_π be the space of π . By definition, π is a subquotient of ρ_ω . Let $\varepsilon\pi$ be the representation of $\mathbf{H}(\mathbb{A}_F)$ on the space V_π of π defined as follows:

$$\varepsilon\pi := \varepsilon \otimes \pi : h \mapsto \varepsilon(h)\pi(h), \quad \forall h \in \mathbf{H}(\mathbb{A}_F).$$

Note that the space $V_{\varepsilon\pi} = V_\pi$ of $\varepsilon\pi$ is a subquotient of $L(\mathbf{H}(\mathbb{A}_F), \omega)$, but $\varepsilon\pi$ is not the right regular representation of $\mathbf{H}(\mathbb{A}_F)$ on $V_{\varepsilon\pi}$.

We say that π is ε -invariant if $\pi \cong \varepsilon\pi$; i.e. there exists an automorphism A on V_π such that $A\pi(h) = \varepsilon\pi(h)A$ for all $h \in \mathbf{H}(\mathbb{A}_F)$. By Schur's lemma, A^2 is a scalar, and we normalize A (multiplying it by $(A^2)^{-1/2}$ if necessary) so that $A^2 = 1$. By this choice of normalization, the operator A is unique up to a sign.

Define an operator $\rho_\omega(\varepsilon)$, or $\rho(\varepsilon)$ for brevity, on $L(\mathbf{H}(\mathbb{A}_F), \omega)$ as follows:

$$(\rho_\omega(\varepsilon)\phi)(h) := \phi(h)\varepsilon(h), \quad \forall \phi \in L(\mathbf{H}(\mathbb{A}_F), \omega), h \in \mathbf{H}(\mathbb{A}_F).$$

The restriction of $\rho(\varepsilon)$ to the space V_π is a nontrivial intertwining operator of $\varepsilon\pi$ into ρ . In particular, $\varepsilon\pi$ is equivalent to the automorphic representation $\rho|_{\rho(\varepsilon)V_\pi}$ of $\mathbf{H}(\mathbb{A}_F)$.

Assuming that multiplicity one holds for $\mathbf{H}$, any two irreducible inequivalent submodules of ρ_ω in $L(\mathbf{H}(\mathbb{A}_F), \omega)$ are either orthogonal or equal to each other, each being generated by $\rho(\mathbf{H}(\mathbb{A}_F))$ from a single vector.

Suppose π is irreducible, and $\pi \cong \varepsilon\pi$. In principle, $\rho_\omega(\varepsilon)$ needs not intertwine π with $\varepsilon\pi$, namely $\rho_\omega(\varepsilon)V_\pi$ may be equivalent to V_π as an $\mathbf{H}(\mathbb{A}_F)$ -module but different from it as a subspace of $L(\mathbf{H}, \omega)$. However, since we assume that the multiplicity one property holds, we have $\rho_\omega(\varepsilon)V_\pi = V_\pi$. It then follows that the restriction of $\rho_\omega(\varepsilon)$ to V_π is a nontrivial intertwining operator in $\text{Hom}_{\mathbf{H}(\mathbb{A}_F)}(\pi, \varepsilon\pi)$. By Schur's lemma, $\rho_\omega(\varepsilon)|_{V_\pi}$ is a nontrivial scalar multiple of A ; hence, it is A or $-A$.

For an irreducible automorphic representation π and a function f in $C(\mathbf{G}(\mathbb{A}_F), \omega)$, put

$$\langle \pi, f \rangle_\varepsilon := \text{tr } \pi(f)\rho_\omega(\varepsilon) = \text{tr } \rho_\omega(\varepsilon)\pi(f).$$

If $\pi \not\cong \varepsilon\pi$, then the spaces V_π and $\rho_\omega(\varepsilon)V_\pi$ are disjoint, and hence $\langle \pi, f \rangle_\varepsilon = 0$. Conversely, if π is ε -invariant, we shall show in Lemma 4.39 that the distribution $f \mapsto \langle \pi, f \rangle_\varepsilon$ is nonzero.

By definition, ρ_ω is the direct sum of all automorphic representations with $\mathbf{Z}_0(\mathbb{A}_F)$ -character ω . If we could express $\langle \rho_\omega, f \rangle_\varepsilon$ as the sum (discrete plus continuous)

$$\sum_{\pi \in \mathcal{A}(\mathbf{H}, \omega)} \langle \pi, f \rangle_\varepsilon,$$

then those automorphic representations which are not ε -invariant would drop out from the sum. Thus, at least formally, the distribution $\langle \rho_\omega, \cdot \rangle_\varepsilon : f \mapsto \langle \rho_\omega, f \rangle_\varepsilon$ on the space $C(\mathbf{H}(\mathbb{A}_F), \omega)$ detects those automorphic representations which are ε -invariant. However, the situation is more delicate.

For $f \in C(\mathbf{H}(\mathbb{A}_F), \omega)$, $\rho_\omega(f)\rho_\omega(\varepsilon)$ is an integral operator on $L(\mathbf{H}(\mathbb{A}_F), \omega)$. Let $K_{f, \varepsilon} : \mathbf{H}(\mathbb{A}_F) \times \mathbf{H}(\mathbb{A}_F) \rightarrow \mathbb{C}$ be the kernel of $\rho_\omega(f)\rho_\omega(\varepsilon)$. The trace $\langle \rho_\omega, f \rangle_\varepsilon$ is obtained by integrating $K_{f, \varepsilon}(h, h)$ over $h \in \mathbf{Z}_0(\mathbb{A}_F)H \backslash \mathbf{H}(\mathbb{A}_F)$. The function $h \mapsto K_{f, \varepsilon}(h, h)$ may be expressed as a sum over conjugacy classes in H , and also as a sum over automorphic representations of $\mathbf{H}(\mathbb{A}_F)$. For applications, we would like to change the order of summation and integration in $\int_{\mathbf{Z}_0(\mathbb{A}_F)H \backslash \mathbf{H}(\mathbb{A}_F)} K_{f, \varepsilon}(h, h) dh$. However, this change of order is not justified unless $\mathbf{H}$ is anisotropic over F . Extending Arthur's results in [A], the authors

of [CLL] introduce a truncation $K_{f,\varepsilon}^T(h)$ of $K_{f,\varepsilon}(h, h)$. They show that we may pull the sum out of the integral in $\int_{\mathbf{Z}_0(\mathbb{A}_F)H \backslash \mathbf{H}(\mathbb{A}_F)} K_{f,\varepsilon}^T(h)$. The ε -**twisted Arthur trace formula** (or ε -trace formula) is the following equality of two different ways to express the integral $\int_{\mathbf{Z}_0(\mathbb{A}_F)H \backslash \mathbf{H}(\mathbb{A}_F)} K_{f,\varepsilon}^T(h) dh$:

$$\sum_{\{\mathcal{O}\}} J_{\mathcal{O},\varepsilon}^T(f) = \sum_{\{\chi\}} J_{\chi,\varepsilon}^T(f).$$

The left hand side of the equation is a sum of (weighted) orbital integrals over the set of semisimple conjugacy classes in H . It is called the $\mathcal{O}$ -**expansion** or the **geometric side** of the trace formula. The right hand side is a sum of traces of automorphic representations over a set of spectral data of $\mathbf{H}(\mathbb{A}_F)$. These are data which catalogue the ε -invariant automorphic representations of $\mathbf{H}(\mathbb{A}_F)$. We call $\sum_{\{\chi\}} J_{\chi,\varepsilon}^T(f)$ the **fine χ -expansion** or the **spectral side** of the trace formula.

1.2.2. The Case of $\mathrm{GSp}(2)$. Let $J = \begin{pmatrix} & w \\ -w & \end{pmatrix} \in \mathrm{GL}(4)$, where $w = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \in \mathrm{GL}(2)$. Let $\mathbf{G} = \mathrm{GSp}(2)$, the **symplectic group of similitudes of rank 2**, be the algebraic group over $\mathbb{Z}$ defined as follows:

$$\mathrm{GSp}(2) = \{g \in \mathrm{GL}(4) : {}^t g J g = \lambda(g) J \text{ for some } \lambda(g) \in \mathbb{G}_m\}.$$

We call λ the **similitude character**.

Let F be a number field, and consider $\mathbf{G}$ as a reductive F -group. For any $g \in \mathbf{G}$, we call $\lambda(g)$ its **similitude factor**. The center of $\mathbf{G}$ is $\mathbf{Z} = \{\mathrm{diag}(z, z, z, z) : z \in \mathbb{G}_m\}$. It is F -split. Fix a character ω of $\mathbf{Z} \backslash \mathbf{Z}(\mathbb{A}_F)$. Let ε be a quadratic character of $G \backslash \mathbf{G}(\mathbb{A}_F)$ which factors through the similitude factor. Since ε is quadratic, and $\lambda(\mathrm{diag}(z, z, z, z)) = z^2$, ε is trivial on $\mathbf{Z} \backslash \mathbf{Z}(\mathbb{A}_F)$. Consider the ε -twisted Arthur trace formula for $\mathbf{G}(\mathbb{A}_F)$. As a distribution on $C(\mathbf{G}(\mathbb{A}_F), \omega)$, the $\mathcal{O}$ -expansion $\sum_{\{\mathcal{O}\}} J_{\mathcal{O},\varepsilon}^T(f)$ of the ε -trace formula is not invariant with respect to stable conjugacy. In [KS], Kottwitz and Shelstad express the elliptic regular part of $\sum_{\{\mathcal{O}\}} J_{\mathcal{O},\varepsilon}^T(f)$ (i.e. the part of the sum which is indexed by the elliptic regular conjugacy classes $\mathcal{O}$) in terms of stably invariant $\mathcal{O}$ -expansions of lower rank groups. We call these lower rank groups the ε -endoscopic groups of $\mathbf{G}$. We compute them in Chapter 2.

On the other hand, Arthur's trace formula for the endoscopic groups equate their $\mathcal{O}$ -expansions with their fine χ -expansions. We impose in Section 3.2.1 a condition on the test function $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$ so that all nonelliptic or singular terms in the geometric expansion of the ε -trace formula vanish. Using Kottwitz-Shelstad's formula, we obtain a global trace identity relating the fine χ -expansion of the ε -trace formula of $\mathbf{G}(\mathbb{A}_F)$ with the fine χ -expansions of the trace formulas of the ε -endoscopic groups. We compute in Section 3.1 the fine χ -expansions of the groups, and we obtain in Section 3.3 a global trace identity relating these spectral expansions. We then use the global trace identity to deduce global lifting results in Sections 4.4 and 4.5

1.3. Local Character Identities

Let k now be a local p -adic field. Let $G = \mathbf{G}(k)$ denote the group of F -points of $\mathbf{G}$. Let ε be a quadratic character of $k^\times$. Then ε defines a character of G as follows:

$$\varepsilon(g) := \varepsilon(\lambda(g)), \quad \forall g \in G.$$

For any admissible ([BZ]) representation π of G , define a representation $\varepsilon\pi$ on the space of π by

$$\varepsilon\pi := \varepsilon \otimes \pi : g \mapsto \varepsilon(g)\pi(g), \quad \forall g \in G.$$

We are interested in irreducible admissible representations of G which are ε -**invariant**: A representation π of G is said to be ε -invariant if $\pi \cong \varepsilon\pi$, namely there is an invertible linear operator $A : V_\pi \rightarrow V_\pi$ on the space V_π of π such that $A\pi(g) = \varepsilon(g)\pi(g)A$ for all $g \in G$.

We fix a character ω of the center Z of G . Let $C(G, \omega)$ be the space of smooth functions f on G which are compactly supported modulo Z and satisfy:

$$f(zg) = \omega^{-1}(z)f(g), \quad \forall z \in Z, g \in G.$$

We fix once and for all a right-invariant Haar measure ([BZ]) dg on G . For any admissible representation π of G with central character ω , the following convolution operator on the space of π has finite rank:

$$\pi(f) := \int_{Z \backslash G} \pi(g)f(g) dg.$$

Suppose π is an irreducible, admissible ε -invariant representation. Let A be a nontrivial intertwining operator in $\text{Hom}_G(\pi, \varepsilon\pi)$. By Schur's lemma ([BZ]), A^2 is a scalar multiplication on π . Replacing A with $(A^2)^{-1/2}A$ if necessary, we assume that A satisfies $A^2 = 1$. Then, A is unique up to a sign.

Put $\langle \pi, f \rangle_A := \text{tr} \int_{Z \backslash G} \pi(g)f(g)A dg$. We call the distribution $f \mapsto \langle \pi, f \rangle_A$ the ε -**character** of π . The ε -character depends on the choice of the intertwining operator A . Since $A^2 = 1$, the ε -character is unique up to a sign.

Chapter 5 is devoted to the classification of the ε -invariant representations of G . More precisely, we utilize the global lifting results established in Chapter 4 to express the ε -characters of these representations in terms of the (nontwisted) characters of representations of the twisted endoscopic groups.

1.4. Statement of Main Results

1.4.1. ε -Endoscopic Groups. Let F be a number field, let ε be a quadratic character of the idèle class group C_F of F . Let E be the quadratic extension of F which corresponds to ε via global class field theory. In Chapter 2, we compute the quasi-split ε -endoscopic groups of $\mathbf{G}$ over F . They are as follows:

- (1) $\mathbf{H}_1 = (\text{GL}(2) \times \text{R}_{E/F}\mathbb{G}_m)'$, where $\text{R}_{E/F}$ is the restriction of scalars (from E to F) functor. The prime denotes the condition that the determinant of the $\text{GL}(2)$ -factor is equal to the norm of the $\text{R}_{E/F}\mathbb{G}_m$ -factor in $\mathbb{G}_m$. In particular, the group of F -points of $\mathbf{H}_1$ is

$$\mathbf{H}_1(F) = \{(g, x) \in \text{GL}(2, F) \times E^\times : \det g = \text{N}_{E/F}x\}.$$

- (2) $\mathbf{H}_2$: the unique quasi-split reductive F -group whose group of F -points is

$$\mathbf{H}_2(F) = (\text{GL}(2, E) \times F^\times) / \{(\text{diag}(z, z), \text{N}_{E/F}z^{-1}) : z \in E^\times\}.$$

1.4.2. Global Lifting. The global lifting results are expressed in terms of **global (quasi-)packets**. A global (quasi-)packet consists of adèlic representations which share the same unramified local components (with respect to fixed maximal compact subgroups) at almost every finite place. Equivalently, a global (quasi-)packet may be expressed as a restricted tensor product of **local (quasi-)packets**. We say that a global packet is a **discrete spectrum packet** if it contains a representation which occurs in the discrete spectrum. We say that a discrete spectrum global packet is **stable** if all its members occur with the same nonzero multiplicity in the discrete spectrum. Otherwise we say that it is **unstable**.

Associated with any algebraic F -group $\mathbf{H}$ is an L -**group**, denoted ${}^L H$ (see [Bo]). It is a split extension of a complex Lie group by the Weil group W_F of F (see [T]). Each unramified local component of a global (quasi-)packet of $\mathbf{H}(\mathbb{A}_F)$ is parametrized by a conjugacy class in ${}^L H$, called a **Frobenius-Hecke class** (see [Bo]). We compute these conjugacy classes for the ε -endoscopic groups in Chapter 4.2.

As we shall discuss in Chapter 2, associated with each ε -endoscopic group $\mathbf{H}$ is an L -group embedding $\xi_H : {}^L H \rightarrow {}^L G$. We say that a global packet $\{\Pi_H\}$ of an ε -endoscopic group $\mathbf{H}(\mathbb{A}_F)$ **lifts** to a packet $\{\Pi\}$ of $\mathbf{G}(\mathbb{A}_F)$, if ξ_H maps the Frobenius-Hecke classes in ${}^L H$ parametrizing $\{\Pi_H\}$ to the Frobenius-Hecke classes in ${}^L G$ parametrizing $\{\Pi\}$.

Let π be an irreducible automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, χ a character of C_E . Let V be the set of places of F . For any place $v \in V$, let $\pi_v \otimes_1 \chi_v$ denote the representation of $\mathbf{H}_1(F_v)$ defined as follows.

$$\pi_v \otimes_1 \chi_v : (g, x) \mapsto \chi_v(x) \pi_v(g), \quad \forall (g, x) \in \mathbf{H}_1(F_v).$$

Then, $\pi_v \otimes_1 \chi_v$ may be a reducible representation of $\mathbf{H}_1(F_v)$ with at most two irreducible constituents ([LL]). The set of irreducible constituents of $\pi_v \otimes_1 \chi_v$ form a local packet. If $\pi_v \otimes_1 \chi_v$ is reducible, we name its two constituents π_v^+, π_v^- . For each finite place v , let $K_{1,v}$ be the hyperspecial maximal compact subgroup $\mathbf{H}_1(\mathcal{O}_v)$, where $\mathcal{O}_v$ is the ring of integers in F_v . We say that a representation of $\mathbf{H}_1(F_v)$ is unramified if it contains a nonzero $K_{1,v}$ -fixed vector. If $\pi_v \otimes_1 \chi_v$ is unramified, we let π_v^+ denote its unique irreducible unramified constituent. If $\pi_v \otimes_1 \chi_v$ is irreducible, we put $\pi_v^+ := \pi_v \otimes_1 \chi_v$ and $\pi_v^- := 0$. From [LL], a global packet of $\mathbf{H}_1(\mathbb{A}_F)$ has the form

$$\pi \otimes_1 \chi := \{ \otimes_v \pi_v : \pi_v \in \{ \pi_v^+, \pi_v^- \}, \forall v; \pi_v = \pi_v^+ \text{ for almost all } v \}.$$

For $f_1 \in C(\mathbf{H}_1(\mathbb{A}_F), \omega)$, we have $\langle \pi \otimes_1 \chi, f_1 \rangle = \sum_{\pi' \in \pi \otimes_1 \chi} \langle \pi', f_1 \rangle$.

The group $\mathbf{H}_2(\mathbb{A}_F)$ is a quotient of $\mathrm{GL}(2, \mathbb{A}_E) \times \mathbb{A}_F^\times$. Since rigidity, or strong multiplicity one theorem, holds for $\mathrm{GL}(2)$ and $\mathrm{GL}(1)$, it also holds for $\mathbf{H}_2$. Consequently, each global packet of $\mathbf{H}_2(\mathbb{A}_F)$ consists of a single irreducible automorphic representation. For any automorphic representation π of $\mathrm{GL}(2, \mathbb{A}_E)$ and character μ of C_F such that the central character ω_π of π is equal to $\mu \circ N_{E/F}$, let $\pi \otimes_2 \mu$ denote the representation

$$\pi \otimes_2 \mu : (g, x) \mapsto \mu(x) \pi(g), \quad \forall (g, x) \in \mathbf{H}_2(\mathbb{A}_F).$$

1.4.2.1. Summary of Global Lifting. We now give a summary of our global lifting results. For simplicity's sake, some of these statements are accurate only up to certain nuanced conditions. We mark them with an *. The full, accurate versions of these statements may be found in Sections 4.4, 4.5, and 4.6.

A more concise version of this summary is in Appendix A.

We classify packets of $\mathrm{GSp}(2, \mathbb{A}_F)$ by stating what automorphic representations of $\mathrm{GL}(4, \mathbb{A}_F)$ they lift to via the natural L -group embedding $\mathrm{GSp}(2, \mathbb{C}) \rightarrow \mathrm{GL}(4, \mathbb{C})$. This classification is not one to one. There are cases in which two inequivalent packets of $\mathrm{GSp}(2, \mathbb{A}_F)$ lift to the same representation in $\mathrm{GL}(4, \mathbb{A}_F)$.

We say that an automorphic representation π of $\mathrm{GL}(2, \mathbb{A}_F)$ is E -**monomial** if it is the monomial representation $\pi(\theta)$ associated with a character θ of C_E , or more precisely with the induced two-dimensional representation $\mathrm{Ind}_{W_{E/E}}^{W_{E/F}}(\theta)$ of W_F (see [JL], [K]). The first global lifting result which we examine involves global packets of $\mathbf{H}_1$ of the form $\pi_1 = \pi \otimes_1 \chi$, where π is a cuspidal non- E -monomial or one dimensional automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, and $\chi = \mu \circ N_{E/F}$ for some character μ of C_F . From the way $\mathbf{H}_1$ is defined, $\pi \otimes_1 \mu \circ N_{E/F}$ is equal to $\mu \pi \otimes_1 1$; hence, we assume without

loss of generality that $\pi_1 = \pi \otimes_1 1$. Let ω_π be the central character of π . Let $B_{E/F}\pi$ be the representation of $\mathrm{GL}(2, \mathbb{A}_E)$ which is the base change of π (see [L], [F1]). Then, $\omega_{B_{E/F}\pi} = \omega_\pi \circ N_{E/F}$.

1.1. PROPOSITION. *The packets $\pi \otimes_1 1$ of $\mathbf{H}_1(\mathbb{A}_F)$ and $B_{E/F}\pi \otimes_2 \omega_\pi$ of $\mathbf{H}_2(\mathbb{A}_F)$ lift to a packet $\{\Pi\}$ of $\mathrm{GSp}(2, \mathbb{A}_F)$, which in turn lifts to the induced representation $I_{(2,2)}(\pi, \varepsilon\pi)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.*

REMARK. (i) According to [A3, Sect. 5], the packet $\{\Pi\}$ is unstable. In the case where the central character of $\{\Pi\}$ is trivial, the instability of $\{\Pi\}$ is proven in [F4, V. 10.]. (ii) In [A3], Arthur denotes $\{\Pi\}$ by the symbol $(\pi \boxtimes 1) \boxplus (\varepsilon\pi \boxtimes 1)$ if π is cuspidal non- E -monomial and calls it a Yoshida type packet. If $\pi = \mu 1_{\mathrm{GL}(2, \mathbb{A}_F)}$ for some character μ of C_F , Arthur calls $\{\Pi\}$ a Howe, Piatetski-Shapiro type packet and denotes it by the symbol $(\mu \boxtimes \nu(2)) \boxplus (\varepsilon\mu \boxtimes \nu(2))$, where ν is the (normalized) absolute value function on $\mathbb{A}_F$.

Let σ be the generator of $\mathrm{Gal}(E/F)$. For any character θ of C_E , put ${}^\sigma\theta := \theta \circ \sigma$.

1.2. PROPOSITION. *Let θ, χ be characters of C_E such that none of $\theta, \chi, \theta\chi, \theta{}^\sigma\chi$ is fixed by the action of $\mathrm{Gal}(E/F)$. Let $\pi(\theta), \pi(\chi)$ be the cuspidal monomial representations of $\mathrm{GL}(2, \mathbb{A}_F)$ associated with θ, χ , respectively. The global packets*

- $\pi(\theta) \otimes_1 \chi$,
- $\pi(\theta) \otimes_1 {}^\sigma\chi$,
- $\pi(\chi) \otimes_1 \theta$,
- $\pi(\chi) \otimes_1 {}^\sigma\theta$

all lift to an unstable packet $\{\Pi\}$, which in turn lifts to the induced representation

$$I_{(2,2)}(\pi(\theta\chi), \pi(\theta{}^\sigma\chi))$$

of $\mathrm{GL}(4, \mathbb{A}_F)$. Moreover, no discrete spectrum representation of $\mathbf{H}_2(\mathbb{A}_F)$ lifts to $\{\Pi\}$.

REMARK. Under the notation of [A3], $\{\Pi\}$ is the unstable Yoshida type packet denoted by $(\pi(\theta\chi) \boxtimes 1) \boxplus (\pi(\theta{}^\sigma\chi) \boxtimes 1)$.

For an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$, let $\pi(\tau)$ denote the automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$ which is obtained from τ via the twisted endoscopic lifting from $\mathrm{R}_{E/F}\mathrm{GL}(2)$ to $\mathrm{GL}(4)$ (see [AC, Sect. 3.6]).

1.3. PROPOSITION*. *Let χ be a character of C_E not fixed by $\mathrm{Gal}(E/F)$. Let π be an irreducible, cuspidal, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$ which is not E -monomial. Suppose there does not exist a character ε' of C_F such that ${}^\sigma\chi/\chi$ is equal to $\varepsilon' \circ N_{E/F}$ and $\pi \cong \varepsilon'\pi$. The packet $\pi \otimes_1 \chi$ lifts to a stable packet $\{\Pi\}$ which lifts to the cuspidal automorphic representation $\pi(\chi B_{E/F}\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$. No discrete spectrum representation of $\mathbf{H}_2(\mathbb{A}_F)$ lifts to $\{\Pi\}$.*

1.4. PROPOSITION*. *Let χ be a character of C_E not fixed by $\mathrm{Gal}(E/F)$. The one dimensional representation $1_{\mathrm{GL}(2, \mathbb{A}_F)} \otimes_1 \chi$ lifts to a stable quasi-packet, denoted by*

$$\{L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))\},$$

which lifts to the Langlands quotient representation $J(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. No discrete spectrum representation of $\mathbf{H}_2(\mathbb{A}_F)$ lifts to $\{L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))\}$.

For a representation π of $\mathrm{GL}(2, \mathbb{A}_E)$, let ${}^\sigma\pi$ denote the representation

$${}^\sigma\pi : (g_{ij}) \mapsto \pi(\sigma g_{ij}), \quad \forall (g_{ij}) \in \mathrm{GL}(2, \mathbb{A}_E).$$

1.5. PROPOSITION*. Let π be an irreducible discrete spectrum automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$, μ a character of C_F , such that $\pi \neq {}^\sigma\pi$ and $\omega_\pi = \mu \circ N_{E/F}$.

- If π is cuspidal, the representation $\pi \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ lifts to a stable packet $\{\Pi\}$ which in turn lifts to the cuspidal automorphic representation $\pi(\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.
- Suppose $\pi = \chi 1_{\mathrm{GL}(2, \mathbb{A}_E)}$, where χ is a character of C_E such that $\chi \neq {}^\sigma\chi$ and ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ for some nontrivial quadratic character ε' of C_F . Let ζ be either ε' or $\varepsilon\varepsilon'$. Then, $\pi \otimes_2 \chi|_{\mathbb{A}_F^\times} \cdot \zeta$ lifts to a stable quasi-packet $\{\Pi\}$ which in turn lifts to the Langlands quotient representation $J(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. We denote $\{\Pi\}$ by $\{L(\nu\varepsilon\zeta, \nu^{-1/2}\pi(\chi))\}$.

In either case, no discrete spectrum packet of $\mathbf{H}_1(\mathbb{A}_F)$ lifts to the (quasi-)packet $\{\Pi\}$.

Additional global lifting results involving parabolically induced representations of $\mathbf{G}(\mathbb{A}_F)$ are stated in Section 4.5.

1.6. PROPOSITION*. Suppose $\{\Pi\}$ is a global packet of $\mathrm{GSp}(2, \mathbb{A}_F)$ which is the lift of some automorphic representation/packet of one of the ε -endoscopic groups. Then, $\{\Pi\}$ contains at least one representation which is ε -invariant.

If $\{\Pi\}$ is a discrete spectrum global packet of $\mathrm{GSp}(2, \mathbb{A}_F)$ which is not the lift of any discrete spectrum representation/packet of the ε -endoscopic groups, then $\{\Pi\}$ does not contain any discrete spectrum automorphic representation which is ε -invariant.

1.4.3. Local Lifting. Let k be a local nonarchimedean field. Let ε be a nontrivial quadratic character of $k^\times$. Let $\mathcal{K}$ be the quadratic extension of k which corresponds via local class field theory to ε . Let $G = \mathrm{GSp}(2, k)$. Let $Z = \{\mathrm{diag}(z, z, z, z) : z \in k^\times\}$ be the center of G . Let

- (1) $H_1 = (\mathrm{GL}(2, k) \times \mathcal{K}^\times)' = \{(g, x) \in \mathrm{GL}(2, k) \times \mathcal{K}^\times : \det g = N_{\mathcal{K}/k}x\},$
- (2) $H_2 = (\mathrm{GL}(2, \mathcal{K}) \times k^\times) / \{(\mathrm{diag}(z, z), N_{\mathcal{K}/k}z^{-1}) : z \in \mathcal{K}^\times\}.$

Recall that the character ε defines a character of G via $\varepsilon(g) := \varepsilon(\lambda(g)), \forall g \in G$. For any representation π of G , we say that π is ε -invariant if $\pi \cong \varepsilon\pi$.

Fix a character ω of Z . Recall that $C(G, \omega)$ is the space of smooth functions on G which are compactly supported modulo Z and transform under Z via ω^{-1} . We fix once and for all a Haar measure dg on G . For any irreducible ε -invariant admissible representation π of G with central character ω , and for any function $f \in C(G, \omega)$, recall that

$$\langle \pi, f \rangle_A := \mathrm{tr} \int_{Z \backslash G} \pi(g) f(g) A dg,$$

where A is an intertwining operator in $\mathrm{Hom}_G(\pi, \varepsilon\pi)$ such that $A^2 = 1$.

For $i = 1, 2$, let $Z_0(H_i)$ denote the maximal k -split component of the center of H_i . We shall see in Section 2.4 that $Z_0(H_i)$ is isomorphic to Z ; thus, ω is defines a character on $Z_0(H_i)$. Let $C(H_i, \omega)$ denote the space of smooth functions on H_i which are compactly supported modulo $Z_0(H_i)$ and transform under $Z_0(H_i)$ via ω^{-1} . Fix once and for all a Haar measure dh_i on H_i . For any admissible representation π_i of H_i , and for any function $f_i \in C(H_i, \omega)$, put

$$\langle \pi_i, f_i \rangle := \mathrm{tr} \int_{Z_0(H_i) \backslash H_i} \pi_i(h_i) f_i(h_i) dh_i.$$

Using global lifting results, we deduce in Chapter 5 local character identities relating quantities of the form $\langle \pi, f \rangle_A$ and $\langle \pi_i, f_i \rangle$ for **matching** functions f, f_i . Functions are matching if their orbital integrals are compatible with the norm correspondence between G and its

ε -endoscopic groups. We shall discuss norm correspondence in Section 2.4. In general, the existence of matching functions is conjectural. It is related to a conjecture called the Fundamental Lemma. We prove the Fundamental Lemma in the context of G and its ε -endoscopic groups in Appendix B.

1.4.3.1. *Notations:* For a character θ of $\mathcal{K}^\times$, let $\pi(\theta)$ denote the cuspidal monomial representation of $\mathrm{GL}(2, k)$ associated with θ , or more precisely with the representation $\mathrm{Ind}_{W_{\mathcal{K}/\mathcal{K}}}^{W_{\mathcal{K}/k}}(\theta)$ of W_k . If θ is invariant under the action of $\mathrm{Gal}(\mathcal{K}/k)$, $\pi(\theta)$ is induced; otherwise, $\pi(\theta)$ is cuspidal ([JL]).

Let ν be the normalized absolute value function on k ; that is, $\nu(x) = q^{-\mathrm{ord} x}$, where q is the cardinality of the residue field of k and $\mathrm{ord} x$ is the p -adic valuation of x .

Let P_0 be the minimal upper triangular parabolic subgroup of G . It has a decomposition $P_0 = TN$, where T is the maximal diagonal torus of G and N is the unipotent component of P_0 . For characters μ_1, μ_2, μ of $k^\times$, let $\mu_1 \times \mu_2 \rtimes \mu$ denote the representation of $\mathrm{GL}(2, k)$ (normalizedly) parabolically induced from the following representation of P_0 :

$$\mu_1 \otimes \mu_2 \otimes \mu : \begin{pmatrix} a & & \\ & b & \\ & \lambda/b & \\ & & \lambda/a \end{pmatrix} n \mapsto \mu_1(a)\mu_2(b)\mu(\lambda), \quad \forall a, b, \lambda \in k^\times, n \in N.$$

Let P_β be the standard Heisenberg parabolic subgroup of G containing P_0 . Its Levi component is

$$M_\beta = \left\{ \begin{pmatrix} a & & \\ & g & \\ & & \frac{\det g}{a} \end{pmatrix} : a \in k^\times, g \in \mathrm{GL}(2, k) \right\}.$$

Let N_β be the unipotent component of P_β . It is a Heisenberg group. For a character μ of $k^\times$ and an admissible representation π of $\mathrm{GL}(2, k)$, let $\mu \rtimes \pi$ denote the representation of G (normalizedly) parabolically induced from the following representation of P_β :

$$\mu \otimes \pi : \begin{pmatrix} a & & \\ & g & \\ & & \frac{\det g}{a} \end{pmatrix} n \mapsto \mu(a)\pi(g), \quad \forall a \in k^\times, g \in \mathrm{GL}(2, k), n \in N_\beta.$$

1.4.3.2. *Summary of Local Character Identities.* Below is a summary of our twisted character identities for the ε -invariant representations of $G = \mathrm{GSp}(2, k)$.

1. Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial, admissible $\mathrm{GL}(2, k)$ -module whose central character ω_τ is trivial. Let $\{\pi^+, \pi^-\}$ be the local packet of G which lifts to the induced representation $I_{(2,2)}(\tau, \varepsilon\tau)$ of $\mathrm{PGL}(4, k)$ according to [F4]. There exist intertwining operators A^+ in $\mathrm{Hom}_G(\pi^+, \varepsilon\pi^+)$, and A^- in $\mathrm{Hom}_G(\pi^-, \varepsilon\pi^-)$, such that the following holds for matching functions:

$$\begin{aligned} \langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} &= \langle \tau \otimes_1 1, f_1 \rangle, \\ \langle \pi^+, f \rangle_{A^+} - \langle \pi^-, f \rangle_{A^-} &= \langle B_{\mathcal{K}/k}\tau \otimes_2 \omega_\tau, f_2 \rangle. \end{aligned}$$

2. Let θ, χ be characters of $\mathcal{K}^\times$ such that none of $\theta, \chi, \theta\chi, \theta^\sigma\chi$ is fixed by the action of $\mathrm{Gal}(\mathcal{K}/k) = \langle \sigma \rangle$. Suppose $(\theta\chi)|_{k^\times} = (\theta^\sigma\chi)|_{k^\times} = \varepsilon$, then the distinct $\mathcal{K}$ -monomial representations $\pi(\theta\chi), \pi(\theta^\sigma\chi)$ are cuspidal with trivial central characters.

Let $\{\pi^+, \pi^-\}$ be the local packet which lifts to the induced $\mathrm{PGL}(4, k)$ -module

$$I_{(2,2)}(\pi(\theta\chi), \pi(\theta^\sigma\chi)).$$

There exist intertwining operators A^+ in $\text{Hom}_G(\pi^+, \varepsilon\pi^+)$, A^- in $\text{Hom}_G(\pi^-, \varepsilon\pi^-)$, and a constant $\epsilon = \pm 1$, such that the following holds for matching functions:

$$\begin{aligned}\langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} &= \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle, \\ \langle \pi^+, f \rangle_{A^+} - \langle \pi^-, f \rangle_{A^-} &= \epsilon \langle \pi(\chi) \otimes_1 \theta, f_1 \rangle.\end{aligned}$$

3. Let χ be a character of $\mathcal{K}^\times$ not fixed by $\text{Gal}(\mathcal{K}/k)$. Let π^+, π^- be the two inequivalent, tempered, irreducible subrepresentations of $1 \rtimes \pi(\chi)$ ([ST]). There exist intertwining operators A^+ in $\text{Hom}_G(\pi^+, \varepsilon\pi^+)$ and A^- in $\text{Hom}_G(\pi^-, \varepsilon\pi^-)$, such that the following holds for matching functions:

$$\langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} = \langle \pi(\chi) \otimes_1 1, f_1 \rangle.$$

4. Let $1_{2,l}$ ($l = k, \mathcal{K}$) denote the nontempered trivial representation of $\text{GL}(2, l)$. Let $\text{St}_{2,l}$ denote the square integrable Steinberg representation of $\text{GL}(2, l)$.

Suppose ε is unramified. Let ξ be a (possibly trivial) quadratic character of $k^\times$. Let $\delta = \delta(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ be the unique square integrable constituent of the induced G -module $\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\xi$. Let $\delta^- = \delta^-(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ be the cuspidal member of the local packet containing δ . In particular, the local packet $\{\delta, \delta^-\}$ lifts to $I_{(2,2)}(\xi\text{St}_{2,k}, \varepsilon\xi\text{St}_{2,k})$ of $\text{PGL}(4, k)$.

Let $L = L(\nu\varepsilon, \varepsilon \rtimes \nu^{-1/2}\xi)$ be the nontempered quotient of $\nu^{1/2}\varepsilon 1_{2,k} \rtimes \nu^{-1/2}\xi$ ([ST]). The local quasi-packet $\{L, \delta^-\}$ lifts to $I_{(2,2)}(\xi 1_{2,k}, \varepsilon\xi 1_{2,k})$ of $\text{PGL}(4, k)$.

There exist intertwining operators $A^L \in \text{Hom}_G(L, \varepsilon L)$, $A^\delta \in \text{Hom}_G(\delta, \varepsilon\delta)$, and $A^- \in \text{Hom}_G(\delta^-, \varepsilon\delta^-)$, such that the following holds for matching functions:

$$\begin{aligned}\langle L, f \rangle_{A^L} + \langle \delta^-, f \rangle_{A^-} &= \langle \xi 1_{2,k} \otimes_1 1, f_1 \rangle, \\ \langle L, f \rangle_{A^L} - \langle \delta^-, f \rangle_{A^-} &= \langle (\xi \circ \text{N}_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes_2 1, f_2 \rangle;\end{aligned}$$

$$\begin{aligned}\langle \delta, f \rangle_{A^\delta} - \langle \delta^-, f \rangle_{A^-} &= \langle \xi \text{St}_{2,k} \otimes_1 1, f_1 \rangle, \\ \langle \delta, f \rangle_{A^\delta} + \langle \delta^-, f \rangle_{A^-} &= \langle (\xi \circ \text{N}_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes_2 1, f_2 \rangle.\end{aligned}$$

5. Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial, admissible representation of the group $\text{GL}(2, k)$, χ a character of $\mathcal{K}^\times$, such that: There does not exist a character ε' of $k^\times$ for which ${}^\sigma\chi/\chi = \varepsilon' \circ \text{N}_{\mathcal{K}/k}$ and $\tau \cong \varepsilon'\tau$. Then, there exists a collection $[\pi]$ of distinct ε -invariant, irreducible, cuspidal, admissible G -modules, and an intertwining operator $A(\pi') \in \text{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes_1 \chi, f_1 \rangle.$$

REMARK. From the proof of the above twisted character identity, one should expect that $[\pi]$ is a subset of a local packet which lifts to $\pi(\chi B_{\mathcal{K}/k}\tau)$, the cuspidal $\text{GL}(4, k)$ -module which is the twisted endoscopic lift of the representation $\chi B_{\mathcal{K}/k}\tau$ of $\text{GL}(2, \mathcal{K})$ (see [AC]).

6. Let χ be a character of $\mathcal{K}^\times$ which is not fixed by $\text{Gal}(\mathcal{K}/k)$. Let $L = L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$ be the nontempered quotient of the parabolically induced G -module $\nu\varepsilon \rtimes \nu^{-1/2}\pi(\chi)$. Let $\delta = \delta(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$ be the square integrable subrepresentation of $\nu\varepsilon \rtimes \nu^{-1/2}\pi(\chi)$. There exist intertwining operators A^L in $\text{Hom}_G(L, \varepsilon L)$, and A^δ in $\text{Hom}_G(\delta, \varepsilon\delta)$ such that

the following holds for matching functions:

$$\begin{aligned}\langle L, f \rangle_{A^L} &= \langle 1_{2,k} \otimes_1 \chi, f_1 \rangle, \\ \langle \delta, f \rangle_{A^\delta} &= \langle \text{St}_{2,k} \otimes_1 \chi, f_1 \rangle.\end{aligned}$$

7. Let τ be an irreducible, cuspidal, admissible representation of $\text{GL}(2, \mathcal{K})$, not fixed by the action of $\text{Gal}(\mathcal{K}/k)$, such that its central character ω_τ is equal to $\mu \circ \mathbf{N}_{\mathcal{K}/k}$ for some character μ of $k^\times$. Then, there exists a collection $[\pi]$ of distinct ε -invariant, irreducible, cuspidal, admissible G -modules, and an intertwining operator $A(\pi')$ in $\text{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes_2 \mu, f_2 \rangle.$$

REMARK. From the proof of the above twisted character identity, one should expect that $[\pi]$ is a subset of a local packet which lifts to $\pi(\tau)$, the cuspidal $\text{GL}(4, k)$ -module which is the twisted endoscopic lift of τ .

8. Let τ be a cuspidal non- $\mathcal{K}$ -monomial, or one dimensional, irreducible, admissible representation of $\text{GL}(2, k)$. There exists an operator A , intertwining the induced G -module $\varepsilon \rtimes \tau$ with $\varepsilon \rtimes \varepsilon\tau$, such that

$$\langle \varepsilon \rtimes \tau, f \rangle_A = \langle B_{\mathcal{K}/k}\tau \otimes_2 \omega_\tau \varepsilon, f_2 \rangle$$

for matching functions.

9. Suppose the field extension $\mathcal{K}/k$ is unramified. Let χ be a character of $\mathcal{K}^\times$ such that $\chi \neq {}^\sigma\chi$ and ${}^\sigma\chi/\chi = \varepsilon' \circ \mathbf{N}_{\mathcal{K}/k}$ for some nontrivial quadratic character ε' of $k^\times$. Let $\zeta = \varepsilon'$ or $\varepsilon'\varepsilon$. Let $L = L(\varepsilon\zeta\nu, \nu^{-1/2}\pi(\chi))$ and $\delta = \delta(\varepsilon\zeta\nu, \nu^{-1/2}\pi(\chi))$. There exist intertwining operators A^L in $\text{Hom}_G(L, \varepsilon L)$, and A^δ in $\text{Hom}_G(\delta, \varepsilon\delta)$, such that the following holds for matching functions:

$$\begin{aligned}\langle L, f \rangle_{A^L} &= \langle \chi 1_{2,\mathcal{K}} \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \rangle, \\ \langle \delta, f \rangle_{A^\delta} &= \langle \chi \text{St}_{2,\mathcal{K}} \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \rangle.\end{aligned}$$

Additional character identities involving parabolically induced G -modules are stated in Sections 5.5.1 and 5.5.2.

1.5. Assumptions and Restrictions

Before we continue further, we list here the assumptions on which this work rests and the restrictions on our results:

Assumptions.

- (1) The multiplicity one theorem for the discrete spectrum of $\text{GSp}(2)$. As mentioned earlier in this chapter, the multiplicity one theorem has been established for $\text{PGSp}(2)$ by Flicker in [F4].
- (2) The results of Arthur's announced in [A3], including the multiplicity one theorem for the automorphic representations of $\text{GSp}(2)$ which are associated with semisimple Arthur parameters.
- (3) The existence of the transfer of functions from $\text{GSp}(2)$ to its ε -twisted endoscopic groups, and the precise forms which matching functions take (see remarks after the proofs of Theorem 4.10 and Corollary 4.16). In the nontwisted

case, Waldspurger has shown that the Fundamental Lemma implies the transfer of functions [Wa1]. In all likelihood, the twisted analogue of Waldspurger's theorem also holds.

Restrictions. We are only able to classify those discrete spectrum, ε -invariant, automorphic representations of $\mathrm{GSp}(2)$ which have at least two elliptic local components. The restriction “two elliptic components” could presumably be reduced to “one elliptic component,” through a technique developed in [F2] which involves a class of test functions called regular functions. It also seems likely that the work of Arthur's announced in [A3] contains a method to do away with any restriction on the automorphic representations.

CHAPTER 2

ε -Endoscopy for $\mathrm{GSp}(2)$

2.1. Endoscopic Data

In this chapter, F is either a local field of characteristic zero or a number field. If F is a number field, let C_F denote the idèle class group of F . For a reductive F -group $\mathbf{H}$, let $\hat{H}$ denote the identity component of the L -group ${}^L H$ of $\mathbf{H}$ (see [Bo]). Let $\mathbf{G}$ be the reductive F -group $\mathrm{GSp}(2)$.

Kottwitz and Shelstad have defined in [KS] the endoscopic data attached to a triple $(\mathbf{G}, \theta, \mathbf{a})$, where θ is an automorphism of $\mathbf{G}$ over F . We now recall this definition in the special case where θ is trivial.

2.1. DEFINITION. The quadruple $(\mathbf{H}, \mathcal{H}, s, \xi)$ is a set of ε -**endoscopic data** (or in the terminology of [KS], the endoscopic data attached to the triple $(\mathbf{G}, 1, \mathbf{a})$) if it satisfies each of the following conditions:

- (1) $\mathbf{H}$ is a quasi-split reductive group over F .
- (2) $\mathcal{H}$ is a split extension of W_F by $\hat{H}$ such that the L -action $\rho_{\mathcal{H}}$ of W_F on $\hat{H}$ determined by this extension coincides with the Weil group action determined by ${}^L H$ (see [Bo]).
- (3) s is a semisimple element in $\hat{G}$.
- (4) $\xi : \mathcal{H} \rightarrow {}^L G$ is an L -homomorphism satisfying the following two conditions.
 - (a) $\mathrm{Int}(s) \circ \xi = a \cdot \xi$;
 - (b) ξ maps $\hat{H}$ isomorphically onto the identity component of $\mathrm{Cent}(s, \hat{G})$, the centralizer of s in $\hat{G}$.

We call $\mathbf{H}$ an ε -**endoscopic group** of $\mathbf{G}$.

REMARK. The action ρ of W_F on $\hat{H}$ determined by $\mathcal{H}$ may not fix any choice of splitting for $\hat{H}$. However, there exists a set $\{h_w \in \hat{H}\}_{w \in W_F}$ such that the action $\rho_{\mathcal{H}} : w \mapsto \mathrm{Int}(h_w)\rho(w)$ does fix a choice of splitting for $\hat{H}$. As such, $\rho_{\mathcal{H}}$ constitutes an L -action. For our purpose, $\mathcal{H}$ may be taken to be ${}^L H$; hence, condition 2 is largely moot.

2.1.1. Elliptic ε -Endoscopic Data of $\mathbf{G}$. In general, an L -group is a semidirect product $\hat{H} \rtimes W_F$. As we shall show, every group considered in this work is split over some quadratic extension of F . For simplicity, we let ${}^L H$ denote the finite Galois form $\hat{H} \rtimes \mathrm{Gal}(K/F)$ of the L -group, where K is the smallest extension of F over which $\mathbf{H}$ splits. In particular, since $\mathbf{G}$ is F -split, we let ${}^L G$ denote $\hat{G}$, which we identify with $\mathrm{GSp}(2, \mathbb{C})$ (see [Bo], [A3]).

Let Γ be the absolute Galois group $\mathrm{Gal}(\bar{F}/F)$. Since W_F is dense in Γ , the action of W_F on any group $\hat{H}$ extends to an action of Γ on $\hat{H}$. A set of endoscopic data $(H, \mathcal{H}, s, \xi)$ is said to be **elliptic** if $\xi(Z(\hat{H})^\Gamma)^0$ is contained in $Z(\hat{G})$ (Here, upper 0 denotes identity

component). We may take Γ to be $\mathrm{Gal}(K/F)$ if $\mathbf{H}$ splits over some finite extension K of F .

Two sets of ε -endoscopic data $(H, \mathcal{H}, s, \xi)$ and $(H', \mathcal{H}', s', \xi')$ are said to be **equivalent** if there exists $g \in \hat{G}$ such that:

- (1) $g\xi(\mathcal{H})g^{-1} = \xi'(\mathcal{H}')$,
- (2) $gs g^{-1} \equiv s' \pmod{Z(\hat{G})}$.

With regard to the application of Kottwitz-Shelstad's stabilization of the trace formula, we will compute elliptic endoscopic data only up to equivalence (hence the earlier remark that we may take $\mathcal{H}$ to be ${}^L H$).

Let ε be a nontrivial quadratic character of $F^\times$ if F is local, or of C_F if F is global. Let W_F denote the Weil group of F (see [T]). As explained in [KS], corresponding via Langlands correspondence to ε is a cocycle $\mathbf{a}$ in $H^1(W_F, Z(\hat{G}))$ if F is local, or in $H^1(W_F, Z(\hat{G}))/\ker^1(W_F, Z(\hat{G}))$ if F is global.

Let E be the quadratic extension of F which corresponds to ε via local or global class field theory (see [La]). Then, $\mathbf{a}|_{W_E}$ is trivial, and we identify $\mathbf{a}$ with a representative in $H^1(\mathrm{Gal}(E/F), Z(\hat{G}))$.

Since $\mathbf{G}$ is F -split, the Galois action of $\mathrm{Gal}(E/F)$ on $Z(\hat{G}) = \mathbb{C}^\times$ is trivial. Let σ be the generator of $\mathrm{Gal}(E/F)$. The cohomology class of $\mathbf{a}$ consists of a single character $a \in \mathrm{Hom}(\mathrm{Gal}(E/F), \mathbb{C}^\times)$, defined by $a(\sigma) = -1$.

Suppose $(\mathbf{H}, \mathcal{H}, s, \xi)$ is a set of ε -endoscopic data. Then, for any $h_w \in \mathcal{H}$ with image w in W_F , we have:

$$\xi(h_w)^{-1} s \xi(h_w) = \begin{cases} -s & \text{if } w \equiv \sigma \pmod{W_E}, \\ s & \text{otherwise.} \end{cases}$$

Since we shall list the sets of endoscopic data only up to equivalence, we may assume that the semisimple element s lies in the diagonal torus $\hat{T}$ of $\hat{G}$. Since ${}^L G = \hat{G}$, if $\xi(h_w)^{-1} s \xi(h_w) = -s$ for some $h_w \in \mathcal{H}$, then $\mathrm{Int}(\xi(h_w))$ must correspond to an element of the Weyl group $W = W(\hat{T}, \hat{G})$ of $\hat{T}$ in $\hat{G}$.

Given any $t = \mathrm{diag}(a, b, \lambda/b, \lambda/a)$ in $\hat{T}$, let the numbers 1, 2, 3, 4 represent the entries $a, b, \lambda/b, \lambda/a$, respectively. Then, the actions of W on t are represented by the following set of permutations on $\{1, 2, 3, 4\}$:

$$\{1, (12)(34), (23), (3421), (2431), (42)(31), (23)(41), (14)\}.$$

Hence, W is isomorphic to D_4 , the dihedral group of order 8.

By going through all possible images of s under W , we see that, up to equivalence of endoscopic data, s must be equal to one of:

- (1) $s_1 = \mathrm{diag}(1, 1, -1, -1)$,
- (2) $s_2 = \mathrm{diag}(1, -1, -1, 1)$,
- (3) $s_3 = \mathrm{diag}(1, -1, -d, d)$ for some $d \neq \pm 1$.

2.2. CLAIM. *No elliptic ε -endoscopic data is of the form $(\mathbf{H}, \mathcal{H}, s_3, \xi)$.*

PROOF. Suppose $(\mathbf{H}, \mathcal{H}, s_3, \xi)$ constitutes a set of endoscopic data. Then, $\hat{H}$ is isomorphic to

$$\mathrm{Cent}(s, \hat{G})^0 = \hat{T} = \{\mathrm{diag}(a, b, \lambda/b, \lambda/a) : a, b, \lambda \in \mathbb{C}^\times\}.$$

By condition 4a in Definition 2.1, if $w \in W_F$ is equivalent to σ modulo W_E , then the action of w on $\hat{H}$ must correspond to the permutation $(12)(34)$. If w is not equivalent to σ

modulo W_E , then the action of w on $\hat{H}$ is trivial. Consequently,

$$Z(\hat{H})^\Gamma = \hat{H}^\Gamma = \{\text{diag}(a, a, \lambda/a, \lambda/a) : a, \lambda \in \mathbb{C}^\times\},$$

which is connected and not contained in $Z(\hat{G})$, which means $(H, \mathcal{H}, s_3, \xi)$ is not elliptic. $\square$

In the following sections, we consider the ε -endoscopic data $(\mathbf{H}, \mathcal{H}, s, \xi)$ where s is equal to s_1 or s_2 . It turns out that there is only one quasi-split $\mathbf{H}$ such that $(\mathbf{H}, \mathcal{H}, s, \xi)$ is elliptic for each choice of s .

2.2. Endoscopic group $\mathbf{H}_1$

2.2.1. The data $(\mathbf{H}_1, \mathcal{H}_1, s_1, \xi_1)$. We now construct a set of elliptic ε -endoscopic data $(\mathbf{H}_1, \mathcal{H}_1, s_1, \xi_1)$, where

$$s_1 = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}.$$

Let

$$\hat{H}_1 = (\text{GL}(2, \mathbb{C}) \times \mathbb{C}^\times \times \mathbb{C}^\times) / \{(\text{diag}(z, z)^{-1}, z, z) : z \in \mathbb{C}^\times\}.$$

Fix a splitting $\text{spl}_{\hat{H}_1} = \{\hat{B}_1, \hat{T}_1, \{X_1\}\}$ of $\hat{H}_1$, where $\hat{B}_1 = \{((\begin{smallmatrix} * & * \\ * & * \end{smallmatrix}), *, *)\}$ is a Borel subgroup of $\hat{H}_1$, $\hat{T}_1$ is the diagonal torus of $\hat{H}_1$, and $\{X_1\}$ is a singleton consisting of the image of the vector $X_1 = ((\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}), 0, 0) \in \mathfrak{gl}(2, \mathbb{C}) \times \mathbb{C} \times \mathbb{C}$ in the Lie algebra of $\hat{H}_1$. Here, lower case, gothic type denotes the Lie algebra of a Lie group.

Let σ be the generator of $\text{Gal}(E/F)$. Define an action of W_F on $\hat{H}_1$ by

$$w(A, x, y) = \begin{cases} (A, y, x) & \text{if } w \equiv \sigma \pmod{W_E}, \\ (A, x, y) & \text{otherwise.} \end{cases}$$

This defines a semi-direct product $\mathcal{H}_1 = \hat{H}_1 \rtimes W_F$. Since the action of W_F fixes $\text{spl}_{\hat{H}_1}$, $\mathcal{H}_1$ is an L -group with $\hat{H}_1$ as its identity component.

Let $e = (\begin{smallmatrix} 1 & \\ & -1 \end{smallmatrix})$. For any $A \in \text{GL}(2, \mathbb{C})$, $x, y \in \mathbb{C}^\times$, let $d(A, x, y)$ be the element $(\begin{smallmatrix} xA & \\ & yeAe \end{smallmatrix})$ in $\hat{G}$. Then, d induces an isomorphism

$$d : \hat{H}_1 \xrightarrow{\sim} \text{Cent}(s_1, \hat{G})^0.$$

Define an L -embedding $\xi_1 : \mathcal{H}_1 \rightarrow \hat{G}$ by $\xi_1|_{\hat{H}_1} = d$ and

$$\xi_1(1 \rtimes w) = \begin{cases} (\begin{smallmatrix} e & \\ & e \end{smallmatrix}) & \text{if } w \equiv \sigma \pmod{W_E}, \\ 1 & \text{otherwise.} \end{cases}$$

Let $\mathbf{H}_1$ be the unique quasi-split reductive group over F whose L -group ${}^L H_1$ is equal to $\mathcal{H}_1$. One can check to see that $(\mathbf{H}_1, \mathcal{H}_1, s_1, \xi_1)$ is a set of elliptic ε -endoscopic data.

2.3. CLAIM. *Up to equivalence, $(\mathbf{H}_1, \mathcal{H}_1, s_1, \xi_1)$ is the unique elliptic ε -endoscopic data attached to s_1 .*

PROOF. If $(\mathbf{H}, \mathcal{H}, s_1, \xi)$ is another set of elliptic ε -endoscopic data attached to s_1 , then $\xi(\mathcal{H}) = \xi_1(\mathcal{H}_1)$. It follows from the definitions that $(\mathbf{H}, \mathcal{H}, s_1, \xi)$ is equivalent to $(\mathbf{H}_1, \mathcal{H}_1, s_1, \xi_1)$. $\square$

2.2.2. Explicit Description of $\mathbf{H}_1$. We now describe explicitly the quasi-split reductive F -group $\mathbf{H}_1$. Let σ be the generator of $\mathrm{Gal}(E/F)$. Recall that $\mathcal{H}_1 = \hat{H}_1 \rtimes W_F$, and the action of W_F factors through $W_F/W_E = \mathrm{Gal}(E/F)$. Consequently, $\mathbf{H}_1$ must split over E .

To construct $\mathbf{H}_1$, we make use of the fact that there exists a maximal torus $\mathbf{T}_1$ of $\mathbf{H}_1$ dual to the maximal diagonal torus $\hat{T}_1$ of $\hat{H}_1$; namely, $X_*(\mathbf{T}_1) = X^*(\hat{T}_1)$, where $X_*(\mathbf{T}_1) := \mathrm{Hom}(\mathbb{G}_m, \mathbf{T}_1)$ and $X^*(\hat{T}_1) := \mathrm{Hom}(\hat{T}_1, \mathbb{C}^\times)$. We have:

$$X^*(\hat{T}_1) = \{(x, y; z, t) \in \mathbb{Z}^2 \times \mathbb{Z}^2 : x + y = z + t\},$$

where $(x, y; z, t)(\mathrm{diag}(a, b), c, d) := a^x b^y c^z d^t$ for all $(\mathrm{diag}(a, b), c, d) \in \hat{T}_1$.

The Galois action of σ on $\hat{T}_1$ is given by $\sigma(\mathrm{diag}(a, b), c, d) = (\mathrm{diag}(a, b), d, c)$. This induces an action on $X^*(\hat{T}_1)$ and consequently on $X_*(\mathbf{T}_1)$, namely:

$$\sigma(x, y; z, t) = (x, y; t, z), \quad \forall (x, y; z, t) \in X_*(\mathbf{T}_1).$$

We conclude that

$$\mathbf{T}_1 = (\mathbb{G}_m \times \mathbb{G}_m \times \mathrm{R}_{E/F}\mathbb{G}_m)',$$

where $\mathrm{R}_{E/F}\mathbb{G}_m$ is the F -group obtained from $\mathbb{G}_m$ on restricting scalars from E to F , and the prime indicates that the product of the first two factors is equal to the norm of the third factor in $\mathbb{G}_m$.

The algebraic group $\mathbf{H}_1$ is defined by its root datum, which is the dual of the root datum of $\hat{H}_1$ (see [Sp]). Having found the dual of $\hat{T}_1$, it remains to find the dual of the root lattice and make a choice of splitting. We skip this routine procedure. The resulting group is

$$\mathbf{H}_1 = (\mathrm{GL}(2) \times \mathrm{R}_{E/F}\mathbb{G}_m)',$$

where the prime indicates that the determinant of the first factor is equal to the norm of the second factor in $\mathbb{G}_m$.

Since $\mathbf{H}_1$ is split over E and the group of E -points of $\mathrm{R}_{E/F}\mathbb{G}_m$ is $E^\times \times E^\times$, the group of E -points of $\mathbf{H}_1$ is $\mathbf{H}_1(E) =$

$$(\mathrm{GL}(2, E) \times E^\times \times E^\times)' := \{(g, \gamma, \delta) \in \mathrm{GL}(2, E) \times E^\times \times E^\times : \det g = \gamma\delta\}.$$

For any $\gamma \in E$, put $\bar{\gamma} := \sigma\gamma$. For $g = (g_{ij}) \in \mathrm{GL}(2, E)$, put $\bar{g} := (\bar{g}_{ij})$. The action of σ on $\mathbf{H}_1(E)$ is induced by the action of σ on $\hat{H}_1$, namely:

$$\sigma^* : (g, \gamma, \delta) \mapsto (\bar{g}, \bar{\delta}, \bar{\gamma}), \quad \forall (g, \gamma, \delta) \in (\mathrm{GL}(2, E) \times E^\times \times E^\times)'.$$

The F -points of $\mathbf{H}_1$ are the elements in $\mathbf{H}_1(E)$ which are fixed by σ^* ; thus,

$$\mathbf{H}_1(F) = (\mathrm{GL}(2, F) \times E^\times)' := \boxed{\{(g, \gamma) \in \mathrm{GL}(2, F) \times E^\times : \det g = \mathrm{N}_{E/F}\gamma\}}.$$

Suppose F is a number field. The norm mapping $\mathrm{N}_{E/F} : E \rightarrow F$ induces a norm mapping $\mathrm{N}_{E/F} : \mathbb{A}_E \rightarrow \mathbb{A}_F$ on the adèles, and the group of $\mathbb{A}_F$ -points of $\mathbf{H}_1$ is

$$\mathbf{H}_1(\mathbb{A}_F) = (\mathrm{GL}(2, \mathbb{A}_F) \times \mathbb{A}_E^\times)' := \boxed{\{(g, \gamma) \in \mathrm{GL}(2, \mathbb{A}_F) \times \mathbb{A}_E^\times : \det g = \mathrm{N}_{E/F}\gamma\}}.$$

At each place v of F , the Galois action on $\mathbf{H}_1$ as an F_v -group is compatible with the Galois action on $\mathbf{H}_1$ as an F -group. At a place v which does not split in E , $E_v := E \otimes_F F_v$ is a field and $\mathrm{Gal}(E_v/F_v)$ embeds isomorphically into $\mathrm{Gal}(E/F)$. Consequently, the F_v -group $\mathbf{H}_1$ splits over E_v and its group of F_v -points is as follows:

$$\mathbf{H}_1(F_v) = (\mathrm{GL}(2, F_v) \times E_v^\times)' := \{(g, \gamma) \in \mathrm{GL}(2, F_v) \times E_v^\times : \det g = \mathrm{N}_{E/F}\gamma\}.$$

If v splits into two places v_1, v_2 in E , then $E_{v_1} = E_{v_2} = F_v$ and the Galois action on $\mathbf{H}_1$ as an F_v -group is trivial, i.e. $\mathbf{H}_1$ is split over F_v . We have $\mathbf{H}_1(F_v) =$

$$(\mathrm{GL}(2, F_v) \times F_v^\times \times F_v^\times)' := \{(g, a, b) \in \mathrm{GL}(2, F_v) \times F_v^\times \times F_v^\times : \det g = ab\}.$$

2.3. Endoscopic group $\mathbf{H}_2$

2.3.1. The Data $(\mathbf{H}_2, \mathcal{H}_2, s_2, \xi_2)$. We now construct a set of elliptic ε -endoscopic data $(\mathbf{H}_2, \mathcal{H}_2, s_2, \xi_2)$, where

$$s_2 = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}.$$

Let

$$\hat{H}_2 = \{(A, B, \lambda) \in \mathrm{GL}(2, \mathbb{C})^2 \times \mathbb{C}^\times : \det A = \det B = \lambda\}.$$

Fix a splitting $\mathbf{spl}_{\hat{H}_2} = \{\hat{B}_2, \hat{T}_2, \{X_a, X_b\}\}$ of $\hat{H}_2$, where

$$\hat{B}_2 = \{((\begin{smallmatrix} * & * \\ * & * \end{smallmatrix}), (\begin{smallmatrix} * & * \\ * & * \end{smallmatrix}), *)\} \subset \hat{H}_2,$$

$\hat{T}_2$ is the maximal diagonal torus of $\hat{H}_2$, and

$$X_a = ((\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}), (\begin{smallmatrix} 0 & 0 \\ 0 & 0 \end{smallmatrix}), 0), \quad X_b = ((\begin{smallmatrix} 0 & 0 \\ 0 & 0 \end{smallmatrix}), (\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}), 0)$$

are elements in the Lie algebra of $\hat{H}_2$.

Define an action of W_F on $\hat{H}_2$ by

$$w(A, B, \lambda) = \begin{cases} (B, A, \lambda) & \text{if } w \equiv \sigma \pmod{W_E}, \\ (A, B, \lambda) & \text{otherwise.} \end{cases}$$

This defines a semi-direct product $\mathcal{H}_2 = \hat{H}_2 \rtimes W_F$. Since the action of W_F fixes $\mathbf{spl}_{\hat{H}_2}$, $\mathcal{H}_2$ is an L -group, with $\hat{H}_2$ as its identity component.

We define an L -group embedding $\xi_2 : \mathcal{H}_2 \rightarrow \hat{G}$ as follows: Put

$$\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} e & f \\ g & h \end{pmatrix} \right] := \begin{pmatrix} a & e & f & b \\ & c & g & h \\ & & & d \end{pmatrix}.$$

Define the restriction of ξ_2 to $\hat{H}_2$ by

$$\xi_2|_{\hat{H}_2}(A, B, \lambda) = [A, B], \quad \forall (A, B, \lambda) \in \hat{H}_2.$$

Then, $\hat{H}_2$ is isomorphic to $\mathrm{Cent}(s_2, \hat{G})^0$ via $\xi_2|_{\hat{H}_2}$. Put $w_0 := (\begin{smallmatrix} & 1 \\ 1 & \end{smallmatrix}) \in \mathrm{GL}(2, \mathbb{C})$. Let

$$\xi_2(1 \rtimes w) = \begin{cases} (\begin{smallmatrix} w_0 & \\ & w_0 \end{smallmatrix}) & \text{if } w \equiv \sigma \pmod{W_E}, \\ 1 & \text{otherwise.} \end{cases}$$

Let $\mathbf{H}_2$ be the unique quasi-split reductive group over F whose L -group ${}^L H_2$ is $\mathcal{H}_2$. One can check to see that $(\mathbf{H}_2, \mathcal{H}_2, s_2, \xi_2)$ is a set of elliptic ε -endoscopic data.

2.4. CLAIM. *Up to equivalence, $(\mathbf{H}_2, \mathcal{H}_2, s_2, \xi_2)$ is the unique elliptic ε -endoscopic data attached to s_2 .*

PROOF. As in the case of s_1 , if $(\mathbf{H}, \mathcal{H}, s_2, \xi)$ is another set of elliptic ε -endoscopic data attached to s_2 , then $\xi(\mathcal{H}) = \xi_2(\mathcal{H}_2)$. It follows from the definitions that $(\mathbf{H}, \mathcal{H}, s_2, \xi)$ is equivalent to $(\mathbf{H}_2, \mathcal{H}_2, s_2, \xi_2)$. $\square$

2.3.2. Explicit Description of $\mathbf{H}_2$. We now give an explicit description of the quasi-split, reductive F -group $\mathbf{H}_2$. The action of W_F on $\hat{H}_2$ factors through $W_F/W_E = \mathrm{Gal}(E/F)$. Consequently, the group $\mathbf{H}_2$ is split over the quadratic extension E .

To find $\mathbf{H}_2$, let $\mathbf{T}_2$ be the maximal torus of $\mathbf{H}_2$ which is dual to the maximal diagonal torus $\hat{T}_2$ of $\hat{H}_2$. Thus, $X_*(\mathbf{T}_2)$ is equal to

$$X^*(\hat{T}_2) = \{(x, y; z, t; w) \in \mathbb{Z}^2 \times \mathbb{Z}^2 \times \mathbb{Z}\} / \{(n, n; m, m; -n - m) : m, n \in \mathbb{Z}\},$$

where

$$(x, y; z, t; w)(\mathrm{diag}(a, b), \mathrm{diag}(c, d), \lambda) = a^x b^y c^z d^t \lambda^w, \\ \forall (\mathrm{diag}(a, b), \mathrm{diag}(c, d), \lambda) \in \hat{T}_2.$$

The action of σ on $\hat{T}_2$ induces an action on $X^*(\hat{T}_2)$; namely,

$$\sigma(x, y; z, t; w) = (z, t; x, y; w), \quad \forall (x, y; z, t; w) \in X^*(\hat{T}_2).$$

We conclude that $\mathbf{T}_2$ is a torus whose group of E -points is

$$\left\{ \left(\begin{pmatrix} \zeta_1 & \\ & \zeta_2 \end{pmatrix}, \begin{pmatrix} \gamma_1 & \\ & \gamma_2 \end{pmatrix}, \eta \right) : \zeta_1, \zeta_2, \gamma_1, \gamma_2, \eta \in E^\times \right\} / \{(\alpha I_2, \beta I_2, (\alpha\beta)^{-1}) : \alpha, \beta \in E^\times\},$$

where I_2 is the 2×2 identity matrix. Put $\bar{\gamma} := \sigma\gamma$ for $\gamma \in E$. The action of σ on $\hat{T}_2$ gives rise to an action of σ on $\mathbf{T}_2(E)$ define by:

$$\sigma^*(\mathrm{diag}(\zeta_1, \zeta_2), \mathrm{diag}(\gamma_1, \gamma_2), \eta) = (\mathrm{diag}(\bar{\gamma}_1, \bar{\gamma}_2), \mathrm{diag}(\bar{\zeta}_1, \bar{\zeta}_2), \eta).$$

Having found the the dual of $\hat{T}_2$, we conclude without going through the details here that:

$$\mathbf{H}_2(E) = (\mathrm{GL}(2, E) \times \mathrm{GL}(2, E) \times E^\times) / \{(\alpha I_2, \beta I_2, (\alpha\beta)^{-1}) : \alpha, \beta \in E^\times\}.$$

For $g = (g_{ij}) \in \mathrm{GL}(2, E)$, let $\bar{g} = (\overline{g_{ij}})$. The Galois action on $\hat{H}_2$ induces an action on $\mathbf{H}_2(E)$ defined by:

$$\sigma^* : (g_1, g_2, \zeta) \mapsto (\bar{g}_2, \bar{g}_1, \bar{\zeta}), \quad \forall (g_1, g_2, \zeta) \in \mathbf{H}_2(E).$$

2.5. CLAIM. *The group of F -points of $\mathbf{H}_2$ is*

$$\mathbf{H}_2(F) = \{(g, \bar{g}, c) : g \in \mathrm{GL}(2, E), c \in F^\times\} / \{(z I_2, \bar{z} I_2, N_{E/F} z^{-1}) : z \in E^\times\} \\ \cong \boxed{(\mathrm{GL}(2, E) \times F^\times) / \{(z I_2, N_{E/F} z^{-1}) : z \in E^\times\}}.$$

PROOF. The group of F -points of $\mathbf{H}_2$ consists of the set of equivalence classes represented by $(g, h, \zeta) \in \mathrm{GL}(2, E) \times \mathrm{GL}(2, E) \times F^\times$ such that $(g, h, \zeta) \equiv (\bar{h}, \bar{g}, \bar{\zeta})$ modulo $\{(\alpha I_2, \beta I_2, (\alpha\beta)^{-1}) : \alpha, \beta \in E^\times\}$.

Suppose $(g, h, \zeta) \equiv (\bar{h}, \bar{g}, \bar{\zeta})$, then there exist $\alpha, \beta \in E^\times$ such that

$$\alpha g = \bar{h}, \\ \beta h = \bar{g}, \\ (\alpha\beta)^{-1} \zeta = \bar{\zeta}.$$

This implies that $\alpha = \bar{\beta}^{-1}$ and $\beta^{-1} \zeta \in F^\times$. Since

$$(g, h, \zeta) \equiv (g, \bar{g}, \beta^{-1} \zeta) = (g, h, \zeta)(1, \beta, \beta^{-1}) \mod \{(\alpha I_2, \beta I_2, (\alpha\beta)^{-1})\},$$

the claim follows. $\square$

Suppose F is a number field. The norm mapping $N_{E/F} : E \rightarrow F$ induces a mapping $N_{E/F} : \mathbb{A}_E \rightarrow \mathbb{A}_F$. The group of $\mathbb{A}_F$ -points of $\mathbf{H}_2$ is:

$$\mathbf{H}_2(\mathbb{A}_F) = (\mathrm{GL}(2, \mathbb{A}_E) \times \mathbb{A}_F^\times) / \{(zI_2, N_{E/F}z^{-1}) : z \in \mathbb{A}_E^\times\}.$$

At a place v of F which does not split in E , the Galois group $\mathrm{Gal}(E_v/F_v)$ is isomorphic to $\mathrm{Gal}(E/F)$. Consequently, we have:

$$\mathbf{H}_2(F_v) = (\mathrm{GL}(2, E_v) \times F_v^\times) / \{(zI_2, N_{E/F}z^{-1}) : z \in E_v^\times\}.$$

At a place v which splits in E , $\mathbf{H}_2$ is split over F_v , and $\mathbf{H}_2(F_v)$ is equal to

$$\{(g_1, g_2, c) \in \mathrm{GL}(2, F_v) \times \mathrm{GL}(2, F_v) \times F_v^\times\} / \{(aI_2, bI_2, (ab^{-1})) : a, b \in F_v^\times\}.$$

2.3.2.1. *Summary.* In summary, we work with the the following quasi-split reductive F -groups:

- $\mathbf{G} = \mathrm{GSp}(2) := \{g \in \mathrm{GL}(4) : {}^t g J g = \lambda(g) J \text{ for some } \lambda(g) \in \mathbb{G}_m\}$, where $J = \begin{pmatrix} & & & 1 \\ & & 1 & \\ & -1 & & \\ -1 & & & \end{pmatrix}$.
- $\mathbf{H}_1 = (\mathrm{GL}(2) \times \mathrm{R}_{E/F} \mathbb{G}_m)'$, whose group of F -points is:

$$\mathbf{H}_1(F) = \{(g, \gamma) \in \mathrm{GL}(2, F) \times E^\times : \det g = N_{E/F} \gamma\}.$$

- The reductive group $\mathbf{H}_2$, whose group of F -points is:

$$\mathbf{H}_2(F) = (\mathrm{GL}(2, E) \times F^\times) / \{(zI_2, N_{E/F}z^{-1}) : z \in E^\times\}.$$

2.4. Norm Correspondence

Let F be a local field or a number field.

Terminology:

- For any algebraic F -group $\mathbf{H}$, we say that two elements h, h' in $\mathbf{H}(F)$ are **conjugate** (or F -conjugate) if there exists an element $g \in \mathbf{H}(F)$ such that $g^{-1}hg = h'$. A conjugacy class in $\mathbf{H}(F)$ consists of all elements in $\mathbf{H}(F)$ which are conjugate to one another.
- We say that $h, h' \in \mathbf{H}(F)$ are **stably conjugate** (or $\bar{F}$ -conjugate) if there exists an element $g \in \mathbf{H}(\bar{F})$ such that $g^{-1}hg = h'$. A **stable conjugacy class** in $\mathbf{H}(F)$ consists of all elements in $\mathbf{H}(F)$ which are stably conjugate to a given one. We say that a conjugacy class is stable if it coincides with the stable conjugacy class which contains it.
- For our purpose, an element in $\mathbf{H}(F)$ is **semisimple** if it is diagonalizable over $\bar{F}$. A semisimple element is **regular** if its centralizer in $\mathbf{H}(F)$ is a maximal torus.
- A torus in $\mathbf{H}(F)$ is **elliptic** if it is not contained in any proper parabolic subgroup of $\mathbf{H}(F)$. An element in $\mathbf{H}(F)$ is **elliptic** if it is contained in an elliptic torus.

The goal of this work is to relate admissible representations of $\mathbf{G}(F)$, $\mathbf{H}_1(F)$, and $\mathbf{H}_2(F)$ if F is local, or automorphic representations of $\mathbf{G}(\mathbb{A}_F)$, $\mathbf{H}_1(\mathbb{A}_F)$, and $\mathbf{H}_2(\mathbb{A}_F)$ if F is a number field. To do so, we need to introduce the notion of transfer of stable conjugacy classes from $\mathbf{H}_1$, $\mathbf{H}_2$ to $\mathbf{G}$.

Let $\mathbf{H}$ be an ε -endoscopic group of $\mathbf{G}$ with L -group embedding $\xi : {}^L H \rightarrow {}^L G$. Let $\mathrm{CL}_{\mathrm{ss}}(\mathbf{H})$, $\mathrm{CL}_{\mathrm{ss}}(\mathbf{G})$ denote the set of stable conjugacy classes of semisimple elements in $\mathbf{H}(F)$, $\mathbf{G}(F)$, respectively.

Let $\mathbf{T} = \{\mathrm{diag}(a, b, l/b, l/a) : a, b, l \in \mathbb{G}_m\}$ be the maximal diagonal torus of $\mathbf{G}$. Let $\mathbf{T}_{\mathbf{H}}$ be a maximal torus of $\mathbf{H}$ defined over F . Let λ be the unique map from $X^*(\mathbf{T}_{\mathbf{H}})$ to $X_*(\hat{T})$ such that the following diagram commutes:

$$\begin{array}{ccc} X^*(\mathbf{T}_{\mathbf{H}}) & \xrightarrow{\lambda} & X^*(\mathbf{T}) \\ \downarrow & & \downarrow \\ X_*(\hat{T}_H) & \xrightarrow{\xi} & X_*(\hat{T}) \end{array}$$

Here, the vertical isomorphisms are given by the correspondence between root data and dual root data (see [Bo]).

Any semisimple $\delta_H \in \mathbf{H}(F)$ is $\bar{F}$ -conjugate to some $t_H \in \mathbf{T}_{\mathbf{H}}(\bar{F})$. Likewise, any semisimple $\delta \in \mathbf{G}(F)$ is $\bar{F}$ -conjugate to some $t \in \mathbf{T}(\bar{F})$. Let $\tilde{\delta}_H \in \mathrm{CL}_{\mathrm{ss}}(\mathbf{H})$ denote the stable conjugacy class of $\delta_H \in \mathbf{H}(F)$. Let $\tilde{\delta}$ denote the stable conjugacy class of $\delta \in \mathbf{G}(F)$.

Define a map $\mathcal{A}_{\mathbf{H}/\mathbf{G}} : \mathrm{CL}_{\mathrm{ss}}(\mathbf{H}) \rightarrow \mathrm{CL}_{\mathrm{ss}}(\mathbf{G})$ as follows: $\mathcal{A}_{\mathbf{H}/\mathbf{G}}(\tilde{\delta}_H) = \tilde{\delta}$ if there exists $t_H \in \mathbf{T}_{\mathbf{H}}(\bar{F})$, $t \in \mathbf{T}(\bar{F})$ such that

- t_H is conjugate to δ_H in $\mathbf{H}(\bar{F})$, t is conjugate to δ in $\mathbf{G}(\bar{F})$;
- $\chi(t_H) = (\lambda(\chi))(t)$ for all $\chi \in X^*(\mathbf{T}_{\mathbf{H}})$.

2.6. DEFINITION. We say that $\delta_H \in \mathbf{H}(F)$ is a **norm** of $\delta \in \mathbf{G}(F)$ if $\tilde{\delta} = \mathcal{A}_{\mathbf{H}/\mathbf{G}}(\tilde{\delta}_H)$. If F is global, we say that $\delta_H = (\delta_{H,v}) \in \mathbf{H}(\mathbb{A}_F)$ is a norm of $\delta = (\delta_v) \in \mathbf{G}(\mathbb{A}_F)$ if $\delta_{H,v} \in \mathbf{H}(F_v)$ is a norm of $\delta_v \in \mathbf{G}(F_v)$ for every place v of F .

In general, $\mathcal{A}_{\mathbf{H}/\mathbf{G}}$ may not be one-to-one. Moreover, a regular element in $\mathbf{H}$ may be a norm of a non-regular element in $\mathbf{G}$. If $\delta_H \in \mathbf{H}$ is a norm of a regular element in $\mathbf{G}$, we say that δ_H is **$\mathbf{G}$ -regular**. (Remark: A $\mathbf{G}$ -regular element in $\mathbf{H}$ is necessarily regular).

Note that $\mathcal{A}_{\mathbf{H}/\mathbf{G}}$ is equivalent to a map

$$\mathbf{N} : \mathbf{T}_{\mathbf{H}}(\bar{F})/W_H \rightarrow \mathbf{T}(\bar{F})/W,$$

where W, W_H are the Weyl groups associated with $\mathbf{T}(\bar{F}), \mathbf{T}_{\mathbf{H}}(\bar{F})$, respectively. More precisely, given any class $\tilde{t}_H \in \mathbf{T}_{\mathbf{H}}(\bar{F})/W_H$ represented by $t_H \in \mathbf{T}_{\mathbf{H}}(\bar{F})$, the class $\mathbf{N}(\tilde{t}_H)$ is represented by some $t \in \mathbf{T}(\bar{F})$ such that $(\lambda(\chi))(t) = \chi(t_H)$ for all $\chi \in X^*(\mathbf{T}_{\mathbf{H}})$. We shall describe $\mathbf{N}$ explicitly in the next two sections.

In this work, we identify $\hat{G}$ with $\mathrm{GSp}(2, \mathbb{C})$ (see [Bo]). Write:

$$X^*(\mathbf{T}) = \{(x, y, z, t) \in \mathbb{Z}^4\} / \{(m, -m, -m, m) : m \in \mathbb{Z}\},$$

where $(x, y, z, t)\delta := a^x b^y (l/b)^z (l/a)^t$ for all $\delta = \mathrm{diag}(a, b, l/b, l/a) \in \mathbf{T}$. Write:

$$X_*(\hat{T}) = \{(x, y, z, t) \in \mathbb{Z}^4 : x + t = y + z\},$$

where $(x, y, z, t)\zeta := \mathrm{diag}(\zeta^x, \zeta^y, \zeta^z, \zeta^t) \in \mathbf{T}$ for all $\zeta \in \mathbb{G}_m$.

The isomorphism $\iota : X^*(\mathbf{T}) \rightarrow X_*(\hat{T})$, determined from the identification of $\hat{G}$ with $\mathrm{GSp}(2, \mathbb{C})$, is defined as follows:

$$\iota : (x, y, z, t) \mapsto (x + y, x + z, t + y, t + z),$$

$$\iota^{-1} : (x, y, z, t) \mapsto (x - z, z, t, 0).$$

For $i = 1, 2$, let $\mathbf{T}_i$ denote the maximal diagonal torus of $\mathbf{H}_i$. Let W_i be the Weyl group of $\mathbf{T}_i(\bar{F})$ in $\mathbf{H}_i(\bar{F})$. Let $\mathbf{N}_i$ denote the map $\mathbf{N}$ from $\mathbf{T}_i(\bar{F})/W_i$ to $\mathbf{T}(\bar{F})/W$ defined above.

2.4.1. Norm Correspondence for $\mathbf{H}_1$.

Recall that

$$\hat{H}_1 = (\mathrm{GL}(2, \mathbb{C}) \times \mathbb{C}^\times \times \mathbb{C}^\times) / \{(\mathrm{diag}(z, z), z^{-1}, z^{-1}) : z \in \mathbb{C}^\times\}.$$

We have chosen the embedding $\xi_1 : \hat{H}_1 \rightarrow \hat{G}$ to be $\xi_1 : (A, u, v) \mapsto ({}^u A_{veAe})$, where $e := \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$. Write:

$$X^*(\mathbf{T}_1) = \{(x_1, y_1; z_1, t_1) \in \mathbb{Z}^4\} / \{(z, z, -z, -z) : z \in \mathbb{Z}\},$$

where

$$(x_1, y_1; z_1, t_1)(\mathrm{diag}(a, b), c, d) := a^{x_1} b^{y_1} c^{z_1} d^{t_1}, \quad \forall (\mathrm{diag}(a, b), c, d) \in \mathbf{T}_1.$$

2.7. CLAIM. The map $N_1 : \mathbf{T}_1(\bar{F})/W_1 \rightarrow \mathbf{T}(\bar{F})/W$ is given by

$$N_1 : (\mathrm{diag}(a, l/a); b, l/b) \mapsto \mathrm{diag}(b, a, l/a, l/b).$$

PROOF. The lattice $X^*(\mathbf{T}_1)$ is isomorphic to $X_*(\hat{T}_1)$ in the following way: For any $(x_1, y_1; z_1, t_1) \in X^*(\mathbf{T}_1)$, its image in $X_*(\hat{T}_1)$ is the homomorphism which maps any $\zeta \in \mathbb{C}^\times$ to $\mathrm{diag}((\zeta^{x_1}, \zeta^{y_1}), \zeta^{z_1}, \zeta^{t_1})$. The embedding of $X_*(\hat{T}_1)$ in $X_*(\hat{T})$ is given by

$$\xi_1 : (x_1, y_1; z_1, t_1) \mapsto (z_1 + x_1, z_1 + y_1, t_1 + x_1, t_1 + y_1).$$

The map $\lambda : X^*(\mathbf{T}_1) \rightarrow X^*(\mathbf{T})$ is therefore given by

$$\lambda : (x_1, y_1; z_1, t_1) \mapsto \iota^{-1} \circ \xi_1(x_1, y_1; z_1, t_1) = (z_1 - t_1, t_1 + x_1, t_1 + y_1, 0)$$

Suppose an element $t_1 = (\mathrm{diag}(a_1, l_1/a_1); b_1, l_1/b_1)$ in $\mathbf{T}_1(\bar{F})$ is a norm of an element $t = (a, b, l/b, l/a)$ in $\mathbf{T}(\bar{F})$. By definition, this means that $\chi(t_1) = (\lambda(\chi))(t')$ for some t' conjugate to t and for all $\chi \in X^*(\mathbf{T}_1)$. Without loss of generality, assume $t' = t$. We have:

$$a_1^{x_1} (l_1/a_1)^{y_1} b_1^{z_1} (l_1/b_1)^{t_1} = a_1^{x_1 - y_1} b_1^{z_1 + t_1} (l/b)^{t_1 + y_1} = a^{z_1 - t_1} b^{x_1 - y_1} (l/b)^{t_1 + y_1}$$

for all $(x_1, y_1, z_1, t_1) \in X^*(\mathbf{T}_1)$; hence, $a = b_1, b = a_1, l = l_1$, which completes the proof. $\square$

2.4.2. Norm Correspondence for $\mathbf{H}_2$. To compute the norm correspondence between $\mathbf{G}$ and $\mathbf{H}_2$, we carry out the same procedure employed for $\mathbf{H}_1$. Let I_2 be the 2×2 identity matrix. Write:

$$\mathbf{H}_2(\bar{F}) = (\mathrm{GL}(2, \bar{F}) \times \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times) / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in \bar{F}^\times\}.$$

A typical element in $\mathbf{T}_2(\bar{F})$ is represented by

$$(\mathrm{diag}(a, b), \mathrm{diag}(c, d), 1) \in \mathrm{GL}(2, \bar{F}) \times \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times.$$

2.8. CLAIM. The map $N_2 : \mathbf{T}_2(\bar{F})/W_2 \rightarrow \mathbf{T}(\bar{F})/W$ is given by

$$N_2 : (\mathrm{diag}(a, b), \mathrm{diag}(c, d), 1) \mapsto \mathrm{diag}(ac, ad, bc, bd).$$

PROOF. We have:

$$X^*(\mathbf{T}_2) = \{(x_2, y_2; z_2, t_2) \in \mathbb{Z}^4 : x_2 + y_2 = z_2 + t_2\},$$

where

$$(x_2, y_2; z_2, t_2)(\mathrm{diag}(a, b), \mathrm{diag}(c, d), 1) := a^{x_2} b^{y_2} c^{z_2} d^{t_2},$$

$$\forall (\mathrm{diag}(a, b), \mathrm{diag}(c, d), 1) \in \mathbf{T}_2.$$

The embedding of $X_*(\hat{T}_2) = X^*(\mathbf{T}_2)$ in $X_*(\hat{T})$ is given by

$$\xi_2 : (x_2, y_2; z_2, t_2) \mapsto (x_2, z_2, t_2, y_2).$$

Thus, $\lambda : X^*(\mathbf{T}_2) \rightarrow X^*(\mathbf{T})$ is given by

$$\lambda : (x_2, y_2; z_2, t_2) \mapsto \iota^{-1}(x_2, z_2, t_2, y_2) = (x_2 - t_2, t_2, y_2, 0).$$

Suppose an element $t_2 = (\mathrm{diag}(a_2, b_2), \mathrm{diag}(c_2, d_2), 1)$ in $\mathbf{T}_2(\bar{F})$ is a norm of some $t = (a, b, l/b, l/a) \in \mathbf{T}(\bar{F})$. Then by definition we must have $\chi(t_2) = (\lambda(\chi))(t')$ for some t' conjugate to t and for all $\chi \in X^*(\mathbf{T}_2)$. Without loss of generality, assume that $t' = t$. We have:

$$a_2^{x_2} b_2^{y_2} c_2^{z_2} d_2^{t_2} = (a_2 c_2)^{x_2} (b_2 c_2)^{y_2} (d_2 / c_2)^{t_2} = a^{x_2 - t_2} b^{t_2} (l/b)^{y_2} = a^{x_2} (b/a)^{t_2} (l/b)^{y_2}$$

for arbitrary $(x_2, y_2, z_2, t_2; w) \in X^*(\mathbf{T}_2)$ (The first equality is valid because of the condition that $x_2 + y_2 = z_2 + t_2$). We conclude that:

$$a = a_2 c_2, \quad l/b = b_2 c_2, \quad b/a = d_2 / c_2;$$

hence, $t = \mathrm{diag}(a_2 c_2, a_2 d_2, b_2 c_2, b_2 d_2)$. $\square$

2.4.3. Norm Correspondence for F -Points. Let $G = \mathbf{G}(F)$, $H_i = \mathbf{H}_i(F)$ ($i = 1, 2$). We have described the norm correspondence among the $\bar{F}$ -points of the groups. We now describe the norm correspondence among non-elliptic or central elements in G , H_1 , H_2 . Knowing the norm correspondence for these types of elements is a prerequisite to stating certain useful character identities in Chapter 5. The norm correspondence among elliptic regular elements is addressed in Appendix B.

Recall that E is the quadratic extension of F which corresponds to ε . Suppose $E = F(\sqrt{A})$ for some element $A \in F^\times - F^{\times 2}$. For any element $\gamma = a + b\sqrt{A}$ in $E^\times$, where $a, b \in F$, put

$$\phi^A(\gamma) := \begin{pmatrix} a & bA \\ b & a \end{pmatrix} \in \mathrm{GL}(2, F).$$

The (not necessarily distinct) eigenvalues of $\phi^A(\gamma)$ are $\gamma, \sigma\gamma$, where σ is the generator of $\mathrm{Gal}(E/F)$. Observe that $\det \phi^A(\gamma) = a^2 - b^2 A = N_{E/F} \gamma$, and ϕ^A defines an embedding of $E^\times$ onto a maximal elliptic torus in $\mathrm{GL}(2, F)$. Moreover, $\phi^A(\sigma\gamma)$ is equal to $\begin{pmatrix} 1 & \\ & -1 \end{pmatrix}^{-1} \phi^A(\gamma) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$.

2.9. CLAIM. *Let $\delta_1 = (\mathrm{diag}(a, b), \gamma)$ be an element in H_1 . Then, δ_1 is a norm of $\begin{pmatrix} a & \\ & \phi^A(\gamma) \end{pmatrix} \in G$.*

PROOF. The image of δ_1 in $\mathbf{T}_1(\bar{F})$ is $(\mathrm{diag}(a, b), \gamma, \sigma\gamma)$. The rest follows from Claim 2.7. $\square$

Note that since norm correspondence is defined up to stable conjugacy, the elements $(\mathrm{diag}(a, b), \gamma)$ and $(\mathrm{diag}(a, b), \sigma\gamma)$ in H_1 are both norms of the same element in G .

2.10. CLAIM. *Let $(\mathrm{diag}(\alpha, \beta), c)$ be an element in $\mathrm{GL}(2, E) \times F^\times$, and δ_2 its image in H_2 . Then, δ_2 is a norm of $\begin{pmatrix} cN_{E/F}\alpha & \\ & c\phi^A(\alpha\sigma\beta) \\ & & cN_{E/F}\beta \end{pmatrix} \in G$.*

PROOF. The image of δ_2 in $\mathbf{T}_2(\bar{F})$ is represented by

$$((\begin{smallmatrix} \alpha & \\ & \beta \end{smallmatrix}), c(\begin{smallmatrix} \sigma\alpha & \\ & \sigma\beta \end{smallmatrix}), 1) \in \mathrm{GL}(2, \bar{F}) \times \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times.$$

The rest follows from Claim 2.8. $\square$

Note that the elements $(\mathrm{diag}(\alpha, \beta), c)$ and $(\mathrm{diag}(\sigma\alpha, \sigma\beta), c)$ in H_2 are both norms of the same element in G .

2.4.3.1. *Central Elements.* The center of G is $Z = \{\text{diag}(z, z, z, z) : z \in F^\times\}$. The maximal F -split component of the center of H_1 is

$$Z_0(H_1) = \{(\text{diag}(z, z), z) : z \in F^\times\}.$$

The maximal F -split component of the center of H_2 is

$$Z_0(H_2) = \{(\text{diag}(1, 1), z)_* : z \in F^\times\},$$

where lower $*$ denotes the equivalence class of the element modulo the group

$$\{(\text{diag}(x, x), N_{E/F}x^{-1}) : x \in E^\times\}.$$

2.11. COROLLARY. *The elements $(\text{diag}(z, z), z)$ in $Z_0(H_1)$ and $(\text{diag}(1, 1), z)_*$ in $Z_0(H_2)$ are norms of $\text{diag}(z, z, z, z) \in Z$.*

2.4.3.2. *Split Case.* Suppose F is a number field. Let v be a place of F where $\varepsilon_v = 1$. Then, $\mathbf{H}_1(F_v) = \{(g, a, b) \in \text{GL}(2, F_v) \times F_v^\times \times F_v^\times : \det g = ab\}$, and $\mathbf{H}_2(F_v)$ is

$$\{(g_1, g_2, c) \in \text{GL}(2, F_v) \times \text{GL}(2, F_v) \times F_v^\times\} / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in F_v^\times\}.$$

2.12. CLAIM. *The element $\delta_1 = (\text{diag}(c, d), a, b) \in \mathbf{H}_1(F_v)$ is a norm of the element $\text{diag}(c, a, b, d) \in \mathbf{G}(F_v)$.*

PROOF. This follows from Claim 2.7. $\square$

2.13. CLAIM. *The element $\delta_2 = (\text{diag}(a, b), \text{diag}(c, d), 1)_* \in \mathbf{H}_2(F_v)$ is a norm of $\text{diag}(ac, ad, bc, bd) \in \mathbf{G}(F_v)$.*

PROOF. This follows from Claim 2.8. $\square$

2.5. Matching Functions

Suppose F is a local field. Let $G = \mathbf{G}(F)$. Let ε be a quadratic character of $F^\times$. Fix a character ω of the center of G . For any regular element $t \in G$, let $Z_G(t)$ denote the centralizer of t in G . In particular, since t is regular, $Z_G(t)$ is a maximal torus. Let $C(G, \omega)$ denote the space of functions on G which are smooth, compactly supported modulo center, and transform under the center of G via ω^{-1} . For $f \in C(G, \omega)$, put

$$O_G(f, t) := \int_{Z_G(t) \backslash G} f(g^{-1}tg) \varepsilon(g) dg.$$

Let $\mathbf{H}$ be either $\mathbf{H}_1$ or $\mathbf{H}_2$. Let $H = \mathbf{H}(F)$. The character ω defines a character of $Z_0(H) \cong Z$. Let $C(H, \omega)$ denote the space of functions on H which are smooth, compactly supported modulo $Z_0(H)$, and transform under $Z_0(H)$ via ω^{-1} . For any $\mathbf{G}$ -regular element $t_H \in H$ and function $f_H \in C(H, \omega)$, put

$$O_H(f_H, t_H) := \int_{Z_H(t_H) \backslash H} f_H(h^{-1}t_H h) dh$$

and

$$SO_H(f_H, t_H) := \sum_{t'_H} O_H(f_H, t'_H),$$

where the sum is taken over a set of representatives for the conjugacy classes of elements $t'_H \in H$ in the stable conjugacy class of t_H .

Let H' denote the subgroup of $\mathbf{G}$ -regular elements in H . A conjecture of Langlands asserts that there exists a function $\Delta_{G/H}$ on $H' \times G$ with the following property: For every $f \in C(G, \omega)$, there exists a function $f_H \in C(H, \omega)$ such that

$$(2.1) \quad SO_H(f_H, t_H) = \sum_t \Delta_{G/H}(t_H, t) O_G(f, t), \quad \forall t_H \in H',$$

where the sum is over a set of representatives for the F -conjugacy classes of regular elements $t \in G$. We say that f, f_H are **matching functions** if the above identity holds. The function $\Delta_{G/H}$, called the **transfer factor**, is defined in [LS] in the standard case (where $\varepsilon = 1$) and in [KS] in a more general twisted context. In certain special cases, one may find a description of it in [H], [H1]. The transfer factor is expected to satisfy a host of properties (see [KS]). A full discussion of the transfer factor is beyond the scope of this work, but we do specify it explicitly in the unramified case, which is needed for the Fundamental Lemma (see below). We list here some properties of $\Delta_{G/H}$ which are of interest to us:

- $\Delta_{G/H}(t_H, t) = 0$ if t_H is not a norm of t .
- For all $g \in G$, $\Delta(t_H, g^{-1}tg) = \varepsilon(g)\Delta(t_H, t)$.

Note that, without the second property, the right hand side of (2.1) is not well defined.

Suppose now that F is a number field. Let ε be a quadratic character of C_F . Let $\mathbf{H}$ be an elliptic ε -endoscopic group of $\mathbf{G}$ over F . Let $G = \mathbf{G}(F)$ and $G_v = \mathbf{G}(F_v)$ for any place v of F . Define H, H_v likewise. Suppose $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$ and $f_H \in C(\mathbf{H}(\mathbb{A}_F), \omega)$ are products of local functions $f_v, f_{H,v}$, respectively. We say that f and f_H are **matching functions** if f_v and $f_{H,v}$ are matching for all $v \in V$.

2.5.1. The Fundamental Lemma. An ingredient in the conjecture on the existence of matching functions is the “Fundamental Lemma”, which is actually a conjecture. Let F be a local field. Let ε be a quadratic character of $F^\times$. Let $\mathbf{H}$ be an elliptic ε -endoscopic group of $\mathbf{G}$ over F . Let $\mathcal{O}$ be the ring of integers in F . Assume that $\mathbf{H}$ is defined over $\mathcal{O}$. Let $K = \mathbf{G}(\mathcal{O})$, $K_H = \mathbf{H}(\mathcal{O})$. Let $\mathcal{H}(G)$ be the Hecke algebra of smooth, compactly supported, K -biinvariant functions on G . Define $\mathcal{H}(H)$ likewise, with K replaced with K_H . The L -group embedding $\xi : {}^L H \rightarrow \hat{G}$ induces a map

$$b_\xi : \mathcal{H}(G) \rightarrow \mathcal{H}(H).$$

The Fundamental Lemma asserts that f and $b_\xi(f)$ are matching functions for any f in $\mathcal{H}(G)$.

Suppose F is a number field. Hales shows in [H2] that if one can show that the unit elements of $\mathcal{H}(G_v)$ and $\mathcal{H}(H_v)$ have matching orbital integrals for almost all finite places v , then the Fundamental Lemma follows. Note that even though the title of [H2] reads “standard endoscopy,” the paper does in fact cover our situation. Using this result of Hales’ and the results of Flicker’s in [F3], we prove in Appendix B the Fundamental Lemma in the context of $\mathbf{G}$ and its ε -endoscopic groups.

The Trace Formula

3.1. The fine χ -expansion

3.1.1. An Overview. Let F be a number field. Let V be the set of places of F . For any finite place v , let $\mathcal{O}_v$ be the ring of integers of F_v . Let $\mathbf{H}$ be a reductive F -group. Let $\mathbf{Z}_0$ be the maximal F -split component of the center $\mathbf{Z}$ of $\mathbf{H}$. Let ε be a character of $\mathbf{H}(\mathbb{A}_F)$ whose restriction to $\mathbf{Z}(\mathbb{A}_F)$ is trivial. From now on, by a group we mean an F -group.

Fix a character ω of $\mathbf{Z}_0(F) \backslash \mathbf{Z}_0(\mathbb{A}_F)$. For any place v of F , let $H_v = \mathbf{H}(F_v)$. Let $C(H_v, \omega_v)$ be the space of smooth functions f_v on H_v such that f_v is compactly supported modulo $Z_{0,v}$ and $f_v(zh) = \omega_v^{-1}(z)f_v(h)$ for all $z \in Z_{0,v}, h \in H_v$.

At a finite place v where $\mathbf{H}$ is defined over $\mathcal{O}_v$, let K_v denote the hyperspecial, maximal compact subgroup $\mathbf{H}(\mathcal{O}_v)$ of H_v . Let $\mathcal{H}(H_v, \omega_v)$ denote the Hecke algebra of K_v -biinvariant functions in $C(H_v, \omega_v)$. If v is archimedean, fix a maximal compact subgroup of H_v , and let $\mathcal{H}(H_v, \omega_v)$ be the set of K_v -finite functions in $C(H_v, \omega_v)$.

Let $C(\mathbf{H}(\mathbb{A}_F), \omega)$ denote the span of the smooth, compactly supported functions on $\mathbf{H}(\mathbb{A}_F)$ which are of the form $\otimes_v f_v$, where $f_v \in C(H_v, \omega_v)$ for all v and f_v is a unit in the Hecke algebra $\mathcal{H}(H_v, \omega_v)$ for almost all finite v . In this work, whenever we mention a function $f \in C(\mathbf{H}(\mathbb{A}_F), \omega)$, we assume that f is the product of local components f_v . Since such functions span $C(\mathbf{H}(\mathbb{A}_F), \omega)$, our assumption has no ill effect on our attempt to understand the spectrum of $\mathbf{H}(\mathbb{A}_F)$.

Fix a minimal (F -)parabolic subgroup $\mathbf{P}_0$ of $\mathbf{H}$. Let $\mathbf{A}_0$ be the maximal F -split component of the (F -)Levi subgroup of $\mathbf{P}_0$. Let $W(\mathbf{A}_0, H)$ be the Weyl group of $\mathbf{A}_0$ in $\mathbf{H}$. For any Levi subgroup $\mathbf{M}$ of $\mathbf{H}$, let $\mathbf{A}_M$ denote the split component of the center of $\mathbf{M}$. Let $X_*(\mathbf{A}_M) = \text{Hom}(\mathbb{G}_m, \mathbf{A}_M)$. Let $\mathfrak{a}_M$ denote $X_*(\mathbf{A}_M) \otimes_{\mathbb{Z}} \mathbb{R}$. Let $\mathfrak{a}_M^*$ denote the dual of $\mathfrak{a}_M$. Let $\mathcal{P}(M)$ denote the set of all (F -)parabolic subgroups of $\mathbf{H}$ containing $\mathbf{M}$.

We fix a maximal compact subgroup $\mathbb{K} = \otimes_{v \in V} K_v$ in $\mathbf{H}(\mathbb{A}_F)$, where $K_v = \mathbf{H}(\mathcal{O}_v)$ for every finite place v such that $\mathbf{H}$ is defined over $\mathcal{O}_v$, and K_v is a fixed maximal compact subgroup in $\mathbf{H}(F_v)$ for the rest of the places v .

Let $\mathbf{M}$ be a Levi subgroup in $\mathbf{H}$. For any $\zeta \in \mathfrak{a}_M^*$, let χ_ζ denote the character of $\mathbf{M}(\mathbb{A}_F)$ associated with ζ . For any $m \in \mathbf{M}(\mathbb{A}_F)$, let $H_P(m)$ be the element in $\mathfrak{a}_M$ uniquely defined by the condition: $e^{\langle \zeta, H_P(m) \rangle} = \chi_\zeta(m)$ for all $\zeta \in \mathfrak{a}_M^*$.

For any parabolic subgroup $\mathbf{P} \in \mathcal{P}(M)$, representation τ of $\mathbf{M}(\mathbb{A}_F)$, and element $\zeta \in \mathfrak{a}_M^*$, let $I_{P,\tau}(\zeta)$ denote the $\mathbf{H}(\mathbb{A}_F)$ -module normalizedly induced from the $\mathbf{M}(\mathbb{A}_F)$ -module which sends an element m in $\mathbf{M}(\mathbb{A}_F)$ to $\tau(m)e^{\langle \zeta, H_P(m) \rangle}$. In other words, $I_{P,\tau}(\zeta)$ is the right regular action on the space of smooth functions ϕ on $\mathbf{H}(\mathbb{A}_F)$ which satisfy: $\phi(mg) = (\delta^{1/2}\tau)(m)\phi(g)$ for all $m \in \mathbf{M}(\mathbb{A}_F)$ and $g \in \mathbf{H}(\mathbb{A}_F)$. Here, $\delta(m)$ is defined as $|\det(\text{Ad } m|_{\mathfrak{n}})|$, where $\mathfrak{n}$ denotes the Lie algebra of the unipotent component of $\mathbf{P}$. For any function f in $C(\mathbf{H}(\mathbb{A}_F), \omega)$, let $I_{P,\tau}(\zeta, f)$ denote the convolution operator

$$\int_{\mathbf{Z}_0(\mathbb{A}_F) \backslash \mathbf{H}(\mathbb{A}_F)} (I_{P,\tau}(\zeta))(h) f(h) dh,$$

where dh is a fixed Tamagawa measure on $\mathbf{H}(\mathbb{A}_F)$.

Let $W_H(M)$ denote the group of automorphisms of $\mathfrak{a}_M$ obtained by restricting elements $s \in W(A_0, H)$ which satisfy $s(\mathbf{M}) = \mathbf{M}$. We define an action of $W_H(M)$ on the set of all $\mathbf{M}(\mathbb{A}_F)$ -modules as follows: For any element s in $W_H(M)$ and any $\mathbf{M}(\mathbb{A}_F)$ -module τ , let $s\tau$ denote the $\mathbf{M}(\mathbb{A}_F)$ -module:

$$m \mapsto \tau((s')^{-1}m), \quad \forall m \in \mathbf{M}(\mathbb{A}_F),$$

where s' is a preimage of s in $W(A_0, H)$. Up to equivalence of representations, $s\tau$ is independent of this choice of preimage.

Suppose τ is equivalent to $s\tau$ for some $s \in W_G(M)$. Let $I_{P,\tau}(\varepsilon)$ be the operator on the space of $I_{P,\tau}$ which sends $\phi(g)$ to $\varepsilon(g)\phi(g)$. Then, $I_{P,\tau}(\varepsilon)$ intertwines $I_{P,\tau}(\zeta)$ with $\varepsilon I_{P,\tau}(\zeta)$ for any $\zeta \in \mathfrak{a}_M$. Let $M_P(s, \zeta)$ be an operator which intertwines $I_{P,\tau}(\zeta)$ with $I_{P,s\tau}(s\zeta)$. Then, $I_{P,s\tau}(\varepsilon)M_P(s, \zeta)$ intertwines $I_{P,\tau}(\zeta)$ with $\varepsilon I_{P,s\tau}(s\zeta) = \varepsilon I_{P,\tau}(s\zeta)$. Put

$$I_{P,\tau}(\zeta, f \times \varepsilon) := I_{P,\tau}(\zeta, f)I_{P,\tau}(\varepsilon).$$

The ε -twisted trace formula ([CLL]) is the equality

$$(3.1) \quad \sum_{\{\mathcal{O}\}} J_{\mathcal{O},\varepsilon}^T(f) = \sum_{\{\chi\}} J_{\chi,\varepsilon}^T(f),$$

where the left (resp. right) hand side of the equation is the integral over the diagonal subgroup of the modified geometric (resp. spectral) kernel of the operator $\rho(f)\rho(\varepsilon)$ (see Chapter 1 for the definitions of symbols). The modification depends on an element T , called a **truncation parameter**, in the Lie algebra of a fixed minimal Levi subgroup of $\mathbf{H}$. Each side of (3.1) is defined only when T is sufficiently regular (see [CLL]). The sum $\sum_{\{\chi\}} J_{\chi,\varepsilon}^T(f)$ is called the (twisted-)fine χ -**expansion** of $\mathbf{H}$.

It is important to note that the authors of [CLL] have skipped giving a precise form of the fine χ -expansion for the twisting by $\rho(\varepsilon)$. We take the liberty here to describe without proof the sum $\sum_{\{\chi\}} J_{\chi,\varepsilon}^T(f)$ (For instance, we can bypass the twisting by considering the untwisted trace formula for the group described in Lemma 4.39).

For f in $C(\mathbf{H}(\mathbb{A}_F), \omega)$, the fine χ -expansion is a sum over the set of quadruples $\{\chi\} = (\mathbf{M}, \mathbf{L}, \tau, s)$ consisting of Levi subgroups $\mathbf{M}, \mathbf{L}$ of $\mathbf{H}$, an element $s \in W_G(M)$, and a discrete spectrum automorphic representation τ of $\mathbf{M}(\mathbb{A}_F)$ such that:

- $\mathbf{M} \subset \mathbf{L}$;
- $\mathfrak{a}_L$ is the subspace of $\mathfrak{a}_M$ fixed pointwise by s ;
- τ is trivial on $\exp \mathfrak{a}_M$;
- the restriction of τ to $\mathbf{Z}_0(\mathbb{A}_F)$ is equal to ω ;
- $s\tau$ is equivalent to $\varepsilon\tau$.

We say that a parabolic subgroup is *standard* if it contains the fixed minimal parabolic subgroup $\mathbf{P}_0$. Let $\mathbf{P} \in \mathcal{P}(M)$ be the standard parabolic subgroup with Levi component $\mathbf{M}$. The term associated with $(\mathbf{M}, \mathbf{L}, \tau, s)$ in the fine χ -expansion is the product of

$$(3.2) \quad \frac{|W(A_0, M)|}{|W(A_0, H)|} |\det(1 - s)|_{\mathfrak{a}_M/\mathfrak{a}_L}^{-1}$$

and

$$(3.3) \quad (2\pi)^{-\dim(\mathfrak{a}_L/\mathfrak{a}_H)} \int_{i\mathfrak{a}_L^*/i\mathfrak{a}_H^*} \mathrm{tr} \mathcal{M}_L^T(P, \zeta) I_{P,\tau}(\zeta, f \times \varepsilon) M_P(s, \zeta) d\zeta.$$

Here, $\mathcal{M}_L^T(P, \zeta)$ is the logarithmic derivative of an intertwining operator. We will not reproduce here the definition of $\mathcal{M}_L^T(P, \zeta)$. We only use the fact that $\mathcal{M}_L^T(P, \zeta) = 1$ when

$\mathbf{L}$ is equal $\mathbf{H}$. In particular, the term in the expansion associated with a quadruple of the form $(\mathbf{M}, \mathbf{H}, \tau, s)$ is

$$(3.4) \quad \frac{|W(A_0, M)|}{|W(A_0, H)|} |\det(1-s)|_{\mathfrak{a}_M/\mathfrak{a}_L}^{-1} \operatorname{tr} I_{P,\tau}(\zeta, f \times \varepsilon) M_P(s, 0).$$

We call the sum of all the terms in the form (3.4) the **discrete part** of the fine χ -expansion of the ε -twisted trace formula. We denote it by $I^d(H, f, \varepsilon)$. We call the sum of the rest of the terms the **continuous part** and denote it by $I^c(H, f, \varepsilon)$. If $\varepsilon = 1$, we simply write $I^d(H, f)$ (resp. $I^c(H, f)$) for $I^d(H, f, 1)$ (resp. $I^c(H, f, 1)$). Observe that the ε -twisted trace of a discrete spectrum representation π of $\mathbf{H}(\mathbb{A}_F)$ corresponds to the quadruple $(\mathbf{H}, \mathbf{H}, \pi, 1)$. For any quadruple of the form $(\mathbf{M}, \mathbf{H}, \tau, s)$ and parabolic subgroup $\mathbf{P}$ in $\mathcal{P}(M)$, we say that the (normalizedly) parabolically induced representation $I_{P,\tau}$ **occurs ε -discretely** in the spectrum of $\mathbf{H}(\mathbb{A}_F)$.

In this work, we only study the discrete parts of the spectral expansions of the groups. We now list the terms which occur in $I^d(\operatorname{GSp}(2), f, \varepsilon)$ and $I^d(H_i, f_i)$ ($i = 1, 2$).

Notation:

- For any number field L , let C_L denote the idèle class group $L^\times \backslash \mathbb{A}_L^\times$.
- For any character θ of C_E , put ${}^\sigma\theta(x) := \theta(\sigma x)$ for all $x \in \mathbb{A}_E^\times$. We say that θ is **σ -invariant** if $\theta = {}^\sigma\theta$.
- Let ω_π denote the central character of a representation π .

Terminology:

- Let L be a quadratic extension of F . We say that an automorphic representation π of $\operatorname{GL}(2, \mathbb{A}_F)$ is **L -monomial** if it is the monomial representation $\pi(\theta)$ associated with a character θ of C_L . In particular, if μ is the class field character of L/F , then $\mu\pi(\theta)$ is equivalent to $\pi(\theta)$, and the central character $\omega_{\pi(\theta)}$ is equal to $\theta|_{\mathbb{A}_F^\times} \cdot \mu$ (see [JL, Thm. 4.6], [K]).
- Let **(ε) -DOR** stand for “ (ε) -discretely occurring representation.”

3.1.2. ε -DOR of $\operatorname{GSp}(2, \mathbb{A}_F)$. Let $\mathbf{G} = \operatorname{GSp}(2)$. Let $\mathbf{Z}$ be the center of $\mathbf{G}$. It is isomorphic to $\mathbb{G}_m$. In particular, the maximal F -split component $\mathbf{Z}_0$ of the center is $\mathbf{Z}$ itself. We fix a character ω of $\mathbf{Z}_0(F) \backslash \mathbf{Z}_0(\mathbb{A}_F)$.

Recall that we have fixed a quadratic character ε of C_F . It induces a character of $\operatorname{GSp}(2, \mathbb{A}_F)$ defined by $\varepsilon(g) = \varepsilon(\lambda(g))$. In this section, we list all quadruples $(\mathbf{M}, \mathbf{G}, \tau, s)$ which satisfy the conditions mentioned in the previous section.

If the quadruple has the form $(\mathbf{G}, \mathbf{G}, \tau, s)$, then s is necessarily trivial, and τ can be any discrete spectrum automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ with the property that $\omega_\tau = \omega$. In this case, the intertwining operator $M_G(1, 0)$ is equal to 1. The contribution to the fine χ -expansion corresponding to $(\mathbf{G}, \mathbf{G}, \tau, 1)$ is therefore

$$\operatorname{tr} \tau(f \times \varepsilon),$$

where $\tau(f \times \varepsilon)$ denotes $\tau(f)\rho(\varepsilon)$.

We now list all the valid quadruples for the proper Levi subgroups $\mathbf{M}$ of $\mathbf{G}$. There are 5 proper Levi subgroups: $\mathbf{A}_0, \mathbf{M}_\alpha, \mathbf{M}'_\alpha, \mathbf{M}_\beta, \mathbf{M}'_\beta$, which we shall describe shortly.

We choose $\mathbf{P}_0$ to be the upper triangular subgroup in $\mathbf{G}$. Its Levi subgroup is $\mathbf{A}_0 = \{\operatorname{diag}(a, b, \lambda/b, \lambda/a) : a, b, \lambda \in \mathbb{G}_m\}$. The F -split component of the center of $\mathbf{A}_0$ is $\mathbf{A}_0$ itself. A basis of the root system associated with the pair $(\mathbf{G}, \mathbf{A}_0)$ is

$$\Delta = \Delta(\mathbf{G}, \mathbf{P}_0, \mathbf{A}_0) = \{\alpha, \beta\},$$

where $\alpha(t) := a/b$, $\beta(t) := b^2/\lambda$ for all $t = \text{diag}(a, b, \lambda/a, \lambda/b)$. The root system is $R = R^+ \cup -R^+$, where $R^+ = R^+(\mathbf{G}, \mathbf{P}_0, \mathbf{A}_0) = \{\alpha, \beta, \alpha + \beta, 2\alpha + \beta\}$ is a fixed set of positive roots. Let $X_*(\mathbf{A}_0) = \text{Hom}(\mathbb{G}_m, \mathbf{A}_0)$. Let $\mathfrak{a}_0 = X_*(\mathbf{A}_0) \otimes \mathbb{R}$. We identify $\mathfrak{a}_0$ with the real vector space $\{(x, y, t - x, t - y) : x, y, t \in \mathbb{R}\} \cong \mathbb{R}^3$.

Let $\mathcal{V}$ be the space of weights ([Sp]) associated with the pair $(\mathbf{G}, \mathbf{A}_0)$. The Weyl group $W(A_0, G)$ of $\mathbf{A}_0$ in $\mathbf{G}$ corresponds to the group of automorphisms of $\mathcal{V}$ which are generated by reflections associated with the roots in R^+ . For a root $\gamma \in R^+$, let s_γ denote the Weyl group element which corresponds to the reflection over γ .

The Weyl group $W(A_0, G)$ is isomorphic to D_4 , the dihedral group of order 8. Viewed as a permutation group on four letters (corresponding to the four entries of an element in $\mathbf{A}_0$), it consists of the following elements:

- $s_\alpha = (12)(34)$,
- $s_\beta = (23)$,
- $s_{2\alpha+\beta}s_\alpha = (1243)$,
- $\rho := s_\beta s_\alpha = (1342)$,
- $s_{\alpha+\beta} = (24)(13)$,
- $s_{2\alpha+\beta}s_\beta = (23)(14)$,
- $s_{2\alpha+\beta} = (14)$.

Note that $\rho = s_\beta s_\alpha$ has order 4, and $W(A_0, G) = \langle s_\beta, \rho \rangle$.

Let e denote the element $\text{diag}(1, -1)$ in $\text{GL}(2)$. The proper Levi subgroups of $\mathbf{G}$ which strictly contain $\mathbf{A}_0$ are:

$$\begin{aligned} \mathbf{M}_\alpha &= \left\{ \text{blockdiag} \left(g, \frac{\lambda}{\det g} ege \right) : g \in \text{GL}(2), \lambda \in \mathbb{G}_m \right\}, \quad \mathbf{M}'_\alpha = s_\beta \mathbf{M}_\alpha s_\beta, \\ \mathbf{M}_\beta &= \{ \text{blockdiag}(a, g, (\det g)/a) : a \in \mathbb{G}_m, g \in \text{GL}(2) \}, \quad \mathbf{M}'_\beta = s_\alpha \mathbf{M}_\beta s_\alpha. \end{aligned}$$

3.1.2.1. ε -DOR associated with $\mathbf{A}_0$. Suppose $(\mathbf{A}_0, \mathbf{G}, \tau, s)$ is a quadruple satisfying the conditions outlined in Section 3.1.1. The discrete representation τ of $\mathbf{A}_0(\mathbb{A}_F)$ has the form:

$$\tau = \mu_1 \otimes \mu_2 \otimes \mu : \text{diag}(a, b, \lambda/b, \lambda, a) \mapsto \mu_1(a) \mu_2(b) \mu(\lambda)$$

for some characters μ_1, μ_2, μ of C_F . Let $\mu_1 \times \mu_2 \rtimes \mu$ denote the (normalizedly) parabolically induced representation $I_{P_0}(\mu_1 \otimes \mu_2 \otimes \mu)$ of $\mathbf{G}(\mathbb{A}_F)$.

For $\mathfrak{a}_0^s$ to equal $\mathfrak{a}_G$, the element s in $W_G(A_0) = W(A_0, G)$ must be one of ρ, ρ^2, ρ^3 . By assumption, $s\tau \cong \varepsilon\tau$ and $\omega_\tau = \omega$, which implies that the following holds:

- If $s = \rho$ or ρ^3 , then τ must have the form $\varepsilon \otimes \varepsilon \otimes \mu$ for some character μ of C_F such that $\mu^2 = \omega$.
- If $s = \rho^2$, then τ must have the form $\varepsilon' \otimes \varepsilon' \varepsilon \otimes \mu$ for some characters ε', μ of C_F such that $(\varepsilon')^2 = 1$ and $\varepsilon\mu^2 = \omega$.

We have: $|W(A_0, A_0)| = 1$, $|W(A_0, G)| = 8$. For $s = \rho, \rho^3$, $\det((1-s)|_{\mathfrak{a}_0/\mathfrak{a}_G})$ is equal to 2, while $|\det(1-\rho^2)|_{\mathfrak{a}_0/\mathfrak{a}_G}| = 4$.

Notation: For any representation τ of $\mathbf{M}(\mathbb{A}_F)$ and element w in the Weyl group $W(A_0, G)$, let ${}^w\tau$ denote $\tau \circ w$.

The orbit of $\varepsilon \otimes \varepsilon \otimes \mu$ under $W(A_0, G) = D_4$ is of order 2. Consequently, the set of quadruples $\{(\mathbf{A}_0, \mathbf{G}, {}^w\tau, s) : s = \rho, \rho^2, w \in W(A_0, G)\}$ has 4 distinct elements, and they all give rise to the same representation $I_{P_0}(\varepsilon \otimes \varepsilon \otimes \mu) = \varepsilon \times \varepsilon \rtimes \mu$. Since $\varepsilon \times \varepsilon \rtimes \mu$ is irreducible, by Schur's lemma we may normalize the intertwining operators so that $M_{P_0}(\rho, 0) = M_{P_0}(\rho^2, 0) = 1$.

We conclude that the contribution from $\varepsilon \times \varepsilon \rtimes \mu$ to the fine χ -expansion is

$$4 \cdot \frac{1}{8} \frac{1}{2} \cdot \text{tr} (\varepsilon \times \varepsilon \rtimes \mu) (f \times \varepsilon).$$

Similarly, the orbit of $\varepsilon' \otimes \varepsilon' \varepsilon \rtimes \mu$ under $W(A_0, G)$ is of order 8, and the irreducibility of $\varepsilon' \times \varepsilon' \varepsilon \rtimes \mu$ implies that $M_{P_0}(\rho^2, 0) = 1$. Consequently, the contribution from $\varepsilon' \times \varepsilon' \varepsilon \rtimes \mu$ to the fine χ -expansion is

$$8 \cdot \frac{1}{8} \frac{1}{4} \cdot \text{tr} (\varepsilon' \times \varepsilon' \varepsilon \rtimes \mu) (f \times \varepsilon).$$

To summarize, the contribution to the fine χ -expansion from quadruples of the form $(\mathbf{A}_0, \mathbf{G}, \tau, s)$ is:

$$(3.5) \quad \frac{1}{4} \sum_{\mu^2=\omega} \text{tr} (\varepsilon \times \varepsilon \rtimes \mu) (f \times \varepsilon) + \frac{1}{4} \sum_{\substack{\varepsilon'^2=1, \\ \varepsilon' \mu^2=\omega}} \text{tr} (\varepsilon' \times \varepsilon' \varepsilon \rtimes \mu) (f \times \varepsilon).$$

3.1.2.2. ε -DOR associated with $\mathbf{M}_\alpha$. Suppose $(\mathbf{M}_\alpha, \mathbf{G}, \tau, s)$ is a quadruple satisfying the conditions outlined in Section 3.1.1. The discrete spectrum representation τ of $\mathbf{M}_\alpha(\mathbb{A}_F)$ has the form:

$$\tau = \pi_2 \otimes \mu : \text{blockdiag} \left(g, \frac{\lambda}{\det g} ege \right) \mapsto \mu(\lambda) \pi_2(g),$$

where π_2 is a discrete spectrum representation of $\text{GL}(2, \mathbb{A}_F)$ and μ is a character of C_F . Let $\mathbf{P}_\alpha$ be the standard parabolic subgroup of $\mathbf{G}$ whose Levi component is $\mathbf{M}_\alpha$. Put

$$\pi_2 \rtimes \mu := I_{P_\alpha}(\pi_2 \otimes \mu).$$

The group $W_G(M_\alpha)$ is represented by the elements $\{1, (24)(13)\}$ in $W(A_0, G)$, and $\mathfrak{a}_{M_\alpha}$ is equal to $\{(x, x, t - x, t - x) : x, t \in \mathbb{R}\} \cong \mathbb{R}^2$. For $\mathfrak{a}_M^s$ to equal $\mathfrak{a}_G$, the element $s \in W_G(M_\alpha)$ must be the restriction of $s_{\alpha+\beta} = (24)(13)$. The following statements hold:

- π_2 is ε -invariant; i.e., π_2 is equivalent to the representation

$$\varepsilon \pi_2 : g \mapsto \varepsilon(\det g) \pi_2(g), \quad \forall g \in \text{GL}(2, \mathbb{A}_F).$$

- $\omega_{\pi_2} = \varepsilon$ and $\varepsilon \mu^2 = \omega$.

By [JL, Thm. 4.6], there exists a character η of C_E such that $\eta \neq {}^\sigma \eta$ and π_2 is the cuspidal monomial representation $\pi(\eta)$ associated with η . Moreover, since $\omega_{\pi(\eta)}$ is equal to $\eta|_{\mathbb{A}_F^\times} \cdot \varepsilon$, the character $\eta|_{\mathbb{A}_F^\times}$ must be trivial.

We have: $|W(A_0, M_\alpha)| = 2$, $|\det((1 - s_{\alpha+\beta})|_{\mathfrak{a}_{M_\alpha}/\mathfrak{a}_G})| = 2$. The orbit of the representation $\tau = \pi(\eta) \otimes \mu$ under $W(A_{M_\alpha}, G)$ is of order 2, and both elements in the orbit give rise to the same induced representation $\pi(\eta) \rtimes \mu$ of $\mathbf{G}(\mathbb{A}_F)$. Since $\pi(\eta) \rtimes \mu$ is irreducible ([ST]), we may normalize the intertwining operator $M_{P_\alpha}(s_{\alpha+\beta}, 0)$ to be 1.

For each $\{\chi\} = (\mathbf{M}_\alpha, \mathbf{G}, \tau, s)$, there is a representation τ' of $\mathbf{M}'_\alpha(\mathbb{A}_F)$ and an element s' in $W_G(M'_\alpha)$ such that the quadruple $(\mathbf{M}'_\alpha, \mathbf{G}, \tau', s')$ contributes a term which is equal to the contribution from $\{\chi\}$. The contribution of $\text{tr} (\pi(\eta) \rtimes \mu) (f \times \varepsilon)$ to the fine χ -expansion is therefore:

$$(3.6) \quad 2 \cdot 2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum_{\eta|_{\mathbb{A}_F^\times}=1, \varepsilon \mu^2=\omega} \text{tr} (\pi(\eta) \rtimes \mu) (f \times \varepsilon).$$

3.1.2.3. ε -DOR associated with $\mathbf{M}_\beta$. Suppose $(\mathbf{M}_\beta, \mathbf{G}, \tau, s)$ is a quadruple satisfying the conditions listed in Section 3.1.1. The discrete representation τ of $\mathbf{M}_\beta(\mathbb{A}_F)$ has the form:

$$\tau = \mu \otimes \pi_2 : \text{blockdiag} \left(a, g, \frac{\det g}{a} \right) \mapsto \mu(a) \pi_2(g),$$

where π_2 is a discrete spectrum representation of $\text{GL}(2, \mathbb{A}_F)$ and μ is a character of C_F . Let $\mathbf{P}_\beta$ be the standard parabolic subgroup of $\mathbf{G}$ whose Levi component is $\mathbf{M}_\beta$. Put

$$\mu \rtimes \pi_2 := I_{\mathbf{P}_\beta}(\mu \otimes \pi_2).$$

The group $W_G(M_\beta)$ is represented by $\{1, (14)\}$ in $W(A_0, G)$, and $\mathfrak{a}_{M_\beta}$ is equal to $\{(x, y, y, 2y - x) : x, y \in \mathbb{R}\} \cong \mathbb{R}^2$. For $\mathfrak{a}_M^s$ to equal $\mathfrak{a}_G$, the element $s \in W_G(M_\alpha)$ must be the restriction of $s_{2\alpha+\beta} = (14)$. The following statements hold:

- $\mu^2 = 1$.
- π_2 is $\mu\varepsilon$ -invariant.
- $\mu\omega_{\pi_2} = \omega$.

The classification can be refined further:

- If $\mu = 1$, then $\pi_2 = \pi(\chi)$ for some character χ of C_E such that $\chi \neq {}^\sigma\chi$ and $\chi|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$ (see [JL], [K]). Observe that the orbit of $1 \otimes \pi(\chi)$ under $W_G(M_\beta)$ is of order 1.
- If $\mu \neq 1, \varepsilon$, let $E_{\mu\varepsilon}$ be the quadratic extension of F corresponding to $\mu\varepsilon$ via global class field theory. Suppose $\text{Gal}(E_{\mu\varepsilon}/F) = \langle \sigma' \rangle$. Then, $\pi_2 = \pi(\theta)$ for some character θ of $C_{E_{\mu\varepsilon}}$ such that $\theta \neq {}^{\sigma'}\theta$ and $\theta|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$.

Let E_μ be the quadratic extension of F corresponding to μ . The orbit of $\mu \otimes \pi(\theta)$ under $W_G(M_\beta)$ is of order 2 if $\pi(\theta)$ is not E_μ -monomial (i.e., not a monomial representation associated with a character of C_{E_μ}). It is of order 1 if $\pi(\theta)$ is E_μ -monomial.

- If $\mu = \varepsilon$, then π_2 can be any cuspidal or one dimensional automorphic representation of $\text{GL}(2, \mathbb{A}_F)$ with central character $\omega_{\pi_2} = \varepsilon\omega$. The orbit of $\varepsilon \otimes \pi_2$ under $W_G(M_\beta)$ is of order 1 if π_2 is E -monomial and of order 2 otherwise.

We have: $|W(A_0, M_\beta)| = 2$, $|\det((1 - s_{2\alpha+\beta})|_{\mathfrak{a}_{M_\beta}/\mathfrak{a}_G})| = 2$. As in the case of $\mathbf{M}_\alpha$, for every contribution to the spectral expansion from a quadruple of the form $(\mathbf{M}_\beta, \mathbf{G}, \tau, s)$, there is an equal contribution from a quadruple of the form $(\mathbf{M}'_\beta, \mathbf{G}, \tau', s')$. Moreover, the elements in the orbit of $\tau = \mu \otimes \pi_2$ under $W_G(M_\beta)$ all give rise to the same representation $\mu \rtimes \pi_2$. Consequently, the contribution from representations of the form $\mu \rtimes \pi_2$ to the fine χ -expansion is the sum of the following terms:

- (1) $2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum \text{tr} (1 \rtimes \pi(\chi)) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0)$, where the sum is over the characters χ of C_E such that $\chi|_{\mathbb{A}_F^\times} = \varepsilon\omega$.
- (2) $2 \cdot 2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum \text{tr} (\mu \rtimes \pi(\theta)) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0)$. The sum is over the nontrivial quadratic characters $\mu \neq \varepsilon$ of C_F and cuspidal $E_{\mu\varepsilon}$ -monomial representations $\pi(\theta)$ of $\text{GL}(2, \mathbb{A}_F)$, where θ is a character of $C_{E_{\mu\varepsilon}}$, such that: $\pi(\theta)$ is not E_μ -monomial and $\theta|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$. Let σ' be the generator of the $\text{Gal}(E_{\mu\varepsilon}/F)$. It is known that $\pi(\theta)$ is E_μ -monomial if and only if $\sigma'\theta/\theta = \mu \circ N_{E/F}$ (see [LL]).
- (3) $2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum \text{tr} (\mu \rtimes \pi(\theta)) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0)$. The sum is over the nontrivial quadratic characters $\mu \neq \varepsilon$ of C_F , cuspidal $E_{\mu\varepsilon}$ -monomial representations $\pi(\theta)$

of $\mathrm{GL}(2, \mathbb{A}_F)$, where θ is a character of $C_{E_{\mu\varepsilon}}$, such that: $\sigma'\theta/\theta$ is equal to μ and $\theta|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$.

- (4) $2 \cdot 2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum \mathrm{tr} (\varepsilon \rtimes \pi) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0)$. The sum is over all cuspidal non- E -monomial, or one dimensional, automorphic representations π of $\mathrm{GL}(2, \mathbb{A}_F)$ with central character $\omega_\pi = \varepsilon\omega$.
- (5) $2 \cdot \frac{1}{4} \cdot \frac{1}{2} \sum \mathrm{tr} (\varepsilon \rtimes \pi) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0)$. The sum is over all cuspidal E -monomial automorphic representations π of $\mathrm{GL}(2, \mathbb{A}_F)$ with central character $\omega_\pi = \varepsilon\omega$.

Suppose an automorphic representation $\pi = \otimes_v \pi_v$ of $\mathbf{G}(\mathbb{A}_F)$ is ε -invariant. Assuming that the multiplicity one theorem holds for $\mathrm{GSp}(2)$, the intertwining operator $\rho(\varepsilon)$ maps the space of π back to itself. Consequently, $\rho(\varepsilon)$ defines at each place v a local operator $\rho(\varepsilon)_v$ which intertwines π_v with $\varepsilon_v \pi_v$. Hence, for $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$, the twisted character $\mathrm{tr} \pi(f \times \varepsilon)$ may be expressed as the product $\prod_v \mathrm{tr} \pi_v(f_v \times \varepsilon_v)$ of twisted local characters. Here, $\mathrm{tr} \pi_v(f_v \times \varepsilon_v) := \mathrm{tr} \pi_v(f_v) \rho(\varepsilon)_v$.

We now examine the behavior of the intertwining operator $M_{P_\beta}(s_{2\alpha+\beta}, 0)$. Let π be the induced representation $\mu \rtimes \pi_2$, where μ is a character of C_F , and π_2 is a cuspidal or one dimensional automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. At any place v of F for which $\mu_v = 1$ and $\pi_{2,v}$ is square integrable or one dimensional, the local representation π_v is of length two, with constituents π_v^+, π_v^- (tempered if $\pi_{2,v}$ is square integrable, nontempered otherwise) (see [ST]). We normalize intertwining operators so that the local component $M_{P_\beta}(s_{2\alpha+\beta}, 0)_v$ of $M_{P_\beta}(s_{2\alpha+\beta}, 0)$ acts trivially on π_v^+ , and via scalar multiplication by -1 on π_v^- . At a place v where $\pi_{2,v}$ is parabolically induced and/or $\mu_v \neq 1$, the representation π_v is irreducible. For convenience, for such v we put $\pi_v^+ := \pi_v$ and $\pi_v^- := 0$. Hence, $M_{P_\beta}(s_{2\alpha+\beta}, 0)_v$ acts trivially on π_v^+ and via scalar multiplication by -1 on π_v^- , for all v .

Using the above notation, we write (recall that we assume $f = \otimes_v f_v$):

$$(3.7) \quad \mathrm{tr} \pi(f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0) = \prod_{v \in V} [\mathrm{tr} \pi_v^+(f_v \times \varepsilon_v) - \mathrm{tr} \pi_v^-(f_v \times \varepsilon_v)].$$

3.1.3. DOR for $\mathbf{H}_1(\mathbb{A}_F)$. The norm mapping $N_{E/F} : E \rightarrow F$ induces the norm mapping $N_{E/F} : \mathbb{A}_E \rightarrow \mathbb{A}_F$ on the adèles. Let $\mathrm{GL}(2, \mathbb{A}_F)^E$ denote the subgroup of $\mathrm{GL}(2, \mathbb{A}_F)$ consisting of elements whose determinants lie in $N_{E/F} \mathbb{A}_E^\times$. For any representation τ of $\mathrm{GL}(2, \mathbb{A}_F)$ and character χ of C_E , let $\tau \otimes_1 \chi$ denote the representation of $\mathbf{H}_1(\mathbb{A}_F)$ in the space of τ defined as follows:

$$\tau \otimes_1 \chi : (g, x) \mapsto \chi(x) (\tau|_{\mathrm{GL}(2, \mathbb{A}_F)^E})(g), \quad \forall (g, x) \in \mathbf{H}_1(\mathbb{A}_F).$$

Since any $(g, x) \in \mathbf{H}_1(\mathbb{A}_F)$ satisfies $\det g = N_{E/F} x$, the representations $\tau \otimes_1 \chi$ and $\mu^{-1} \tau \otimes_1 (\mu \circ N_{E/F}) \chi$ are equal to each other for any quasicharacter μ of C_F . Moreover, since the representation $\tau \otimes_1 \chi$ is defined by χ and the restriction of τ to the subgroup $\mathrm{GL}(2, \mathbb{A}_F)^E$ of $\mathrm{GL}(2, \mathbb{A}_F)$, it may be reducible even if τ is an irreducible representation of $\mathrm{GL}(2, \mathbb{A}_F)$ (see [LL]).

3.1.3.1. Discrete Spectrum Representations. Let $(\mathbf{H}_1, \mathbf{H}_1, \pi, 1)$ be a quadruple for the group $\mathbf{H}_1$. Then, π is an irreducible, discrete spectrum, automorphic representation of $\mathbf{H}_1(\mathbb{A}_F)$, and it is an irreducible constituent of $\tau \otimes_1 \eta$ for some irreducible, discrete spectrum, automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_F)$ and character μ of C_E .

For any place v of F which remains prime in E , let

$$\mathrm{GL}(2, F_v)^E = \{g \in \mathrm{GL}(2, F_v) : \det g \in N_{E/F} E_v^\times\}.$$

If v splits in E , let $\mathrm{GL}(2, F_v)^E = \mathrm{GL}(2, F_v)$. The adèlic group $\mathrm{GL}(2, \mathbb{A}_F)^E$ is equal to the restricted tensor product $\otimes_{v \in V} \mathrm{GL}(2, F_v)^E$ (namely, almost all local components lie in $\mathrm{GL}(2, \mathcal{O}_v) \cap \mathrm{GL}(2, F_v)^E$), and $\tau|_{\mathrm{GL}(2, \mathbb{A}_F)^E} = \otimes_v \tau_v|_{\mathrm{GL}(2, F_v)^E}$.

Let v be any place of F . From [LL], the restriction of τ_v to $\mathrm{GL}(2, F_v)^E$ is reducible if and only if v is prime in E , and τ_v is the E_v -monomial representation associated with some character θ_v of $E_v^\times$. In this case, the representation $\tau_v|_{\mathrm{GL}(2, F_v)^E}$ has length two, with constituents τ_v^+ and τ_v^- . For convenience, if τ_v is irreducible, we let $\tau_v^+ = \tau_v$ and $\tau_v^- = 0$. If $\tau_v|_{\mathrm{GL}(2, F_v)^E}$ is unramified, we let τ_v^+ be the constituent which contains the (unique up to scalar multiple) $\mathrm{GL}(2, \mathcal{O}_v)$ -fixed vector. If $\tau_v|_{\mathrm{GL}(2, F_v)^E}$ has length two and is ramified, then which constituent to label τ_v^+ and which to label τ_v^- is irrelevant for our purpose.

We let $\pi_v^+ = \tau_v^+ \otimes_1 \eta_v$ and $\pi_v^- = \tau_v^- \otimes_1 \eta_v$ for any place v of F . Let $\tau_v \otimes_1 \eta_v$ denote the **local packet** $\{\pi_v^+, \pi_v^-\}$. We define the **global packet** $\tau \otimes_1 \eta$ to be the *restricted tensor product*

$$\otimes_{v \in V} \{\tau_v \otimes_1 \eta_v\} := \{\otimes_v \pi'_v : \pi'_v \in \{\pi_v^+, \pi_v^-\}, \pi'_v = \pi_v^+ \text{ almost all } v\}.$$

Recall our assumption that the global test functions are tensor products of local components. For $f_1 = \otimes_{v \in V} f_{1,v}$ in $C(\mathbf{H}_1(\mathbb{A}_F), \omega)$, put

$$\mathrm{tr} (\tau \otimes_1 \eta) (f_1) := \prod_{v \in V} \left[\mathrm{tr} \pi_v^+(f_v) + \mathrm{tr} \pi_v^-(f_v) \right].$$

The distribution $f_1 \mapsto \mathrm{tr} (\tau \otimes_1 \eta) (f_1)$ defines a distribution on $C(\mathbf{H}_1(\mathbb{A}_F), \omega)$ which is invariant under stable conjugacy.

If τ is not an E -monomial representation, then by [LL] each member of $\tau \otimes_1 \eta$ occurs with multiplicity one in the discrete spectrum of $\mathbf{H}_1(\mathbb{A}_F)$. Thus, the contribution of the packet $\tau \otimes_1 \eta$ to the spectral side of the trace formula of $\mathbf{H}_1(\mathbb{A}_F)$ is

$$\mathrm{tr} (\tau \otimes_1 \eta) (f_1).$$

If τ is cuspidal E -monomial, the multiplicity $m(\pi')$ with which each member π' of $\{\pi\}$ occurs in the discrete spectrum of $\mathbf{H}_1(\mathbb{A}_F)$ is given by the following formula:

$$m(\pi') = \frac{1}{2} \left(1 + (-1)^{n(\pi')} \right),$$

where $n(\pi')$ is the number of places v for which $\pi'_v = \pi_v^-$ ([LL]). Taking into account the nontrivial multiplicity formula, the contribution of $\tau \otimes_1 \eta$ to the spectral side of the trace formula of $\mathbf{H}_1(\mathbb{A}_F)$ is equal to $D(\tau \otimes_1 \eta, f_1) :=$

$$\frac{1}{2} \prod_{v \in V} \left[\mathrm{tr} \pi_v^+(f_{1,v}) + \mathrm{tr} \pi_v^-(f_{1,v}) \right] + \frac{1}{2} \prod_{v \in V} \left[\mathrm{tr} \pi_v^+(f_{1,v}) - \mathrm{tr} \pi_v^-(f_{1,v}) \right].$$

Note that the first term in the above sum is equal to $\frac{1}{2} \cdot \mathrm{tr} (\tau \otimes_1 \eta) (f_1)$. If v is prime in E , there exists an $h \in \mathbf{H}_1(\overline{F}_v)$ such that conjugation by h swaps π_v^+ with π_v^- . Hence, $\frac{1}{2} \cdot \mathrm{tr} (\tau \otimes_1 \eta) (f_1)$ defines a stable distribution on $C(\mathbf{H}_1(\mathbb{A}_F), \omega)$, whereas the second term in the sum defines an unstable distribution. We call $\frac{1}{2} \cdot \mathrm{tr} (\tau \otimes_1 \eta) (f_1)$ the **stable part** of $D(\tau \otimes_1 \eta, f_1)$.

3.1.3.2. Induced Representations. There is one proper Levi subgroup in $\mathbf{H}_1$; namely, the diagonal torus

$$\mathbf{M}_1 = \{(\mathrm{diag}(a, b), c) : a, b \in \mathbb{G}_m, c \in \mathrm{R}_{E/F} \mathbb{G}_m, ab = N_{E/F} c\}.$$

Suppose $(\mathbf{M}_1, \mathbf{H}_1, \tau, s)$ is a quadruple in the fine χ -expansion of $\mathbf{H}_1(\mathbb{A}_F)$. Any discrete spectrum representation of $\mathbf{M}_1(\mathbb{A}_F)$ has the form:

$$\tau = \mu_1 \otimes \mu_2 \otimes_1 \eta : (\text{diag}(a, b), c) \mapsto \mu_1(a)\mu_2(b)\eta(c),$$

$$\forall (\text{diag}(a, b), c) \in \mathbf{M}_1(\mathbb{A}_F),$$

where μ_1, μ_2 are characters of C_F and η is a character of C_E . From the way $\mathbf{M}_1$ is defined, for any quasi-character μ of C_F , we have:

$$(3.8) \quad \mu_1 \otimes \mu_2 \otimes_1 \eta = \mu\mu_1 \otimes \mu\mu_2 \otimes_1 (\mu^{-1} \circ N_{E/F}) \eta.$$

The F -split component $\mathbf{A}_{M_1}$ of $\mathbf{M}_1$ is $\{(\text{diag}(a, b), c) : a, b, c \in \mathbb{G}_m, ab = c^2\}$, and $\mathfrak{a}_{M_1} = \{(x, y, t) : x, y, t \in \mathbb{R}, x + y = 2t\} \cong \mathbb{R}^2$. The group $W_{H_1}(M_1)$ is equal to $\mathbb{Z}/2\mathbb{Z}$, generated by $(12) : (\text{diag}(a, b), c) \mapsto (\text{diag}(b, a), c)$. The F -split component of the center of $\mathbf{M}_1$ is $\mathbf{A}_{H_1} = \{(\text{diag}(c, c), \pm c) : c \in \mathbb{G}_m\}$, and $\mathfrak{a}_{H_1} = \{(z, z, z) : z \in \mathbb{R}\}$.

If $\mathfrak{a}_{M_1}^s = \mathfrak{a}_{H_1}$ for some $s \in W_{H_1}(M_1)$, then s must be equal to (12) . Consequently, $\tau = \mu_1 \otimes \mu_2 \otimes_1 \eta$ is equivalent to $\mu_2 \otimes \mu_1 \otimes_1 \eta$. In other words,

$$\mu_1(a)\mu_2(b)\eta(c) = \mu_2(a)\mu_1(b)\eta(c), \quad \forall (\text{diag}(a, b), c) \in \mathbf{M}_1(\mathbb{A}_F).$$

Since $ab = N_{E/F}c$, the above equality may be rewritten as

$$\frac{\mu_1}{\mu_2}(a) ((\mu_2 \circ N_{E/F}) \cdot \eta)(c) = \frac{\mu_2}{\mu_1}(a) ((\mu_1 \circ N_{E/F}) \cdot \eta)(c), \quad \forall a \in \mathbb{A}_F^\times, c \in \mathbb{A}_E^\times.$$

Thus, μ_1/μ_2 is equal to either 1 or ε . By (3.8), we conclude that τ must have one of the following forms:

- $1 \otimes 1 \otimes_1 \eta$,
- $1 \otimes \varepsilon \otimes_1 \eta$.

We have: $|W(M_1, M_1)| = 1$, $|W(M_1, H_1)| = 2$, and $|\det(1 - (12))|_{\mathfrak{a}_{M_1}/\mathfrak{a}_{H_1}}$ is equal to 2.

Let $\mathbf{P}_1$ be the upper triangular parabolic subgroup of $\mathbf{H}_1$ whose Levi component is $\mathbf{M}_1$. By (3.4), with $\varepsilon = 1$, the contribution to the fine χ -expansion of $\mathbf{H}_1(\mathbb{A}_F)$ from $\mathbf{M}_1$ is

$$\frac{1}{2} \cdot \frac{1}{2} \sum_{\eta|_{\mathbb{A}_F^\times} = \omega} \left[\text{tr} (I(1, 1) \otimes_1 \eta)(f_1) M_{P_1}((12), 0) \right. \\ \left. + \text{tr} (I(1, \varepsilon) \otimes_1 \eta)(f_1) M_{P_1}((12), 0) \right].$$

The representation $I(1, 1) \otimes_1 \eta$ is irreducible, and we normalize the intertwining operator $M_{P_1}((12), 0)$ to be 1. For any place v , the distribution on the space $C(\mathbf{H}_1(F_v), \omega_v)$ defined by $f_{1,v} \mapsto \text{tr} (I(1, 1) \otimes_1 \eta)_v(f_{1,v})$ is stable.

In the case of $I(1, \varepsilon) \otimes_1 \eta$, the restriction of $I(1, \varepsilon)_v$ to $\text{GL}(2, F_v)^E$ is irreducible if the place v splits in E , and it is reducible of length two, with constituents I_v^+, I_v^- , if v is prime in E . We let I_v^+ denote the unramified constituent of $I(1, \varepsilon)_v$ if $I(1, \varepsilon)_v$ is unramified, and we let $I_v^+ = I(1, \varepsilon)_v$, $I_v^- = 0$ if v splits in E . Write the intertwining operator $M_{P_1}((12), 0)$ for $I(1, \varepsilon) \otimes_1 \eta$ as the tensor product $\otimes_{v \in V} M_{P_1}((12), 0)_v$ of local intertwining operators. If v splits in E , we normalize $M_{P_1}((12), 0)_v$ to be 1. If v is prime in E , we may normalize $M_{P_1}((12), 0)_v$ so that it acts trivially on I_v^+ and it acts via multiplication with -1 on I_v^- .

(see [LL, Sect. 5]). Hence, for $f_1 \in C(\mathbf{H}_1(\mathbb{A}_F), \omega)$, we have

$$\mathrm{tr} (I(1, \varepsilon) \otimes_1 \eta) (f_1) M_{P_1}((12), 0) = \prod_{v \in V} (\mathrm{tr} I_v^+(f_{1,v}) - \mathrm{tr} I_v^-(f_{1,v})).$$

If v is prime in E , there exists an element of $h \in \mathbf{H}_1(\overline{F_v})$ such that the action of conjugation by h swaps I_v^+ and I_v^- ([LL]), multiplying the trace by a factor of -1 . The distribution $f_{1,v} \mapsto \mathrm{tr} (I(1, \varepsilon) \otimes_1 \eta)_v (f_{1,v}) M_{P_1}((12), 0)_v$ is therefore *unstable*.

3.1.3.3. Stable Spectrum. Recall that $I^d(H_1, f_1)$ is the discrete part of the fine χ -expansion for $\mathbf{H}_1(\mathbb{A}_F)$. We let $SI^d(H_1, f_1)$ denote the stable part of $I^d(H_1, f_1)$; namely, $SI^d(H_1, f_1)$ is the sum of the following:

- $\sum \mathrm{tr} \pi(f_1)$.
The sum is over representations π of the form $\tau \otimes_1 \eta$, where τ is a cuspidal non- E -monomial or one dimensional representation of $\mathrm{GL}(2, \mathbb{A}_F)$, and η is a character of C_E , such that $\eta|_{\mathbb{A}_F^\times} \cdot \omega_\tau = \omega$.
- $\frac{1}{2} \sum \mathrm{tr} \pi(f_1)$.
The sum is over representations π of the form $\pi = \pi(\theta) \otimes_1 \eta$, where θ, η are characters of C_E , $\theta \neq {}^\sigma \theta$, $(\theta\eta)|_{\mathbb{A}_F^\times} = \omega$, and $\pi(\theta)$ is the cuspidal, E -monomial automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$ associated with θ .
- $\frac{1}{4} \sum \mathrm{tr} (I(1, 1) \otimes_1 \eta) (f_1)$.
The sum is over characters η of C_E such that $\eta|_{\mathbb{A}_F^\times} = \omega$.

3.1.4. DOR for $\mathbf{H}_2(\mathbb{A}_F)$. For any representation π of $\mathrm{GL}(2, \mathbb{A}_E)$ and character μ of C_F such that $\omega_\pi = \mu \circ N_{E/F}$, let $\pi \otimes_2 \mu$ denote the representation of $\mathbf{H}_2(\mathbb{A}_F)$ defined as follows:

$$\pi \otimes_2 \mu : (g, c) \mapsto \mu(c)\pi(g), \quad \forall (g, c) \in \mathbf{H}_2(\mathbb{A}_F).$$

There is one proper Levi subgroup in $\mathbf{H}_2$; namely, the diagonal torus whose group of F -points is:

$$\mathbf{M}_2(F) = \{(\mathrm{diag}(a, b), c)_* : a, b \in E^\times, c \in F^\times\},$$

where lower star denotes the equivalence class of the element in $\mathrm{GL}(2, E) \times F^\times$ modulo the subgroup $\{(\mathrm{diag}(z, z), N_{E/F} z^{-1}) : z \in E^\times\}$.

Suppose $(\mathbf{M}_2, \mathbf{H}_2, \tau, s)$ is a quadruple in the fine χ -expansion of $\mathbf{H}_2$. Any discrete spectrum representation of $\mathbf{M}_2(\mathbb{A}_F)$ has the form:

$$\tau = \eta_1 \otimes \eta_2 \otimes_2 \mu : (\mathrm{diag}(a, b), c)_* \mapsto \eta_1(a)\eta_2(b)\mu(c), \quad \forall (\mathrm{diag}(a, b), c)_*,$$

where η_1, η_2 are characters of C_E and μ is a character of C_F . Because of the equivalence relation on $\mathbf{H}_2(\mathbb{A}_F)$, the characters must satisfy $\eta_1\eta_2 = \mu \circ N_{E/F}$.

The F -points of the split component $\mathbf{A}_{M_2}$ of $\mathbf{M}_2$ is $\{(\mathrm{diag}(a, b), c)_* : a, b, c \in F^\times\}$, and

$$\mathbf{a}_{M_2} = \{(x, y, t) : x, y, t \in \mathbb{R}\} / \{(z, z, -2z) : z \in \mathbb{R}\} \cong \mathbb{R}^2.$$

The group $W_{H_2}(M_2) = W(H_2)$ is equal to $\mathbb{Z}/2\mathbb{Z}$, generated by the permutation (12) : $(\mathrm{diag}(a, b), c)_* \mapsto (\mathrm{diag}(b, a), c)_*$. The F -points of the split component of the center of $\mathbf{H}_2$ is $\mathbf{A}_{H_2}(F) = \{(\mathrm{diag}(1, 1), c)_* : c \in F^\times\}$, and $\mathbf{a}_{H_2} = \{(0, 0; z) : z \in \mathbb{R}\}$.

If $\mathbf{a}_{M_2}^s = \mathbf{a}_{H_2}$ for some $s \in W_{H_2}(M_2)$, then s must be (12). Consequently,

$$\tau = \chi \otimes \chi \otimes_1 \omega$$

for some character χ of C_E such that $\chi^2 = \omega \circ N_{E/F}$.

We have: $|W(M_2, M_2)| = 1$, $|W(M_2, H_2)| = 2$, and $|\det(1 - (12))|_{\mathfrak{a}_{M_2}/\mathfrak{a}_{H_2}}|$ is equal to 2.

Let $\mathbf{P}_2$ be the upper triangular parabolic subgroup of $\mathbf{H}_2$ whose Levi component is $\mathbf{M}_2$. Then, the contribution to the fine χ -expansion of $\mathbf{H}_2(\mathbb{A}_F)$ from $\mathbf{M}_2$ is:

$$\frac{1}{2} \cdot \frac{1}{2} \sum_{\chi^2 = \omega \circ N_{E/F}} \text{tr} (I(\chi, \chi) \otimes_2 \omega) (f_2) M_{P_2}((12), 0),$$

where f_2 is a test function in $C(\mathbf{H}_2(\mathbb{A}_F), \omega)$. Since $I(\chi, \chi) \otimes_2 \omega$ is irreducible, we normalize $M_{P_2}((12), 0)$ to be 1.

3.1.5. Summary. In summary, $I^d(G, f, \varepsilon)$ is equal to the sum of following:

- $\sum_{\pi \in L_d(\mathbf{G}(\mathbb{A}_F), \omega)} \text{tr} \pi(f \times \varepsilon).$
- $\frac{1}{4} \sum_{\mu^2 = \omega} \text{tr} (\varepsilon \times \varepsilon \rtimes \mu) (f \times \varepsilon).$
- $\frac{1}{4} \sum_{\varepsilon'^2 = 1, \varepsilon' \mu^2 = \omega} \text{tr} (\varepsilon' \times \varepsilon' \rtimes \mu) (f \times \varepsilon)$
- $\frac{1}{4} \sum_{\chi|_{\mathbb{A}_F^\times} = 1, \varepsilon \mu^2 = \omega} \text{tr} (\pi(\chi) \rtimes \mu) (f \times \varepsilon) M_{P_\alpha}(s_{\alpha+\beta}, 0).$
- $\frac{1}{4} \sum_{\chi \in \widehat{C_E}; \chi|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega} \text{tr} (1 \rtimes \pi(\chi)) (f \times \varepsilon) M_{P_\beta}(s_{s\alpha+\beta}, 0).$
- $\frac{1}{2} \sum \text{tr} (\mu \rtimes \pi(\theta)) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0).$
 The sum is over the nontrivial quadratic characters μ of C_F and cuspidal monomial representations $\pi(\theta)$ of $\text{GL}(2, \mathbb{A}_F)$, where θ is a character of $C_{E_{\mu\varepsilon}}$, such that: $\pi(\theta)$ is not E_μ -monomial and $\theta|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$.
- $\frac{1}{4} \sum \text{tr} (\mu \rtimes \pi(\theta)) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0).$
 The sum is over the nontrivial quadratic characters μ of C_F and cuspidal monomial representations $\pi(\theta)$ of $\text{GL}(2, \mathbb{A}_F)$, where θ is a character of $C_{E_{\mu\varepsilon}}$, such that $\theta|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega$ and $\pi(\theta)$ is E -monomial. In particular, we have $\frac{\sigma' \theta}{\theta} = \mu$, where σ' is the generator of $\text{Gal}(E_{\mu\varepsilon}/F)$.
- $\frac{1}{2} \sum \text{tr} (\varepsilon \rtimes \pi) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0).$
 The sum is over all cuspidal non- E -monomial, or one dimensional, automorphic representations π of $\text{GL}(2, \mathbb{A}_F)$ such that $\varepsilon \omega_\pi = \omega$.
- $\frac{1}{4} \sum \text{tr} (\varepsilon \rtimes \pi) (f \times \varepsilon) M_{P_\beta}(s_{2\alpha+\beta}, 0).$
 The sum is over all cuspidal E -monomial automorphic representations π of $\text{GL}(2, \mathbb{A}_F)$ such that $\varepsilon \omega_\pi = \omega$.

The stable spectral expansion $SI^d(H_1, f_1)$ is equal to:

$$\begin{aligned} \sum_{\substack{\tau \in L_d(\mathrm{GL}(2, \mathbb{A}_F)), \\ \tau \neq \varepsilon \tau, \eta|_{\mathbb{A}_F^\times} \cdot \omega_\tau = \omega}} \mathrm{tr} (\tau \otimes_1 \eta) (f_1) + \frac{1}{2} \sum_{\substack{\theta \neq {}^\sigma \theta, \\ (\theta \eta)|_{\mathbb{A}_F^\times} \cdot \varepsilon = \omega}} \mathrm{tr} (\pi(\theta) \otimes_1 \eta) (f_1) \\ + \frac{1}{4} \sum_{\eta|_{\mathbb{A}_F^\times} = \omega} \mathrm{tr} (I(1, 1) \otimes_1 \eta) (f_1), \end{aligned}$$

and $I^d(H_2, f_2)$ is equal to:

$$\sum_{\pi \in L_d(\mathbf{H}_2(\mathbb{A}_F), \omega)} \mathrm{tr} \pi(f_2) + \frac{1}{4} \sum_{\chi^2 = \omega \circ N_{E/F}} \mathrm{tr} (I(\chi, \chi) \otimes_2 \omega) (f_2).$$

3.2. Comparison of Geometric Sides of Trace Formulas

3.2.1. Weighted Orbital Integrals. Let $\mathbf{H}$ be a reductive F -group. Let $\mathbf{Z}_0$ be the split component of the center $\mathbf{Z}$ of $\mathbf{H}$. Let ε be a character of $\mathbf{H}$ trivial on $\mathbf{Z}$. Fix a minimal parabolic subgroup $\mathbf{P}_0$ of $\mathbf{H}$. Let $\mathbf{A}_0$ be the split component of the Levi subgroup of $\mathbf{P}_0$. Put $\mathfrak{a}_0 := X_*(\mathbf{A}_0) \otimes_{\mathbb{Z}} \mathbb{R}$.

We say that a regular element in $\mathbf{H}(F)$ is **parabolic** if it is not elliptic; in other words, its centralizer in $\mathbf{H}(F)$ is a maximal torus which is contained in a proper (F -)parabolic subgroup. The terms in the geometric side of the (ε -twisted) trace formula associated with the semisimple conjugacy classes of parabolic elements are **ε -twisted-weighted orbital integrals**, defined as follows:

Suppose γ is a parabolic element in $\mathbf{H}(F)$. Let $\mathbf{M}$ be the minimal Levi subgroup of $\mathbf{H}$ containing γ . Let $\mathbf{H}(\mathbb{A}_F)_\gamma$ be the centralizer of γ in $\mathbf{H}(\mathbb{A}_F)$. The weighted orbital integral associated with the semisimple conjugacy class of γ has the following form ([CLL, Lecture 5]):

$$O^T(\gamma, f) = c(\gamma) \int_{\mathbf{H}(\mathbb{A}_F)_\gamma \backslash \mathbf{H}(\mathbb{A}_F)} f(h^{-1} \gamma h) \varepsilon(h) v(h, T) dh.$$

Here, $c(\gamma)$ is some constant dependent on γ , T is the truncation parameter in $\mathfrak{a}_0$, and $v(h, T)$ is the volume of a certain convex hull in $\mathfrak{a}_H \backslash \mathfrak{a}_M$ determined by the projections of T and $H_{P_0}(h)$ onto $\mathfrak{a}_H \backslash \mathfrak{a}_M$.

Let n be the real dimension of $\mathfrak{a}_H \backslash \mathfrak{a}_M$. For each T , the volume function $v(h, T)$ is a degree n polynomial in the coordinates of $H_{P_0}(h)$. Let $\{h_1, h_2, \dots\}$ be the eigenvalues of h . The coordinates of $H_{P_0}(h) \in \mathfrak{a}_M$ are expressed in terms of $\log |h'_i|$, where $h'_i = N_{L_i/F} h_i$, L_i is the smallest extension field containing h_i , and $|\cdot|$ is the absolute value function on $\mathbb{A}_F$. For any place v of F , let $|\cdot|_v$ be the p -adic absolute value on F_v . Then, $|h_i| = \prod_v |h_i|_v$, and $\log |h'_i| = \sum_v \log |h'_i|_v$. We rewrite $O^T(\gamma, f)$ as follows:

$$(3.1) \quad c(\gamma) \sum_S \left[\int_{H_S, \gamma \backslash H_S} f_S(h_S^{-1} \gamma h_S) \varepsilon_S(h_S) v(h, T, S) dh_S \right. \\ \left. \times \prod_{w \notin S} \int_{H_w, \gamma \backslash H_w} f_w(h_w^{-1} \gamma h_w) \varepsilon_w(h_w) dh_w \right].$$

Here, the sum is over the sets S of n distinct places of F . The subscript S in H_S, f_S, ε_S signifies the product of local components H_v, f_v, ε_v over the places v in S . The term

$v(h, T, S)$ is a polynomial in the local components of $v(h, T)$ at the places in S . Observe that the product over $w \notin S$ consists of non-weighted local ε_w -twisted orbital integrals.

We say that a local function f_v on $\mathbf{H}(F_v)$ is **elliptic** if the orbital integral of f_v vanishes at the parabolic elements in $\mathbf{H}(F_v)$. Matrix coefficients of cuspidal representations of $\mathbf{H}(F_v)$ are examples of such functions. By choosing test functions f whose local components at $n + 1$ distinct places are elliptic, the weighted orbital integral $O^T(\gamma, f)$ vanishes.

We say that an element in $\mathbf{H}(F)$ is **singular** if it has two equal eigenvalues. From [A2], the weighted orbital integral at a singular element may be expressed as a limit of a linear combination of weighted orbital integrals at parabolic elements. Hence, by choosing f with $n + 1$ elliptic components with n sufficiently large, the weighted orbital integrals at singular elements also vanish. Here, n is sufficiently large if it is greater than or equal to the semisimple rank of $\mathbf{H}$.

Consider the case of $\mathbf{G} = \mathrm{GSp}(2)$. Suppose γ is a parabolic element in $\mathbf{G}(F)$ for which the ε -twisted orbital integral is nonzero. Then, the centralizer of γ in $\mathbf{G}(\mathbb{A}_F)$ must necessarily lie in the kernel of ε . The centralizer of γ in $\mathbf{G}(\mathbb{A}_F)$ is a maximal non-elliptic torus. The following is a list of representatives of the conjugacy classes of the maximal non-elliptic tori in $\mathbf{G}(\mathbb{A}_F)$:

- (1) $\mathbf{T}(\mathbb{A}_F) = \{\mathrm{diag}(a, b, \lambda/b, \lambda/a) : a, b, \lambda \in \mathbb{A}_F^\times\};$
- (2) $\mathbf{T}(\mathbb{A}_F) = \left\{ \begin{pmatrix} t_2 & \\ & \frac{\lambda}{\det t_2} e t_2 e \end{pmatrix} : t_2 \in \mathbf{T}_2(\mathbb{A}_F), \lambda \in \mathbb{A}_F^\times \right\},$
where $\mathbf{T}_2$ is a maximal elliptic torus in $\mathrm{GL}(2)$ and $e = \mathrm{diag}(1, -1);$
- (3) $\mathbf{T}(\mathbb{A}_F) = \left\{ \begin{pmatrix} a & \\ & t_2 \\ & & b \end{pmatrix} : a, b \in \mathbb{A}_F^\times, t_2 \in \mathbf{T}_2(\mathbb{A}_F), ab = \det t_2 \right\},$
where $\mathbf{T}_2$ is a maximal elliptic torus in $\mathrm{GL}(2)$.

In the first two cases, the similitude factors of the elements in the torus range over all of $\mathbb{A}_F^\times$; hence these tori do not lie in the kernel of ε . In the third case, the similitude factor is $\det t_2$. An elliptic maximal torus in $\mathrm{GL}(2, \mathbb{A}_F)$ which lies in the kernel of $\varepsilon \circ \det$ is conjugate to $\left\{ \begin{pmatrix} c & dA \\ d & c \end{pmatrix} : c, d \in \mathbb{A}_F, (c, d) \neq (0, 0) \right\}$, where A is an element in $F^\times - F^{\times 2}$ such that $E = F(\sqrt{A})$. Consequently, if the centralizer of γ must be conjugate to

$$\mathbf{T}(\mathbb{A}_F) = \left\{ \begin{pmatrix} a & & \\ & c & dA \\ & d & c \\ & & & b \end{pmatrix} : a, b \in \mathbb{A}_F^\times, c, d \in \mathbb{A}_F, ab = c^2 - d^2 A \right\}.$$

Since the semisimple rank of $\mathbf{T}$ is 1, using test functions with two elliptic components suffices to eliminate the weighted orbital integrals from the geometric side of the ε -twisted trace formula for $\mathbf{G}$.

As for the ε -endoscopic groups $\mathbf{H}_1, \mathbf{H}_2$, their semisimple F -ranks are equal to 1. Hence, using test functions with two elliptic components suffices to eliminate the weighted orbital integrals.

Fix two places u_1, u_2 of F which are prime in E . Let $E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$ denote the space of functions in $C(\mathbf{G}(\mathbb{A}_F), \omega)$ whose local components at u_1, u_2 are elliptic. For $i = 1, 2$, define $E(u_1, u_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$ likewise for $\mathbf{H}_i$. Since elliptic regular elements in $\mathbf{H}_i(F_v)$ are norms of elliptic regular elements in $\mathbf{G}(F_v)$ for any place v which is prime in E , it is reasonable to expect that, for any function f in $E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$, there exists matching functions $f_i \in E(u_1, u_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$.

Let $\mathbf{H} = \mathbf{G}, \mathbf{H}_1$ or $\mathbf{H}_2$. Let $\{\mathcal{O}\}_e$ denote the set of F -conjugacy classes of elliptic regular elements in $\mathbf{H}(F)$. For each $\mathcal{O} \in \{\mathcal{O}\}_e$, put

$$J_{\mathcal{O}}(f) := O_{\gamma}(f) = \int_{\mathbf{H}(\mathbb{A}_F)_{\gamma} \backslash \mathbf{H}(\mathbb{A}_F)} f(h^{-1}\gamma h) \varepsilon_H(h) dh/du,$$

where γ is a representative in $\mathbf{H}(F)$ of the conjugacy class $\mathcal{O}$, and ε_H is ε if $\mathbf{H} = \mathbf{G}$, 1 if $\mathbf{H} = \mathbf{H}_i$ ($i = 1, 2$). Here, dh, du are Tamagawa measures on $\mathbf{H}(\mathbb{A}_F)_\gamma$ and $\mathbf{H}(\mathbb{A}_F)$, respectively.

3.1. PROPOSITION. *Let f be a function in $E(u_1, u_2, \mathbf{H}(\mathbb{A}_F), \omega)$. Then, the geometric side of the ε_H -twisted trace formula for $\mathbf{H}$ is equal to*

$$\sum_{\mathcal{O} \in \{\mathcal{O}\}_e} J_{\mathcal{O}}(f).$$

PROOF. This follows from [CLL, Lecture 5] and the previous discussion. $\square$

3.2.2. Kottwitz-Shelstad's Formula. Let $i = 1$ or 2 . For any strongly $\mathbf{G}$ -regular element $\delta \in \mathbf{H}_i(F)$ and test function $f_i \in C(\mathbf{H}_i(\mathbb{A}_F), \omega)$, let

$$SO_\delta(f_i) = \sum_{\{\delta'\}} O_{\delta'}(f_i),$$

where the sum is over a set of representatives for the $\mathbf{H}_i(\mathbb{A}_F)$ -conjugacy classes of elements $\delta' \in \mathbf{H}_i(\mathbb{A}_F)$ in the $\mathbf{H}_i(\mathbb{A}_F)$ -conjugacy class of δ .

If we have an equation relating the geometric sides of the trace formulas for $\mathbf{G}$ and its ε -endoscopic groups $\mathbf{H}_1, \mathbf{H}_2$, then we may deduce a relation of the spectral sides. The tool with which we use to relate the geometric sides of the trace formulas is the following formula of Kottwitz-Shelstad's in [KS, Sect. 7.4] (applied to our situation):

$$(3.2) \quad T_e(G, f) = \iota(G, H_1) ST_e^{**}(H_1, f_1) + \iota(G, H_2) ST_e^{**}(H_2, f_2),$$

where:

- f and f_i ($i = 1, 2$) are matching functions in $C(\mathbf{G}(\mathbb{A}_F), \omega)$ and $C(\mathbf{H}_i(\mathbb{A}_F), \omega)$, respectively;
- $T_e(G, f) = \sum_{\mathcal{O} \in \{\mathcal{O}\}_e} J_{\mathcal{O}}(f)$;
- $ST_e^{**}(H_i, f_i) := a_{\mathbf{H}_i} \sum_{\{\delta_i\}} SO_{\delta_i}(f_i)$, where the sum is over a set of representatives for the $\mathbf{H}_i(\bar{F})$ -conjugacy classes of elliptic $\mathbf{G}$ -regular elements $\delta_i \in \mathbf{H}_i(F)$;
- For any algebraic F -group $\mathbf{H}$,

$$a_{\mathbf{H}} := \left| \pi_0 \left(Z(\hat{H})^\Gamma \right) \right| \cdot \left| \ker^1(F, Z(\hat{H})) \right|^{-1};$$

- For $i = 1, 2$,

$$\iota(G, H_i) := a_{\mathbf{G}} |\text{Out}(\mathbf{H}_i, \mathcal{H}_i, s_i, \xi_i)|^{-1} \cdot a_{\mathbf{H}_i}^{-1},$$

where $\text{Out}(\mathbf{H}_i, \mathcal{H}_i, s_i, \xi_i)$ is the group of outer automorphisms of $\mathbf{H}_i$ determined by automorphisms of the endoscopic data $(\mathbf{H}_i, \mathcal{H}_i, s_i, \xi_i)$ (see [KS, Sect. 2.1]).

The group $Z(\hat{G}) = \mathbb{C}^\times$ is connected and the action of $\text{Gal}(\bar{F}/F)$ on $Z(\hat{G})$ is trivial; hence, $a_{\mathbf{G}} = 1$. For both endoscopic groups, $\left| \pi_0 \left(Z(\hat{H}_i)^\Gamma \right) \right| = 1$; hence, $a_{\mathbf{H}_i} = \left| \ker^1(F, Z(\hat{H}_i)) \right|^{-1}$.

3.2. CLAIM. *For $i = 1, 2$, $\ker^1(F, Z(\hat{H}_i))$ is trivial.*

PROOF. By definition, for $i = 1, 2$, $\ker^1(F, Z(\hat{H}_i))$ is the kernel of the natural embedding

$$H^1(\Gamma_F, Z(\hat{H}_i)) \rightarrow \prod_v H^1(\Gamma_{F_v}, Z(\hat{H}_i)),$$

where $\Gamma_F = \text{Gal}(\bar{F}/F)$ and Γ_{F_v} is the decomposition group of v in $\text{Gal}(\bar{F}/F)$. Observe that the action of Γ_F on $Z(\hat{H}_i)$ factors through $\text{Gal}(E/F) = \langle \sigma \rangle$. Let $\Gamma_E = \text{Gal}(\bar{F}/E) \subset \Gamma_F$, then $H^1(\Gamma_E, Z(\hat{H}_i)) = \text{Hom}_{\text{cts}}(\Gamma_E, Z(\hat{H}_i))$ because Γ_E acts trivially on $Z(\hat{H}_i)$.

For all but finitely many places w of E , w either lies above a place v of F which is prime and unramified in E , or it lies above a place v of F which splits into two places w, w' of E . In the former case, we have $\text{Gal}(E_w/F_v) = \text{Gal}(E/F)$. In the later, we have $\text{Gal}(E_w/F_v) = \text{Gal}(E_{w'}/F_v) = 1$.

Given any $x \in \ker^1(F, Z(\hat{H}_i)) \subset H^1(\Gamma_F, Z(\hat{H}_i))$, the restriction $x|_{\Gamma_E}$ of x to Γ_E is a (continuous) homomorphism from Γ_E to $Z(\hat{H}_i)$. By assumption, the image of x in $\prod_v H^1(\Gamma_{F_v}, Z(\hat{H}_i))$ is trivial, which means, in particular, that the image of $x|_{\Gamma_E}$ in $H^1(\Gamma_{E_w}, Z(\hat{H}_i))$ is trivial for all but finitely many places w of E . Since $x|_{\Gamma_E}$ is a continuous homomorphism, we conclude that $x|_{\Gamma_E} = 1$.

Fix an element $\sigma_0 \in \Gamma_F - \Gamma_E$. Then, any $x \in \ker^1(F, Z(\hat{H}_i))$ produces a cocycle $y \in H^1(\text{Gal}(E/F), Z(\hat{H}_i))$ defined by $y_\sigma := x_{\sigma_0 \sigma}$. Since $x|_{\Gamma_E} = 1$, y is independent of the choice of σ_0 . Pick a place v of F which remains prime in E . Since x is an element of $\ker^1(F, Z(\hat{H}_i))$, the image of y in $H^1(\text{Gal}(E_v/F_v), Z(\hat{H}_i))$ is trivial. But, $H^1(\text{Gal}(E_v/F_v), Z(\hat{H}_i))$ is equal to $H^1(\text{Gal}(E/F), Z(\hat{H}_i))$ because v is prime in E , so y itself is trivial. The claim follows. $\square$

3.3. CLAIM. $|\text{Out}(\mathbf{H}_i, \mathcal{H}_i, s_i, \xi_i)| = 2$ for $i = 1, 2$.

PROOF. For $i = 1, 2$, the only nontrivial element of $\text{Out}(\mathbf{H}_i, \mathcal{H}_i, s_i, \xi_i)$ comes from the action of $\sigma \in \text{Gal}(E/F)$ on $\hat{H}_i$. $\square$

3.4. COROLLARY. *The following holds for matching functions:*

$$(3.3) \quad T_e(G, f) = \frac{1}{2} \sum_{\{\delta_1\}} SO_{\delta_1}(f_1) + \frac{1}{2} \sum_{\{\delta_2\}} SO_{\delta_2}(f_2)$$

where the sums are over stable conjugacy classes of elliptic $\mathbf{G}$ -regular elements in the F -points of the ε -endoscopic groups.

3.3. Application of Kottwitz-Shelstad's Formula

Recall, for $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$, $I^d(G, f, \varepsilon)$ denotes the ε -discrete part of the fine χ -expansion of $\mathbf{G}$, $I^c(G, f, \varepsilon)$ denotes the continuous part of the expansion, and $I^d(H_i, f_i)$, $I^c(H_i, f_i)$ ($i = 1, 2$) denote analogous objects for $\mathbf{H}_i$.

Fix a two places u_1, u_2 of F which are prime in E . By Proposition 3.1 and the trace formula (3.1), for any function f in $E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$, we have

$$T_e(G, f) = I^d(G, f, \varepsilon) + I^c(G, f, \varepsilon).$$

Since $\mathbf{H}_2$ is a quotient of $\text{R}_{E/F}\text{GL}(2) \times \mathbb{G}_m$, there is no distinction between stable and ordinary conjugacy classes. For any function f_2 in $E(u_1, u_2, \mathbf{H}_2(\mathbb{A}_F), \omega)$ which matches a function in $E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$, the orbital integral of f_2 is zero at any element in $\mathbf{H}_2(F)$ which is not elliptic G -regular; hence, for such f_2 we have

$$ST_e^{**}(H_2, f_2) = T_e(H_2, f_2) = I^d(H_2, f_2) + I_{H_2}^c(H_2, f_2).$$

For $\mathbf{H}_1$, where there is a distinction between stable and ordinary conjugacy, the situation is more complicated.

As in the case of $\mathbf{H}_2$, if the function $f_1 \in E(u_1, u_2, \mathbf{H}_1(\mathbb{A}_F), \omega)$ matches some $f \in E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$, then the orbital integral of f_1 is zero at all elliptic regular elements which are not $\mathbf{G}$ -regular. Hence, $ST_e^{**}(H_1, f_1) = ST_e(H_1, f_1)$, where $ST_e(H_1, f_1)$ is the sum of stable orbital integrals over the stable conjugacy classes of elliptic regular elements in $\mathbf{H}_1(F)$.

Applying Kottwitz-Shelstad's formula in the context of standard endoscopy for $\mathbf{H}_1$, we have:

$$(3.1) \quad I^d(H_1, f_1) + I^c(H_1, f_1) = T_e(H_1, f_1) = ST_e(H_1, f_1) + \frac{1}{2}T_e^\kappa(H_1, f_1),$$

where $T_e^\kappa(H_1, f_1)$ is a sum over stable conjugacy classes in $\mathbf{H}(F)$ of κ -orbital integrals $O_{\delta_{H_1}}^\kappa(f_1)$. In this case, there is only one nontrivial κ (in fact, $\kappa = \varepsilon$), and it is associated with a unique (up to equivalence) standard endoscopic group $\mathbf{T}_1$ of $\mathbf{H}_1$. In particular, $\mathbf{T}_1$ is the F -torus whose F -points are:

$$\mathbf{T}_1(F) = \{(x, y) \in E^\times \times E^\times : N_{E/F}x = N_{E/F}y\}.$$

We rewrite (3.1) as:

$$I^d(H_1, f_1) - \frac{1}{2}T_e^\kappa(H_1, f_1) + I^c(H_1, f_1) = ST_e(H_1, f_1).$$

Recall from Section 3.1.3.3 that $SI^d(H_1, f_1)$ denotes the stable part of the discrete spectral expansion $I^d(H_1, f_1)$. From the theory of [LL], $\frac{1}{2}T_e^\kappa(H_1, f_1)$ is equal to the unstable part of $I^d(H_1, f_1)$; thus, $I^d(H_1, f_1) - \frac{1}{2}T_e^\kappa(H_1, f_1)$ is equal to $SI^d(H_1, f_1)$.

We conclude that the following equation holds for any f_1 in $E(u_1, u_2, \mathbf{H}_1(\mathbb{A}_F), \omega)$ matching some function in $E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$:

$$(3.2) \quad ST_e(H_1, f_1) = SI^d(H_1, f_1) + I^c(H_1, f_1).$$

In conclusion, the following equations hold for $f \in E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and matching functions f_i in $E(u_1, u_2, \mathbf{H}_i, \omega)$ ($i = 1, 2$):

$$(3.3) \quad I^d(G, f, \varepsilon) + I^c(G, f, \varepsilon) = \frac{1}{2} \left(SI^d(H_1, f_1) + I^c(H_1, f_1) + I^d(H_2, f_2) + I^c(H_2, f_2) \right).$$

CHAPTER 4

Global Lifting

4.1. The ε -Trace Identity

4.1.1. Some Notation and Terminology.

4.1.1.1. *Global Data.* Let F be a number field. Let $\mathbf{H}$ be a reductive group over F with L -group ${}^L H$. Let V be the set of places of F . Let S be a finite set of places of V . Let $\{c_v : v \in V - S\}$ be a set of conjugacy classes in ${}^L H$.

Define a (countably-)infinite-tuple as follows:

$$\mathcal{C}(S, {}^L H) := [c_v]_{v \in V - S} \subset \prod_{v \in V - S} {}^L H.$$

We call $\mathcal{C}(S, {}^L H)$ a **global datum** in ${}^L H$. For simplicity, we sometimes replace ${}^L H$ by a finite Galois form. For example, if $\mathbf{H}$ is split over F , then we use $\hat{H}$ instead.

For any automorphic representation π of $\mathbf{H}(\mathbb{A}_F)$, there exists a finite set of places S such that π_v is unramified for all $v \notin S$. For each $v \notin S$, denote by $c(\pi_v)$ the Frobenius-Hecke (abbrev. F-H) class in ${}^L H$ which parametrizes π_v (see [Bo]). Then, π defines the global datum

$$\mathcal{C}(\pi, S, {}^L H) := [c(\pi_v)]_{v \in V - S} \subset \prod_{v \in V - S} {}^L H.$$

We call $\mathcal{C}(\pi, S, {}^L H)$ a **global datum** (or the S -**global datum**) of π . The global datum $\mathcal{C}(\pi, S, {}^L H)$ is **defined** if and only if π_v is unramified for all $v \notin S$. Observe that if $\mathcal{C}(\pi, S, {}^L H)$ is defined, then so is $\mathcal{C}(\pi, S', {}^L H)$ for any finite subset S' of V containing S .

4.1.1.2. *Global Packets.* Let $\{\{\Pi_v\} : v \in V\}$ be a collection of local (quasi-)packets $\{\Pi_v\}$ of $\mathbf{G}(F_v)$ such that, for almost all (finite) v , $\{\Pi_v\}$ contains a unique unramified representation Π_v^0 . We define the **(quasi-)packet** $\{\Pi\}$ to be the restricted tensor product

$$\otimes_{v \in V} \{\Pi_v\} := \left\{ \otimes_{v \in V} \Pi'_v : \Pi'_v \in \{\Pi_v\} \text{ for all } v, \Pi'_v = \Pi_v^0 \text{ for almost all } v \right\}.$$

For any automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$, we denote by $\{\Pi\}$ the global packet to which Π belongs. Under this notation, $\{\Pi\} = \{\Pi'\}$ if $\Pi' \in \{\Pi\}$.

We say that a (quasi-)packet of representations of $\mathbf{G}(\mathbb{A}_F)$ is a **discrete spectrum (quasi-)packet** if it contains a discrete spectrum representation. We say that a discrete spectrum packet of $\mathbf{G}(\mathbb{A}_F)$ is ε -**invariant** if it contains an ε -invariant discrete spectrum representation. We say that a discrete spectrum packet $\{\Pi\}$ is **stable** if every member of $\{\Pi\}$ occurs with the same multiplicity in the discrete spectrum of $\mathbf{G}(\mathbb{A}_F)$. Otherwise, we say that $\{\Pi\}$ is **unstable**.

Let $\{\Pi\} = \otimes_{v \in V} \{\Pi_v\}$ be a (quasi-)packet of representations of $\mathbf{G}(\mathbb{A}_F)$. Let S be a finite set of places such that, for all $v \notin S$, the local packet $\{\Pi_v\}$ contains a unique unramified representation Π_v^0 . For $v \notin S$, let $c(\{\Pi_v\})$ denote the F-H class in $\hat{G}$ which parametrizes Π_v^0 . Let $\{\Pi\}_S$ denote the subset of $\{\Pi\}$ of representations Π such that $\Pi_v = \Pi_v^0$ for all $v \notin S$. We call $\{\Pi\}_S$ the **S -part** of $\{\Pi\}$. In general, the cardinality of a

global packet may be infinite, but the cardinality of $\{\Pi\}_S$ is finite. Put $\mathcal{C}(\{\Pi\}, S, \hat{G}) := [c(\{\Pi_v\})]_{v \in V-S} \subset \prod_{v \notin S} \hat{G}$.

Let $V^{\text{un}}(E/F)$ denote the set of finite places of F which are unramified in E . Let $\{\pi_i\} = \otimes_{v \in V} \{\pi_{i,v}\}$ be a global (quasi-)packet of $\mathbf{H}_i(\mathbb{A}_F)$, where $i = 1$ or 2 . Let S be a finite set of places containing $V - V^{\text{un}}(E/F)$ such that the S -part $\{\pi_i\}_S$ of $\{\pi_i\}$ is nonempty. Let ${}^L H_i$ denote the finite Galois form $\hat{H}_i \rtimes \text{Gal}(E/F)$ of the L -group of $\mathbf{H}_i$. Then, $\{\pi_i\}$ defines the global datum

$$\mathcal{C}(\{\pi_i\}, S, {}^L H_i) = [c(\{\pi_{i,v}\})]_{v \notin S} \subset \prod_{v \notin S} {}^L H_i.$$

As we shall discuss in Section 4.2, each $c(\{\pi_{i,v}\}) \subset {}^L H_i$ lifts via the L -group embedding $\xi_i : {}^L H_i \hookrightarrow \hat{G}$ to a conjugacy class $\xi_i(c(\{\pi_{i,v}\}))$ in $\hat{G}$. We obtain the following global datum:

$$\mathcal{C}(\xi_i(\{\pi_i\}), S, \hat{G}) := [\xi_i(c(\{\pi_{i,v}\}))]_{v \notin S} \subset \prod_{v \notin S} \hat{G}.$$

If $\mathcal{C}(\xi_i(\{\pi_i\}), S, \hat{G})$ coincides with the global datum $\mathcal{C}(\{\Pi\}, S, \hat{G})$ for some global (quasi-)packet $\{\Pi\}$ of $\mathbf{G}(\mathbb{A}_F)$, we say that the packet $\{\pi_i\}$ **lifts** to $\{\Pi\}$ and write $\{\Pi\} = \xi_i(\{\pi_i\})$.

4.1.2. Langlands' Trick. For $f \in C(\mathbf{G}(\mathbb{A}_F))$, let $I^d(G, f, \varepsilon)$ denote the discrete part of the ε -twisted fine χ -expansion for $\mathbf{G}$. Let $I^c(G, f, \varepsilon)$ denote the continuous part of the expansion. Define $I^d(H_i, f_i)$, $I^c(H_i, f_i)$ ($i = 1, 2$) likewise (without twisting) for $\mathbf{H}_i$.

The following proposition depends on the validity of the Fundamental Lemma (see Chapter B in the Appendix).

4.1. PROPOSITION. *Fix two finite places u_1, u_2 of F which are prime in E . The following holds for matching functions f, f_1, f_2 whose components at u_1, u_2 are elliptic:*

$$(4.1) \quad I^d(G, f, \varepsilon) - \frac{1}{2} S I^d(H_1, f_1) - \frac{1}{2} I^d(H_2, f_2) = 0$$

We content ourselves with giving only a sketch of the proof, which is essentially a retelling of an argument of Langlands' in [L, Chap. 11].

Recall that $V^{\text{un}}(E/F)$ is the set of finite places of F which are unramified in E , and that ω is a fixed character of the center of $\mathbf{G}$. Let S be a finite set of places of F containing $(V - V^{\text{un}}(E/F)) \cup \{u_1, u_2\}$ such that ω_v is unramified, and $f_v, f_{1,v}, f_{2,v}$ are spherical, for all $v \notin S$.

Rewrite equation (3.3) as follows:

$$(4.2) \quad I^d(G, f, \varepsilon) - \frac{1}{2} S I^d(H_1, f_1) - \frac{1}{2} I^d(H_2, f_2) \\ = - \left(I^c(G, f, \varepsilon) - \frac{1}{2} I^c(H_1, f_1) - \frac{1}{2} I^c(H_2, f_2) \right).$$

Fix a finite place $u \notin S$ which splits in E . Regard $f_v, f_{1,v}, f_{2,v}$ as fixed for $v \notin S \cup \{u\}$, and let f_u vary in $\mathcal{H}(G_u, \omega_u)$ (hence $f_{1,u}$ and $f_{2,u}$ also vary by the matching condition). Fix the element $\text{diag}(1, 1, \lambda, \lambda) \in \hat{G} = \text{GSp}(2, \mathbb{C})$, where $\lambda = \omega_u(\varpi_u)$. Let $\mathcal{T}(G_u, \omega_u)$ be the set of conjugacy classes in $\hat{G}$ whose similitude factors are equal to $\omega_u(\varpi_u)$. Then, any class $\mathcal{C}$ in $\mathcal{T}(G_u, \omega_u)$ is represented by

$$\begin{pmatrix} x & & & \\ & y & & \\ & & \lambda y^{-1} & \\ & & & \lambda x^{-1} \end{pmatrix}$$

for some $x, y \in \mathbb{C}^\times$. If $\mathcal{C}$ corresponds to a unitarizable unramified representation of G_u , then (x, y) must lie in a compact subset X of $\mathbb{C}^2$. More precisely, using the results of [ST], it can be shown that $X = T \cup U$, where

$$T = \left\{ (|\varpi_u|^s, |\varpi_u|^t) : (s, t) \in i\mathfrak{a}_0/i\mathfrak{a}_G = i\mathbb{R} \times i\mathbb{R} \right\} \subset \mathbb{C}^2$$

(recall that $\mathfrak{a}_0 = X_*(\mathbf{A}_0) \otimes_{\mathbb{Z}} \mathbb{R}$, where $\mathbf{A}_0$ is the maximal diagonal torus of $\mathbf{G}$), and U is a compact nondiscrete subset of a real plane (dependent on the choice of a, b) in $\mathbb{C}^2$.

Conversely, any element (x, y) in X gives rise to a conjugacy class in $\hat{G}$ which corresponds to a unitarizable unramified representation.

The Hecke algebra $\mathcal{H}(G_u, \omega_u)$ yields via Satake transform the algebra $\mathcal{H}^\vee$ of Laurent series ϕ in x, y with the property that

$$(4.3) \quad \phi(x, y) = \phi(x, \lambda y^{-1}) = \phi(y, x).$$

The above property holds because the Satake transforms of Hecke functions are defined on the set of conjugacy classes in $\hat{G}$.

Let $C(X)'$ denote the space of complex valued continuous functions ϕ on X satisfying (4.3). By the Stone-Weierstrass theorem, $\mathcal{H}^\vee$ is dense in $C(X)'$ under the sup-norm on $C(X)'$.

To prove the proposition, we use each side of equation (4.2) to define a functional on $C(X)'$. We then argue that these two functionals must be identically zero.

4.2. CLAIM. *The left hand side of equation (4.2) defines a continuous linear functional on $C(X)'$ which is given by an atomic measure on X .*

PROOF. A nonzero term in $I^d(G, f, \varepsilon)$, with corresponding quadruple $(\mathbf{M}, \mathbf{G}, \tau, s)$, has the form:

$$(4.4) \quad c_{(\mathbf{M}, \mathbf{G}, \tau, s)} \operatorname{tr} I_{P, \tau}(f) M_P^\varepsilon(s, 0) \\ = c_{(\mathbf{M}, \mathbf{G}, \tau, s)} \prod_{v \notin S} \operatorname{tr} I_{P, \tau_v}(f_v) M_P^\varepsilon(s, 0)_v \cdot \prod_{v \in S} \operatorname{tr} I_{P, \tau_v}(f_v) M_P^\varepsilon(s, 0)_v,$$

where $c_{(\mathbf{M}, \mathbf{G}, \tau, s)}$ is a constant dependant on $(\mathbf{M}, \mathbf{G}, \tau, s)$. Since by assumption f_v is spherical for all $v \notin S$, the representations I_{P, τ_v} must be unramified, or else (4.4) is equal to zero. For $v \notin S$, the space of $\mathbf{G}(\mathcal{O}_v)$ -fixed vectors in I_{P, τ_v} is one dimensional; hence, we can (and do) normalize $M_P^\varepsilon(s, 0)_v$ to be 1. Let $\mathcal{C}_{\tau_v}$ ($v \notin S$) be the F-H class in $\hat{G}$ which parametrizes I_{P, τ_v} . Then (4.4) may be written as

$$c_{(\mathbf{M}, \mathbf{G}, \tau, s)} \prod_{v \notin S} f_v^\vee(\mathcal{C}_{\tau_v}) \cdot \prod_{v \in S} \operatorname{tr} I_{P, \tau_v}(f_v) M_P^\varepsilon(s, 0)_v,$$

where $f_v^\vee$ is the Satake transform of f . Analogous statements hold for $\mathbf{H}_1$ and $\mathbf{H}_2$, with ε replaced with 1.

Through the embedding of L -groups, every conjugacy class $\mathcal{C}_i$ in ${}^L H_i$ ($i = 1, 2$) lifts to a conjugacy class $\mathcal{C}$ in $\hat{G}$. Moreover, if $\mathcal{C}_i$ corresponds to a unitarizable unramified representation of $H_{i, v}$, then the unramified representation of G_v parametrized by $\mathcal{C}$ must also be unitarizable (see Sect. 4.2, [ST]).

Let v be a place in $V^{\text{un}}(E/F)$. Let $f_{i, v}$ ($i = 1, 2$), f_v be functions in $\mathcal{H}(H_{i, v}, \omega_v)$, $\mathcal{H}(G_v, \omega_v)$, respectively, such that

$$f_{i, v}^\vee(\mathcal{C}_i) = f_v^\vee(\mathcal{C})$$

for every conjugacy class $\mathcal{C}_i$ in ${}^L H_i$ which lifts to a class $\mathcal{C}$ in $\hat{G}$. By the Fundamental Lemma, such Hecke functions have matching orbital integrals.

Given any global datum $\mathcal{G} = \mathcal{C}(S, \hat{G})$, let $Q(G, \mathcal{G})$ denote the set of equivalence classes of ε -invariant, ε -discretely occurring automorphic representations of $\mathbf{G}(\mathbb{A}_F)$ whose components at each $v \notin S$ are unramified and are parametrized by the local component $\mathcal{G}_v$ of $\mathcal{G}$.

Let $Q(H_2, \mathcal{G})$ denote the set of equivalence classes of the discretely occurring automorphic representations π_2 of $\mathbf{H}_2(\mathbb{A}_F)$ such that, for all $v \notin S$, $\pi_{2,v}$ is unramified, and the F-H class in ${}^L H_2$ parametrizing $\pi_{2,v}$ lifts to $\mathcal{G}_v$.

Let $\{\pi_1\}$ be a global (quasi-)packet of representations of $\mathbf{H}_1(\mathbb{A}_F)$. For any function f_1 in $C(\mathbf{H}_1(\mathbb{A}_F), \omega)$, put

$$\mathrm{tr} \{\pi_1\}(f_1) := \sum_{\pi'_1 \in \{\pi_1\}} m(\pi'_1) \mathrm{tr} \pi'_1(f_1),$$

where $m(\pi'_1)$ is the multiplicity of π'_1 in the discrete spectrum of $\mathbf{H}_1(\mathbb{A}_F)$. Let

$$\mathrm{tr}^{\mathrm{stable}} \{\pi_1\}(f_1)$$

denote the stable part of $\mathrm{tr} \{\pi_1\}(f_1)$ (see Section 3.1.3.3). Assume that f_1 is the tensor product of local components $f_{1,v}$. Put

$$\mathrm{tr} \{\pi_{1,v}\}(f_{1,v}) := \sum_{\pi'_{1,v} \in \{\pi_{1,v}\}} \mathrm{tr} \pi'_{1,v}(f_{1,v}).$$

Suppose the members of $\{\pi_1\}$ are the constituents of $\tau \otimes_1 \eta$, where τ is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, and η is a character of C_E . By [LL], we have

$$\mathrm{tr}^{\mathrm{stable}} \{\pi_1\}(f_1) = a_{\{\pi_1\}} \prod_{v \in V} \mathrm{tr} \{\pi_{1,v}\}(f_{1,v}),$$

where $a_{\{\pi_1\}} = \begin{cases} \frac{1}{2} & \text{if } \tau \text{ is } E\text{-monomial,} \\ 1 & \text{otherwise.} \end{cases}$

Let $\bar{Q}(H_1, \mathcal{G})$ denote the collection of equivalence classes of global packets of $\mathbf{H}_1$ whose global data lift to $\mathcal{G}$.

Let

$$\begin{aligned} \alpha_{\mathcal{G}} &= \sum_{\pi \in Q(G, \mathcal{G})} m(\pi) c_{\pi} \prod_{v \in S} \mathrm{tr} \pi_v(f_v \times \varepsilon_v) M(\pi)_v \\ &\quad - \frac{1}{2} \sum_{\{\pi_1\} \in \bar{Q}(H_1, \mathcal{G})} m(\{\pi_1\}) c_{\{\pi_1\}} a_{\{\pi_1\}} \prod_{v \in S} \mathrm{tr} \{\pi_{1,v}\}(f_{1,v}) M(\{\pi_1\})_v \\ &\quad - \frac{1}{2} \sum_{\pi_2 \in Q(H_2, \mathcal{G})} m(\pi_2) c_{\pi_2} \prod_{v \in S} \mathrm{tr} \pi_{2,v}(f_{2,v}) M(\pi_2)_v. \end{aligned}$$

Here, c_{π} is the constant defined by (3.2), $M(\pi)$ is the intertwining operator $M_{\mathcal{P}}^{\varepsilon}(s, 0)$. The symbols c_{π_i} , $M(\pi_i)$ ($i = 1, 2$) are defined likewise. The factor $m(\cdot)$ denotes the multiplicity of the representation/packet.

By the Fundamental Lemma, the left hand side of equation (4.2) may be written as the sum

$$(4.5) \quad \sum_{\mathcal{G} \in \{\mathcal{G}\}} \alpha_{\mathcal{G}} \prod_{v \notin S} f_v^{\vee}(\mathcal{G}_v)$$

over the set $\{\mathcal{G}\}$ consisting of the global data $\mathcal{G} = \mathcal{C}(S, \hat{G})$ for which $\alpha_{\mathcal{G}}$ is nonzero. Note that the set $\{\mathcal{G}\}$ is countable by the definition of the discrete part of the fine χ -expansion (see Sect. 3.1.1).

Recall that we have fixed a finite place u which splits in E such that the test functions have spherical local components at u . Let $\{r_{u,j}\}_{j=1,2,\dots}$ be the set of distinct $\mathcal{G}_u$'s for $\mathcal{G} \in \{\mathcal{G}\}$. Let

$$c_j = \sum_{\{\mathcal{G} \in \{\mathcal{G}\} : \mathcal{G}_u = r_{u,j}\}} \alpha_{\mathcal{G}} \prod_{v \notin S \cup \{u\}} f_v^{\vee}(\mathcal{G}_v).$$

Then, (4.5) is equal to

$$(4.6) \quad \sum_{j=1,2,\dots} c_j f_u^{\vee}(r_{u,j}).$$

The representations which appear in the (ε) -discrete spectra for the groups have unitarizable local components at every place. Hence, the conjugacy classes $r_{u,j}$ in (4.6) correspond to elements (x_j, y_j) in the compact set X . Since $\mathcal{H}^{\vee}$ is dense in $C(X)'$, (4.6) extends to a continuous linear functional $\mathcal{D}_d$ on $C(X)'$ defined by:

$$\mathcal{D}_d(\phi) = \sum_{j=1,2,\dots} c_j \phi(x_j, y_j), \quad \forall \phi \in C(X)'.$$

Note that the correspondence between r_j and (x_i, y_i) involves a choice of representative in $\hat{G}$ of the conjugacy class r_j . However, that choice does not affect the definition of the functional. The claim follows. $\square$

4.3. CLAIM. *The right hand side of equation (4.2) defines a sum of continuous linear functionals on $C(X)'$, each of which is defined by a measure which is absolutely continuous with respect to the Lebesgue measure on a subtorus of T .*

PROOF. The right hand side of (4.2) consists of terms in the continuous parts of the (ε) -twisted fine χ -expansions for $\mathbf{G}$, $\mathbf{H}_1$, and $\mathbf{H}_2$. Let $\mathbf{H}$ be either $\mathbf{G}$, $\mathbf{H}_1$, or $\mathbf{H}_2$. Let $\{\chi\} = (\mathbf{M}_H, \mathbf{L}_H, \tau, s)$ be the quadruple associated with a term in the continuous part of the fine χ -expansion for $\mathbf{H}$. In particular, $\mathbf{L}_H$ is a proper Levi subgroup of $\mathbf{H}$. The term associated with $\{\chi\}$ has the form:

$$(4.7) \quad c_{\{\chi\}} \int_{i\mathfrak{a}_{L_H}^*/i\mathfrak{a}_H^*} \text{tr } \mathcal{M}_{L_H}^T(P_H, \zeta) I_{P_H, \tau}(\zeta, f_H \times \varepsilon) M_{P_H}(s, \zeta) d\zeta,$$

where $c_{\{\chi\}}$ is a constant, $\mathbf{P}_H$ is the standard parabolic with Levi component $\mathbf{M}_H$, and f_H is a test function in $C(\mathbf{H}(\mathbb{A}_F), \omega)$ satisfying the hypothesis of Proposition 4.1. Here, we take ε to be 1 if $\mathbf{H}$ is one of the twisted endoscopic groups. We assume that (4.7) is nonzero. In particular, since f_u is spherical, τ_u is necessarily unramified. For $\zeta \in i\mathfrak{a}_{L_H}^*/i\mathfrak{a}_H^*$, let $\mathcal{C}_{H_u, \tau_u}(\zeta)$ denote the conjugacy class in ${}^L H$ which parametrizes $I_{P_{H,u}, \tau_u}(\zeta)$. Put $\mathcal{C}_{H_u, \tau_u} := \mathcal{C}_{H_u, \tau_u}(0)$.

Let τ_u^0 be the unramified representation of the diagonal torus $A_{0,u}$ of G_u such that $\mathcal{C}_0 := \mathcal{C}_{G_u, \tau_u^0}$ is represented by $\text{diag}(1, 1, \lambda, \lambda)$ in $\hat{G}$. For each quadruple $(\mathbf{M}, \mathbf{L}, \tau, s)$ which appears in the ε -twisted fine χ -expansion for $\mathbf{G}$ such that τ_u is unramified, there is an element ζ_{τ} in $i\mathfrak{a}_0/i\mathfrak{a}_G$ such that $I_{P_{M,u}, \tau_u}(\mathbf{P}_M)$ (being the standard parabolic with Levi component $\mathbf{M}$) is equal to $I_{P_{0,u}, \tau_u^0}(\zeta_{\tau})$ ($\mathbf{P}_0$ being the standard parabolic with Levi component $\mathbf{A}_0$). Note that, for each $\zeta_L \in i\mathfrak{a}_L/i\mathfrak{a}_G$, we have $I_{P_{M,u}, \tau_u}(\zeta_L) = I_{P_{0,u}, \tau_u^0}(\zeta_{\tau} + \zeta_L)$.

For each proper Levi subgroup $\mathbf{L}$ in $\mathbf{G}$, let $Q_G(L)$ denote the set of quadruples (associated with terms in the ε -twisted fine χ -expansion for $\mathbf{G}$) of the form $(\mathbf{M}, \mathbf{L}, \tau, s)$. As in [L, Chap. 11], it may be shown that there exists a function d_L on $Q_G(L) \times i\mathfrak{a}_L/i\mathfrak{a}_G$, integrable with respect to the second argument, such that the sum of all the terms in the ε -twisted fine χ -expansion for $\mathbf{G}$ associated with the quadruples in $Q_G(L)$ is equal to

$$\sum_{\{\chi\}=(\mathbf{M}, \mathbf{L}, \tau, s) \in Q_G(L)} \int_{i\mathfrak{a}_L^*/i\mathfrak{a}_G^*} f_u^\vee(\mathcal{C}_0(\zeta_\tau + \zeta)) d_L(\{\chi\}, \zeta) d\zeta.$$

Here, $\mathcal{C}_0(\zeta_\tau + \zeta_L) := \mathcal{C}_{G_u, \tau_u^0}(\zeta_\tau + \zeta_L)$.

For $j = 1, 2$, let $\mathbf{A}_j$ be the diagonal torus of $\mathbf{H}_j$. Let $\mathcal{C}_j$ be a conjugacy class in ${}^L H_j$ which lifts to $\mathcal{C}_0$. Let $\tau_{j,u}^0$ be the unramified representation of $A_{j,u}$ which is parametrized by $\mathcal{C}_j$. For each quadruple $(\mathbf{M}_j, \mathbf{L}_j, \tau_j, s)$ which appears in the fine χ -expansion for $\mathbf{H}_j$ such that $\tau_{j,u}$ is unramified, there is an element ζ_{τ_j} in $i\mathfrak{a}_j^*/i\mathfrak{a}_{H_j}^*$ such that $I_{P_{M_j, u}, \tau_{j,u}} = I_{P_{j,u}, \tau_{j,u}^0}(\zeta_{\tau_j})$. Here, P_{M_j} is the standard parabolic subgroup of $\mathbf{H}_j$ with Levi component $\mathbf{M}_j$, and P_j is the minimal standard parabolic subgroup of $\mathbf{H}_j$. For each proper Levi subgroup $\mathbf{L}_j$ in $\mathbf{H}_j$, let $Q_{H_j}(L_j)$ denote the set of quadruples (associated with terms in the fine χ -expansion for $\mathbf{H}_j$) of the form $(\mathbf{M}_j, \mathbf{L}_j, \tau_j, s)$. As in the case of $\mathbf{G}$, there is a function d_{j,L_j} on $Q_{H_j}(L_j) \times i\mathfrak{a}_j/i\mathfrak{a}_{H_j}$, integrable with respect to the second argument, such that the sum of all the terms in the fine χ -expansion for $\mathbf{H}_j$ associated with the quadruples in $Q_{H_j}(L_j)$ is equal to

$$\sum_{\{\chi_j\}=(\mathbf{M}_j, \mathbf{L}_j, \tau_j, s) \in Q_{H_j}(L_j)} \int_{i\mathfrak{a}_{L_j}^*/i\mathfrak{a}_{H_j}^*} f_{j,u}^\vee(\mathcal{C}_j(\zeta_{\tau_j} + \zeta)) d_{j,L_j}(\{\chi_j\}, \zeta) d\zeta.$$

For $\zeta_j \in i\mathfrak{a}_j^*/i\mathfrak{a}_{H_j}^*$ ($j = 1, 2$), let ζ_j^G denote the element in $i\mathfrak{a}_0^*/i\mathfrak{a}_G^*$ which is the lift of ζ_j via the L -group embedding $\xi_j : {}^L H_j \hookrightarrow \hat{G}$. For the image $f_j \in \mathcal{H}(H_{j,u}, \omega_u)$ of $f \in \mathcal{H}(G_u, \omega_u)$ under the map between the Hecke algebras which is dual to ξ_j , we have

$$f^\vee(\mathcal{C}_0(\zeta_j^G)) = f_j^\vee(\mathcal{C}_j(\zeta_j)), \quad \forall \zeta_j \in i\mathfrak{a}_j^*/i\mathfrak{a}_{H_j}^*.$$

By the Fundamental Lemma, f_j and f have matching orbital integrals.

Collecting terms on the right hand side of (4.2) and factoring, we obtain for each (up to conjugation) proper Levi subgroup $\mathbf{L}$ of $\mathbf{G}$, a function B_L on $Q_G(L) \times i\mathfrak{a}_L^*/i\mathfrak{a}_G^*$, integrable with respect to the second argument, such that the right hand side of (4.2) is equal to

$$(4.8) \quad \sum_{\mathbf{L}} \sum_{\{\chi\}=(\mathbf{M}, \mathbf{L}, \tau, s) \in Q_L(G)} \int_{i\mathfrak{a}_L^*/i\mathfrak{a}_G^*} f_u^\vee(\mathcal{C}_0(\zeta_\tau + \zeta)) B_L(\{\chi\}, \zeta) d\zeta.$$

The sum is over, up to conjugation, the set of proper Levi subgroups $\mathbf{L}$ of $\mathbf{G}$.

Since by the Stone-Weierstrass Theorem $\mathcal{H}^\vee$ is dense in $C(X)'$, the expression (4.8) extends to a continuous linear functional $\mathcal{D}_c$ on $C(X)'$. Moreover, $\mathcal{D}_c$ is a sum of functionals, each of which given by a measure which is absolutely continuous with respect to the Lebesgue measure on a subtorus of T . $\square$

PROOF OF PROPOSITION 4.1. By equation (4.2), we have an equality of functionals: $\mathcal{D}_d = \mathcal{D}_c$. By Claims 4.2, 4.3, and the Riesz Representation Theorem (see [Fo, Chap. 7]), we conclude that both $\mathcal{D}_d$ and $\mathcal{D}_c$ are zero. $\square$

We have shown that (4.6) is zero. By the generalized linear independence of characters (see [FK1]), we conclude that each

$$c_j = \sum_{\{\mathcal{G}: \mathcal{G}_u = r_{u,j}\}} \alpha_{\mathcal{G}} \prod_{v \notin S \cup \{u\}} f_v^\vee(\mathcal{G}_v)$$

is zero. By applying the same argument repeatedly, it can be shown that if U is any finite set of places disjoint from S , then

$$\sum \alpha_{\mathcal{G}} \prod_{v \notin S \cup U} f_v^\vee(\mathcal{G}_v) = 0,$$

where the sum is over those global data $\mathcal{G}$ with the property that $\mathcal{G}_v$ is equal to a fixed conjugacy class r_v in $\hat{G}$ for each $v \in U$. Using yet another argument of Langlands' in [L, Chap. 11], it can then be shown that each $\alpha_{\mathcal{G}}$ in (4.5) is equal to zero. Equivalently, using the notation introduced earlier, the following corollary to Proposition 4.1 holds:

4.4. COROLLARY. *Fix two places u_1, u_2 of F which are prime in E . For any*

- *finite set of places S containing $(V - V^{\text{un}}(E/F)) \cup \{u_1, u_2\}$,*
- *global datum $\mathcal{G} = \mathcal{C}(S, \hat{G}) = [c_v]_{v \in V-S}$, where the c_v 's are conjugacy classes in $\hat{G}$,*
- *matching functions $f \in E(u_1, u_2, \mathbf{G}(\mathbb{A}_F), \omega)$, $f_i \in E(u_1, u_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$ ($i = 1, 2$) whose local components at all $v \notin S$ are spherical,*

the following holds:

$$\begin{aligned} (4.9) \quad & \sum_{\pi \in Q(G, \mathcal{G})} m(\pi) c_\pi \prod_{v \in S} \text{tr } \pi_v(f_v \times \varepsilon_v) M(\pi)_v \\ &= \frac{1}{2} \sum_{\{\pi_1\} \in \bar{Q}(H_1, \mathcal{G})} m(\{\pi_1\}) c_{\{\pi_1\}} a_{\{\pi_1\}} \prod_{v \in S} \text{tr } \{\pi_{1,v}\} (f_{1,v}) M(\{\pi_1\})_v \\ & \quad + \frac{1}{2} \sum_{\pi_2 \in Q(H_2, \mathcal{G})} m(\pi_2) c_{\pi_2} \prod_{v \in S} \text{tr } \pi_{2,v}(f_{2,v}) M(\pi_2)_v. \end{aligned}$$

REMARK. Note that the above identity is of the trivial form $0 = 0$ if the global datum $\mathcal{G}$ corresponds to representations whose central characters are not equal to the character ω .

We call equation (4.9) an ε -**trace identity** with respect to S . A representation which appears nontrivially on either side of the equation is said to **contribute** to the ε -trace identity. By definition, the representations which appear nontrivially on the left hand side of (4.9) form the S -part of a global (quasi-)packet of $\mathbf{G}(\mathbb{A}_F)$. On the other hand, the representations π_i which contribute to the right hand side may comprise the S -parts of more than one global packet of $\mathbf{H}_i(\mathbb{A}_F)$. For each group $\mathbf{H}_i$, the set of all contributing representations form what we call a **multi-packet** of $\mathbf{H}_i(\mathbb{A}_F)$.

Let $\{\pi\} = \bigcup_{1 \leq j \leq n} \{\pi_j\}$ be a multi-packet of $\mathbf{H}_i(\mathbb{A}_F)$ ($i = 1$ or 2), where $\{\pi_j\}$ ($1 \leq j \leq n$) is a global (quasi-)packet of $\mathbf{H}_i(\mathbb{A}_F)$. Let $\{\pi\}_S = \bigcup_{1 \leq j \leq n} \{\pi_j\}_S$, the union of the S -parts of the global packets $\{\pi_j\}$. We call $\{\pi\}_S$ the **S -part** of the multi-packet $\{\pi\}$.

If the S -part of a global packet $\{\Pi\}$ of $\mathbf{G}(\mathbb{A}_F)$ and the S -parts of multi-packets $\{\pi_i\}$ of $\mathbf{H}_i(\mathbb{A}_F)$ ($i = 1, 2$) contribute to an ε -trace identity, we express the fact in the following table:

G	H_1	H_2
$\{\Pi\}_S$	$\{\pi_1\}_S$	$\{\pi_2\}_S$

Sometimes we drop the subscript S for brevity.

In the next section, we examine the correspondence between global data and automorphic representations. In addition, we demonstrate how the global data of the ε -endoscopic groups lift to those of $\mathbf{G}$.

4.2. Frobenius-Hecke Classes

Let V be the set of places of F . Let $V^{\text{un}}(E/F)$ be the set of places of F which are unramified in E . For each finite place v of F , fix a uniformizer ϖ_v of F_v . We sometimes drop the subscript v when the context is clear. If $v \in V^{\text{un}}(E/F)$ does not split in E , let ϖ_v also be the uniformizer of E_v . If a place v of F splits into two distinct places v_1, v_2 of E , then $E_{v_i} = F_v$ ($i = 1, 2$), and we let ϖ_v be the uniformizer of E_{v_i} .

4.2.1. Frobenius-Hecke Classes for $\mathbf{G}$. We now describe the global data for parabolically induced representations of $\mathbf{G}(\mathbb{A}_F)$. The results are well known (see [F4]) and we state them without proof.

- Let α, β, μ be characters of C_F . Let $\mathbf{T}$ be the maximal diagonal torus of $\mathbf{G}$. Let $\mathbf{P}_0$ be the minimal upper triangular parabolic subgroup of $\mathbf{G}$ containing $\mathbf{T}$. Define a representation of $\mathbf{T}(\mathbb{A}_F)$ as follows:

$$\alpha \otimes \beta \otimes \mu : \text{diag}(a, b, \lambda/b, \lambda/a) \mapsto \alpha(a)\beta(b)\mu(\lambda).$$

We extend $\alpha \otimes \beta \otimes \mu$ to a representation of $\mathbf{P}_0(\mathbb{A}_F)$ by setting it to be 1 on the unipotent component of $\mathbf{P}_0(\mathbb{A}_F)$.

Let $\alpha \times \beta \rtimes \mu$ denote the (normalizedly) parabolically induced representation $I_{\mathbf{P}_0(\mathbb{A}_F)}^{\mathbf{G}(\mathbb{A}_F)}(\alpha \otimes \beta \otimes \mu)$. Let S be a finite set of places such that α_v, β_v, μ_v are unramified for all $v \notin S$. Then,

$$\mathcal{C}(\alpha \times \beta \rtimes \mu, S, \hat{G}) = \left[\left(\begin{array}{ccc} \alpha_v \beta_v \mu_v & & \\ & \alpha_v \mu_v & \\ & \beta_v \mu_v & \\ & & \mu_v \end{array} \right) \right]_{v \notin S}.$$

Here, for any unramified character χ_v of $F_v^\times$, we also let χ_v denote the value of the character at ϖ_v .

- Let $e = \text{diag}(1, -1)$. Let $\mathbf{P}_\alpha$ be the upper triangular Siegel parabolic subgroup of $\mathbf{G}$. Its Levi component is

$$\mathbf{M}_\alpha := \left\{ \left(\begin{array}{c} g \\ \frac{\lambda}{\det g} e g e \end{array} \right) : \lambda \in \mathbb{G}_m, g \in \text{GL}(2) \right\}.$$

Let π be an automorphic representation of $\text{GL}(2, \mathbb{A}_F)$. Let μ be a character of C_F . Define a representation of $\mathbf{M}_\alpha(\mathbb{A}_F)$ as follows:

$$\pi \otimes \mu : \left(\begin{array}{c} g \\ \frac{\lambda}{\det g} e g e \end{array} \right) \mapsto \mu(\lambda)\pi(g).$$

The representation $\pi \otimes \mu$ extends to a representation of $\mathbf{P}_\alpha(\mathbb{A}_F)$. Let $\pi \rtimes \mu$ denote the normalizedly induced representation $I_{\mathbf{P}_\alpha(\mathbb{A}_F)}^{\mathbf{G}(\mathbb{A}_F)}(\pi \otimes \mu)$. Let S be a finite set of places such that μ_v, π_v are unramified for all $v \notin S$. Let $\mathcal{C}(\pi, S, \text{GL}(2, \mathbb{C})) = [c(\pi_v)]_{v \notin S}$ be the global datum in $\text{GL}(2, \mathbb{C})$ which parametrizes π . Then,

$$\mathcal{C}(\pi \rtimes \mu, S, \hat{G}) = \left[\left(\begin{array}{ccc} \mu_v \omega_{\pi_v} & & \\ & \mu_v c(\pi_v) & \\ & & \mu_v \end{array} \right) \right]_{v \notin S}.$$

Here, ω_{π_v} is the central character of π_v .

- Let $\mathbf{P}_\beta$ be the upper triangular Heisenberg parabolic subgroup of $\mathbf{G}$. Its Levi component is

$$\mathbf{M}_\beta := \left\{ \begin{pmatrix} a & & \\ & g & \\ & & \frac{\det g}{a} \end{pmatrix} : a \in \mathbb{G}_m, g \in \mathrm{GL}(2) \right\}.$$

Let μ be a character of C_F . Let π be an automorphic $\mathrm{GL}(2, \mathbb{A}_F)$ -module. Define a representation of $\mathbf{M}_\beta(\mathbb{A}_F)$ as follows:

$$\mu \otimes \pi : \begin{pmatrix} a & & \\ & g & \\ & & \frac{\det g}{a} \end{pmatrix} \mapsto \mu(a)\pi(g).$$

The representation $\pi \otimes \mu$ extends to a representation of $\mathbf{P}_\beta(\mathbb{A}_F)$. Let $\mu \rtimes \pi$ denote the normalizedly induced representation $I_{\mathbf{P}_\beta(\mathbb{A}_F)}^{\mathbf{G}(\mathbb{A}_F)}(\pi \otimes \mu)$. Let S be a finite set of places such that μ_v, π_v are unramified for all $v \notin S$. Let $\mathcal{C}(\pi, S, \mathrm{GL}(2, \mathbb{C})) = [c(\pi_v)]_{v \notin S}$ be the global datum parametrizing π . Suppose for each $v \notin S$ the conjugacy class $c(\pi_v)$ in $\mathrm{GL}(2, \mathbb{C})$ is represented by a diagonal matrix $t(\pi_v) = \mathrm{diag}(c_1(\pi_v), c_2(\pi_v)) \in \mathrm{GL}(2, \mathbb{C})$. Then, $\mathcal{C}(\pi \rtimes \mu, S, \hat{G})$ is represented by

$$\left[\begin{pmatrix} \mu_v t(\pi_v) & \\ & t(\pi_v) \end{pmatrix} \right]_{v \notin S} \in \prod_{v \notin S} \hat{G}.$$

As for the F-H classes parametrizing discrete spectrum automorphic representations of $\mathbf{G}(\mathbb{A}_F)$, we postpone their description to Section 4.3.1.

We now turn to describing the global data parametrizing the automorphic representations of the ε -endoscopic groups. The groups are nonsplit over F , and as a result the description of their global data is less straightforward. As a warmup, we first review the correspondence between F-H classes and automorphic representations in the case of $\mathrm{R}_{E/F}\mathrm{GL}(1)$, the group obtained from $\mathrm{GL}(1)$ upon restriction of scalars from E to F .

4.2.2. The Case of $\mathrm{R}_{E/F}\mathrm{GL}(1)$. Recall that E is the quadratic extension of F which corresponds via global class field theory to ε . Let σ be the generator of $\mathrm{Gal}(E/F)$. The group $E^\times$ may be identified with the group of F -points of $\mathbf{H} = \mathrm{R}_{E/F}\mathrm{GL}(1)$. In particular, $\mathbf{H}$ is the unique elliptic ε -endoscopic group of $\mathrm{GL}(2)$ (see [K]). Recall that endoscopic groups are by definition quasi-split. The L -group of $\mathbf{H}$ is ${}^L\mathbf{H} = (\mathbb{C}^\times \times \mathbb{C}^\times) \rtimes \mathrm{Gal}(E/F)$, and the Galois action is given by $\sigma(a, b) = (b, a)$ for $(a, b) \in \mathbb{C}^\times \times \mathbb{C}^\times$.

If a place $v \in V^{\mathrm{un}}(E/F)$ does not split in E , then $\mathbf{H}(F_v) = E_v^\times$. If v splits into two places v_1, v_2 of E , then $E_{v_1} = E_{v_2} = F_v$, and $\mathbf{H}(F_v) = F_v^\times \times F_v^\times$.

Let χ be a character of $E^\times \backslash \mathbb{A}_E^\times$. Consider χ as an automorphic representation of $\mathbf{H}(\mathbb{A}_F)$. We would like to compute the F-H classes in ${}^L\mathbf{H}$ which parametrize χ . The representation χ_v is unramified for almost all $v \in V^{\mathrm{un}}(E/F)$. This statement is equivalent to the following:

- (1) If v does not split in E , then $\chi_v = \chi'_v \circ N_{E/F}$ for some unramified character χ'_v of $F_v^\times$. Here, to simplify the notation, we let $N_{E/F}$ denote the norm mapping $N_{E_v/F_v} : E_v \rightarrow F_v$.
- (2) If v splits into distinct places v_1, v_2 of E , then $\chi_v = \chi_{v_1} \otimes \chi_{v_2}$, where χ_{v_i} ($i = 1, 2$) is an unramified character of $E_{v_i}^\times = F_v^\times$.

Consider case 1. To maintain consistency with the expression of its L -group, we write: $\mathbf{H}(\bar{F}) = \{(a, b) \in \mathrm{GL}(1, \bar{F}) \times \mathrm{GL}(1, \bar{F})\}$ and $\mathbf{H}(F_v) = \{(x, \sigma x) : x \in E_v^\times\}$. An unramified character χ_v of $\mathbf{H}(F_v)$ has the form

$$\chi_v = \mu_1 \otimes \mu_2 : (x, \sigma x) \mapsto \mu_1(x)\mu_2(\sigma x),$$

where $\mu_{1,v}, \mu_{2,v}$ are unramified characters of $E_v^\times$. Let $\varpi_v \in F_v$ be the uniformizer of both F_v and E_v ; this is possible because v is unramified in E . The F-H class in ${}^L H$ parametrizing χ_v is represented by (see [Bo]):

$$(\mu_1(\varpi_v), \mu_2(\sigma\varpi_v)) \rtimes \sigma = (\mu_1(\varpi_v), \mu_2(\varpi_v)) \rtimes \sigma.$$

The element $(\mu_1(\varpi_v), \mu_2(\varpi_v)) \rtimes \sigma$ is conjugate in ${}^L H$ to

$$(\lambda\mu_1(\varpi_v), \lambda^{-1}\mu_2(\varpi_v)) \rtimes \sigma$$

for any $\lambda \in \mathbb{C}^\times$. Letting $\lambda = \mu_2(\varpi_v)$, we see that the F-H class is represented by

$$t(\chi_v) = (\mu_1\mu_2(\varpi_v), 1) \rtimes \sigma = (\chi_v(\varpi_v), 1) \rtimes \sigma.$$

In particular, the F-H class parametrizing χ_v depends only on the restriction of the character to the maximal F_v -split torus $T_d = \{(x, x) : x \in F_v^\times\}$ of $\mathbf{H}(F_v)$.

In case 2, where v of F splits into v_1, v_2 of E , the group $\mathbf{H}$ is split over F_v , and the decomposition group of v in $\text{Gal}(E/F)$ is trivial. The group of F_v -points of $\mathbf{H}$ is $\mathbf{H}(F_v) = F_v^\times \times F_v^\times$. If $\chi_v = \chi_{v_1} \otimes \chi_{v_2}$ for some unramified characters χ_{v_1}, χ_{v_2} of $F_v^\times$, then the F-H class of χ_v is represented by

$$t(\chi_v) = (\chi_{v_1}(\varpi_v), \chi_{v_2}(\varpi_v)) \rtimes 1.$$

4.2.3. Classes in ${}^L H_1$. Recall from Chapter 2 that ${}^L H_1 = \hat{H}_1 \rtimes \text{Gal}(E/F)$, where

$$\hat{H}_1 = [\text{GL}(2, \mathbb{C}) \times \mathbb{C}^\times \times \mathbb{C}^\times] / \{(\text{diag}(z, z), z^{-1}, z^{-1}) : z \in \mathbb{C}^\times\}.$$

Let τ be an automorphic representation of $\text{GL}(2, \mathbb{A}_F)$. Let χ be a character of C_E . Let S be a finite set of places of F containing $V - V^{\text{un}}(E/F)$ such that τ_v is unramified for all $v \notin S$, and χ_w is unramified whenever the place w of E lies above a place $v \notin S$.

Suppose τ is parametrized by the the global datum

$$\mathcal{C}(\tau, S, \text{GL}(2, \mathbb{C})) = [c(\tau_v)]_{v \notin S},$$

where, for $v \notin S$, $c(\tau_v)$ is a conjugacy class in $\text{GL}(2, \mathbb{C})$ represented by some diagonal matrix $t(\tau_v) = \begin{pmatrix} c_{1,v} & \\ & c_{2,v} \end{pmatrix}$.

If a place $v \in V^{\text{un}}(E/F)$ remains prime in E , let v also denote the (unique) place of E which lies above v . If χ_v is unramified, then $\chi_v = \chi'_v \circ N_{E/F}$ for some unramified character χ'_v of $F_v^\times$. If $v \in V^{\text{un}}(E/F)$ splits into two distinct places v_1, v_2 of E , then $E \otimes_F F_v = F_v \oplus F_v = E_{v_1} \oplus E_{v_2}$. At such a place v , let χ_v denote the character $\chi_{v_1} \otimes \chi_{v_2}$ of $E_{v_1}^\times \oplus E_{v_2}^\times$.

4.5. CLAIM. *The local component at $v \notin S$ of the global datum $\mathcal{C}(\tau \otimes_1 \chi, S, {}^L H_1)$ parametrizing the global packet $\tau \otimes_1 \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ is as follows:*

$$\mathcal{C}(\tau \otimes_1 \chi, S, {}^L H_1)_v \ni \begin{cases} (t(\tau_v), 1, \chi'_v) \rtimes \sigma & \text{if } v \text{ is prime in } E, \\ (t(\tau_v), \chi_{v_1}, \chi_{v_2}) \rtimes 1 & \text{if } v \text{ splits in } E. \end{cases}$$

Here, for an unramified character η_v of $F_v^\times$, we also let η_v denote the value of the character at ϖ_v .

PROOF. At a place $v \notin S$ which does not split in E , we write $\mathbf{H}_1(F_v)$ in the following form:

$$\{(g, x, \sigma x) \in \text{GL}(2, F_v) \times E_v^\times \times E_v^\times : \det g = N_{E/F} x\}.$$

The character χ_v has the form $\chi_v(x) = \mu_{1,v}(x)\mu_{2,v}(\sigma x)$ for all $x \in E_v^\times$, where $\mu_{1,v}, \mu_{2,v}$ are unramified character of $E_v^\times$. The F-H class in ${}^L H_1$ parametrizing the unramified representation $\tau_v \otimes_1 \chi_v$ is represented by $(t(\tau_v), \mu_{1,v}, \mu_{2,v}) \rtimes \sigma$, which is conjugate in ${}^L H_1$ to

$$(t(\tau_v), \lambda^{-1} \mu_{1,v}, \lambda \mu_{2,v}) \rtimes \sigma$$

for any $\lambda \in \mathbb{C}^\times$. Letting $\lambda = \mu_1(\varpi_v)$, we see that the F-H class of $\tau_v \otimes_1 \chi_v$ is represented by

$$(t(\tau_v), 1, \chi_v) \rtimes \sigma.$$

The proof for the split case is similar, and we skip it. $\square$

Let $e = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \in \mathrm{GL}(2, \mathbb{C})$. Recall from Chapter 2 that the L -group embedding $\xi_1 : {}^L H_1 \rightarrow \hat{G}$ sends $1 \rtimes \sigma$ to $\begin{pmatrix} & e \\ e & \end{pmatrix}$ and $(g, a, b) \in \hat{H}_1$ to $d(g, a, b) := \begin{pmatrix} ag & \\ & bge \end{pmatrix}$.

4.6. COROLLARY. *The global datum $\mathcal{C}(\tau \otimes_1 \chi, S, {}^L H_1)$ parametrizing $\tau \otimes_1 \chi$ lifts to the global datum $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ described as follows:*

- *At a place $v \notin S$ which is prime in E , the local component at v of the global datum $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ is represented by*

$$\begin{pmatrix} \chi'^2 t(\tau_v) e & t(\tau_v) e \\ & \chi'_v c_{1,v} \end{pmatrix} \sim \begin{pmatrix} \chi'_v c_{1,v} & & & \\ & -\chi'_v c_{2,v} & & \\ & & -\chi'_v c_{1,v} & \\ & & & \chi'_v c_{2,v} \end{pmatrix} \in \mathrm{GSp}(2, \mathbb{C})$$

(Recall: $t(\tau_v) = \begin{pmatrix} c_{1,v} & \\ & c_{2,v} \end{pmatrix} \in \mathrm{GL}(2, \mathbb{C})$).

- *At a place v which splits into two places v_1, v_2 of E , the local component at v of $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ is represented by*

$$\begin{pmatrix} \chi_{v_1} t(\tau_v) & \\ & \chi_{v_2} t(\tau_v) \end{pmatrix} \in \mathrm{GSp}(2, \mathbb{C}).$$

Note that a priori all we know about $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ is that it is an infinite-tuple of conjugacy classes in $\mathrm{GSp}(2, \mathbb{C})$. At this stage we make no prediction on whether $\mathcal{C}(S, \mathrm{GSp}(\mathbb{C}))$ is the global datum of any (quasi-)packet of $\mathrm{GSp}(2, \mathbb{A}_F)$.

4.2.4. Classes in ${}^L H_2$. Recall from Chapter 2 that ${}^L H_2 = \hat{H}_2 \rtimes \mathrm{Gal}(E/F)$, where

$$\hat{H}_2 = \{(g_1, g_2, \lambda) \in \mathrm{GL}(2, \mathbb{C})^2 \times \mathbb{C}^\times : \det g_1 = \det g_2 = \lambda\},$$

and the generator σ of $\mathrm{Gal}(E/F)$ swaps the two $\mathrm{GL}(2, \mathbb{C})$ -factors.

Let $\pi_2 = \pi \otimes_2 \mu$ be an automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$, where π is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ and μ is a character of C_F such that the central character ω_π of π is equal to $\mu \circ N_{E/F}$.

If $v \in V^{\mathrm{un}}(E/F)$ is prime in E , then π_v is a representation of $\mathrm{GL}(2, E_v)$ which satisfies $\omega_{\pi_v} = \mu_v \circ N_{E/F}$. If v splits into two distinct places v_1, v_2 of E , then

$$\mathbf{H}_2(F_v) = (\mathrm{GL}(2, F_v) \times \mathrm{GL}(2, F_v) \times F_v^\times) / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in F_v^\times\},$$

and $(\pi \otimes_2 \mu)_v$ takes the form $\pi_{v_1} \otimes \pi_{v_2} \otimes_2 \mu_v$, where π_{v_i} ($i = 1, 2$) is a representation of $\mathrm{GL}(2, E_{v_i}) = \mathrm{GL}(2, F_v)$, and $\omega_{\pi_{v_1}} = \omega_{\pi_{v_2}} = \mu_v$.

Identify $\mathrm{GL}(2, \mathbb{A}_E)$ with the group of $\mathbb{A}_F$ -points of $R_{E/F}\mathrm{GL}(2)$, the F -group obtained from $\mathrm{GL}(2)$ via the restriction of scalars from E to F . The L -group (actually, a finite Galois form of the L -group) of $R_{E/F}\mathrm{GL}(2)$ is the semidirect product ${}^L R_{E/F}\mathrm{GL}(2) = \mathrm{GL}(2, \mathbb{C}) \times \mathrm{GL}(2, \mathbb{C}) \rtimes \mathrm{Gal}(E/F)$, where the action of $\sigma \in \mathrm{Gal}(E/F)$ swaps the two $\mathrm{GL}(2, \mathbb{C})$ -factors.

Let S be a finite set of places of F containing $V - V^{\text{un}}(E/F)$ such that $\pi_{2,v}$ is unramified for all $v \notin S$. Consider π as an automorphic representation of $(\text{R}_{E/F}\text{GL}(2))(\mathbb{A}_F)$. Then, π_v is unramified for all $v \notin S$. Suppose the F-H classes $\{c(\pi_v)\}_{v \notin S}$ in ${}^L\text{R}_{E/F}\text{GL}(2)$ parametrizing π are as follows:

$$c(\pi_v) \ni \begin{cases} \left(\begin{pmatrix} t_1(\pi_v) & \\ & t_2(\pi_v) \end{pmatrix}, \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \right) \rtimes \sigma & \text{if } v \text{ is prime in } E, \\ (d(\pi_{v_1}), d(\pi_{v_2})) \rtimes 1 & \text{if } v \text{ splits into } v_1, v_2 \text{ in } E, \end{cases}$$

where $t_1(\pi_v), t_2(\pi_v)$ are elements in $\mathbb{C}^\times$, and $d(\pi_{v_i})$ ($i = 1, 2$) is a diagonal matrix in $\text{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes the unramified representation π_{v_i} of $\text{GL}(2, E_{v_i})$.

4.7. CLAIM. *The global datum $\mathcal{C}(\pi_2, S, {}^L H_2)$ parametrizing the automorphic representation $\pi_2 = \pi \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ is as follows:*

- If $v \notin S$ is prime in E , then the local component $\mathcal{C}(\pi_2, S, {}^L H_2)_v$ of the global datum is represented by

$$\left(\begin{pmatrix} \mu_v^{1/2} & \\ & \mu_v^{1/2} \end{pmatrix}, \begin{pmatrix} \mu_v^{-1/2} t_1(\pi_v) & \\ & \mu_v^{-1/2} t_2(\pi_v) \end{pmatrix}, \mu_v \right) \rtimes \sigma.$$

- If $v \notin S$ splits into v_1, v_2 in E , then $\mathcal{C}(\pi_2, S, {}^L H_2)_v$ is represented by

$$(d(\pi_{v_1}), d(\pi_{v_2}), \mu_v) \rtimes 1.$$

PROOF. First, we deal with the case where v is prime in E . Write $\mathbf{H}_2(F_v)$ in the form

$$\{(g, \sigma g, x) \in \text{GL}(2, E_v)^2 \times F_v^\times\} / \{(zI_2, zI_2, N_{E/F} z^{-1}) : z \in E_v^\times\}.$$

If $\pi_{2,v} = \pi_v \otimes_2 \mu_v$ is unramified, then π_v and μ_v are unramified. In particular, π_v is the induced representation $I(\alpha, \beta)$ of $\text{GL}(2, E_v)$ for some unramified characters α, β of $E_v^\times$. The condition $\omega_\pi = \mu \circ N_{E/F}$ implies that $\omega_{\pi_v}|_{F_v^\times} = (\alpha\beta)|_{F_v^\times} = \mu_v^2$. The F-H class $c(\pi_v)$ in ${}^L\text{R}_{E/F}\text{GL}(2)$ parametrizing π is represented by

$$t(\pi_v) = \left(\begin{pmatrix} \alpha & \\ & \beta \end{pmatrix}, \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \right) \rtimes \sigma.$$

In particular, we may take $t_1(\pi_v) = \alpha(\varpi_v)$ and $t_2(\pi_v) = \beta(\varpi_v)$. Suppose the F-H class $c(\pi_{2,v})$ in ${}^L H_2$ parametrizing $\pi_{2,v}$ is represented by

$$(\text{diag}(\alpha_1, \beta_1), \text{diag}(\alpha_2, \beta_2), \mu_v) \rtimes \sigma \in {}^L H_2,$$

where α_i, β_i ($i = 1, 2$) are complex numbers. The following conditions must be satisfied:

- $\alpha_1 \alpha_2 = \alpha(\varpi_v)$,
- $\beta_1 \beta_2 = \beta(\varpi_v)$,
- $\alpha_1 \alpha_2 \beta_1 \beta_2 = \mu_v^2(\varpi_v)$.

The element $(\text{diag}(\alpha_1, \beta_1), \text{diag}(\alpha_2, \beta_2), \mu_v) \rtimes \sigma$ is conjugate in ${}^L H_2$ to

$$\left(\begin{pmatrix} \lambda_1 \alpha_1 & \\ & \lambda_2 \beta_1 \end{pmatrix}, \begin{pmatrix} \lambda_1^{-1} \alpha_2 & \\ & \lambda_2^{-1} \beta_2 \end{pmatrix}, \mu_v \right) \rtimes \sigma$$

for any $\lambda_1, \lambda_2 \in \mathbb{C}^\times$. Fix a choice of $\mu_v(\varpi_v)^{1/2}$. Letting $\lambda_1 = \mu_v(\varpi_v)^{1/2} \alpha_1^{-1}$ and $\lambda_2 = \mu_v(\varpi_v)^{1/2} \beta_1^{-1}$, we see that $c(\pi_{2,v})$ is represented by

$$\begin{aligned} & \left(\begin{pmatrix} \mu_v^{1/2} & \\ & \mu_v^{1/2} \end{pmatrix}, \begin{pmatrix} \mu_v^{-1/2} \alpha_1 \alpha_2 & \\ & \mu_v^{-1/2} \beta_1 \beta_2 \end{pmatrix}, \mu_v \right) \rtimes \sigma \\ &= \left(\begin{pmatrix} \mu_v^{1/2} & \\ & \mu_v^{1/2} \end{pmatrix}, \begin{pmatrix} \mu_v^{-1/2} \alpha & \\ & \mu_v^{-1/2} \beta \end{pmatrix}, \mu_v \right) \rtimes \sigma. \end{aligned}$$

The nonsplit case of the claim follows. Notice that the conjugacy class of the above element does not change if $\mu_v(\varpi_v)^{1/2}$ is replaced with $-\mu_v(\varpi_v)^{1/2}$.

The proof for the split case is similar, and we skip it. $\square$

Recall the notation $[(a_{ij}), (b_{ij})] := \begin{pmatrix} a_{11} & & a_{12} \\ & b_{11} & b_{12} \\ & b_{21} & b_{22} \\ a_{21} & & a_{22} \end{pmatrix}$. The L -group embedding $\xi_{2,v} : {}^L H_2 \rightarrow \hat{G}$ sends $1 \rtimes \sigma$ to $\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}$ and $(g_1, g_2, c) \rtimes 1$ to $[g_1, g_2]$.

4.8. COROLLARY. *The global datum $\mathcal{C}(\pi \otimes_2 \mu, S, {}^L H_2)$ parametrizing $\pi \otimes_2 \mu$ lifts to the global datum $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ described as follows:*

- At a place $v \notin S$ which is prime in E , $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))_v$ is represented by

$$\begin{pmatrix} & \mu_v^{1/2} & \\ \mu_v^{-1/2} t_1(\pi_v) & & \\ & & \mu_v^{-1/2} t_2(\pi_v) \\ & \mu_v^{1/2} & \end{pmatrix}.$$

- At a place v which splits into two places v_1, v_2 of E , $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))_v$ is represented by

$$[d(\pi_{v_1}), d(\pi_{v_2})].$$

As remarked earlier, at this stage we make no prediction on whether the global datum $\mathcal{C}(S, \mathrm{GSp}(2, \mathbb{C}))$ corresponds to a global (quasi-)packet of $\mathrm{GSp}(2, \mathbb{A}_F)$.

4.3. Packets

In this section, we describe the multi-packets for each ε -endoscopic group; in other words, we describe the inequivalent packets which simultaneously contribute to an ε -trace identity.

4.3.1. (Quasi-)Packets of $\mathrm{GSp}(2, \mathbb{A}_F)$. For any place v , we let ν_v be the normalized absolute value function on F_v . In other words, if v is finite, $\nu_v(x) := q_v^{-\mathrm{ord} x}$ for all $x \in F_v$, where q_v is the cardinality of the residue field of F_v , and $\mathrm{ord} x$ is the p -adic valuation of x . We let $\nu = \otimes_v \nu_v$ be the normalized absolute value function on $\mathbb{A}_F$. For any finite place v of F and unramified representation τ_v of $\mathrm{GL}(2, F_v)$, let $t(\tau_v)$ be a diagonal matrix in $\mathrm{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes τ_v .

Let 1_2 denote the trivial representation $1_{\mathrm{GL}(2, \mathbb{A}_F)}$ of $\mathrm{GL}(2, \mathbb{A}_F)$. For any place v , let $1_{2,v}$ and $\mathrm{St}_{2,v}$ denote the trivial and Steinberg representations of $\mathrm{GL}(2, F_v)$, respectively.

4.3.1.1. Unstable Packets. From [A3], the unstable (quasi-)packets of $\mathrm{GSp}(2, \mathbb{A}_F)$ fall into three types. They are described as follows:

1. Let τ_1, τ_2 be distinct cuspidal automorphic representations of the group $\mathrm{GL}(2, \mathbb{A}_F)$ such that $\omega_{\tau_1} = \omega_{\tau_2}$. Suppose $\tau_{1,v}$ and $\tau_{2,v}$ are unramified for all v outside of some finite set of places S . The packet $[\tau_1, \tau_2]$ is parametrized by the global datum (product of conjugacy classes):

$$\mathcal{C}([\tau_1, \tau_2], S, \hat{G}) \ni \left[[t(\tau_{1,v}), t(\tau_{2,v})] \right]_{v \notin S}.$$

In particular, $[\tau_1, \tau_2]$ lifts via the L -group embedding $\mathrm{GSp}(2, \mathbb{C}) \hookrightarrow \mathrm{GL}(4, \mathbb{C})$ to the induced representation $I_{(2,2)}(\tau_1, \tau_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

If the central characters of τ_1 and τ_2 are trivial, then we know from [F4, V. 10] that $[\tau_1, \tau_2]$ is the restricted tensor product $\{\Pi\} = \otimes_v \{\Pi_v\}$. Here, for all v where both τ_1 and τ_2 are square integrable, the local packet $\{\Pi_v\}$ consists of two square integrable representations Π_v^+, Π_v^- . At the rest of the places, $\{\Pi_v\}$ consists of a single representation, which we denote by Π_v^+ .

The multiplicity formula for $\Pi' \in \{\Pi\}$ is $m(\Pi') = \frac{1}{2}(1 + (-1)^{n(\Pi')})$, where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$.

REMARK. In [A3], $[\tau_1, \tau_2]$ is denoted by $(\tau_1 \boxtimes 1) \boxplus (\tau_2 \boxtimes 1)$ and is called a *Yoshida type* packet.

2. Let τ be a cuspidal automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, μ be a unitary character of C_F , such that $\omega_\tau = \mu^2$. Suppose τ_v, μ_v are unramified for all v outside of some finite set of places S . The quasi-packet $[\tau, \mu 1_2]$ is parametrized by the global datum

$$\begin{aligned} \mathcal{C}([\tau, \mu 1_2], S, \hat{G}) &\ni \left[t(\tau_v), t(\mu_v 1_{2,v}) \right]_{v \notin S} \\ &= \left[\text{blockdiag} \left(\mu_v q_v^{1/2}, t(\tau_v), \mu_v q_v^{-1/2} \right) \right]_{v \notin S}. \end{aligned}$$

Here, $\mu_v := \mu_v(\varpi_v)$.

In particular, the quasi-packet $[\tau, \mu 1_2]$ lifts to the induced representation $I_{(2,2)}(\tau, \mu 1_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

If $\mu^2 = \omega_\tau = 1$, then from [F4, V. 10] we know that $[\tau, \mu 1_2]$ is the global quasi-packet $\{\Pi\} = \otimes_v \{\Pi_v\}$, where $\{\Pi_v\} = \{\Pi_v^\times, \Pi_v^-\}$ for all v where τ_v is square integrable. Here, $\Pi_v^\times$ is the nontempered quotient $L(\nu_v^{1/2} \mu_v \tau_v, \mu_v \nu_v^{-1/2})$ of $\nu_v^{1/2} \mu_v \tau_v \rtimes \mu_v \nu_v^{-1/2}$ and Π_v^- is the cuspidal representation denoted by $\delta^-(\nu_v^{1/2} \mu_v \tau_v, \mu_v \nu_v^{-1/2})$ in [F4, V. 8]. At each place v where τ_v is the induced representation $I(\eta_v, \eta_v^{-1})$ for some character η_v of $F_v^\times$, $\{\Pi_v\}$ is the singleton consisting of the irreducible induced representation $\Pi_v^\times := \eta_v \mu_v 1_{2,v} \rtimes \eta_v^{-1}$.

The multiplicity formula for $\Pi' \in \{\Pi\}$ is

$$m(\Pi') = \frac{1}{2} \left(1 + \varepsilon(\mu\tau, 1/2) (-1)^{n(\Pi')} \right),$$

where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$, and $\varepsilon(\mu\tau, \frac{1}{2})$ is the value at $1/2$ of the epsilon factor in the functional equation of the L -function of $\mu\tau$.

REMARK. In [A3], $[\tau, \mu 1_2]$ is denoted by $(\mu \boxtimes \nu(2)) \boxplus (\tau \boxtimes 1)$ and is called a *Saito, Kurokawa type* packet.

3. Let μ_1, μ_2 be distinct unitary characters of C_F such that $\mu_1^2 = \mu_2^2$. Suppose $\mu_{i,v}$ ($i = 1, 2$) is unramified for all v lying outside of some finite set of places S . The quasi-packet $[\mu_1 1_2, \mu_2 1_2]$ is parametrized by the global datum

$$\begin{aligned} \mathcal{C}([\mu_1 1_2, \mu_2 1_2], S, \hat{G}) &\ni \left[t(\mu_{1,v} 1_{2,v}), t(\mu_{2,v} 1_{2,v}) \right]_{v \notin S} \\ &= \left[\text{diag} \left(\mu_{1,v} q_v^{1/2}, \mu_{2,v} q_v^{1/2}, \mu_{2,v} q_v^{-1/2}, \mu_{1,v} q_v^{-1/2} \right) \right]_{v \notin S}. \end{aligned}$$

In particular, the quasi-packet $[\mu_1 1_2, \mu_2 1_2]$ lifts to the representation $I_{(2,2)}(\mu_1 1_2, \mu_2 1_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

If $\mu_1^2 = \mu_2^2 = 1$, let ζ be the nontrivial quadratic character $\frac{\mu_1}{\mu_2}$ of C_F . In this case, we know from [F4, V. 10] that $[\mu_1 1_2, \mu_2 1_2]$ is the quasi-packet $\{\Pi\} = \otimes_v \{\Pi_v^\times, \Pi_v^-\}$. Here, $\Pi_v^\times$ is the nontempered quotient

$$L(\nu_v \zeta_v, \zeta_v \rtimes \nu_v^{-1/2} \mu_{2,v})$$

of the induced representation $I_v = \nu_v \zeta_v \times \zeta_v \rtimes \nu_v^{-1/2} \mu_{2,v}$ of $\mathrm{GSp}(2, F_v)$, and

$$\Pi_v^- = \begin{cases} \delta^-(\zeta_v \nu_v^{1/2} \mathrm{St}_{2,v}, \zeta_v \mu_{2,v} \nu_v^{-1/2}) & \text{if } \zeta_v \neq 1, \\ L(\zeta_v \nu_v^{1/2} \mathrm{St}_{2,v}, \zeta_v \mu_{2,v} \nu_v^{-1/2}) & \text{if } \zeta_v = 1, \end{cases}$$

where $L(\zeta_v \nu_v^{1/2} \text{St}_{2,v}, \zeta_v \mu_{2,v} \nu_v^{-1/2})$ is the unique nontempered subquotient of the induced representation $\zeta_v \nu_v^{1/2} \text{St}_{2,v} \rtimes \zeta_v \mu_{2,v} \nu_v^{-1/2}$, and $\delta^-(\zeta_v \nu_v^{1/2} \text{St}_{2,v}, \zeta_v \mu_{2,v} \nu_v^{-1/2})$ is the cuspidal member of the local packet which contains the unique square integrable subrepresentation $\delta(\zeta_v \nu_v^{1/2} \text{St}_{2,v}, \zeta_v \mu_{2,v} \nu_v^{-1/2})$ of I_v . At almost every place v , Π_v^+ is unramified.

The multiplicity formula for $\Pi' \in \{\Pi\}$ is $\frac{1}{2}(1 + (-1)^{n(\Pi')})$, where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$.

REMARK. In [A3], $[\mu_1 1_2, \mu_2 1_2]$ is denoted by $(\mu_1 \boxtimes \nu(2)) \boxplus (\mu_2 \boxtimes \nu(2))$ and is called a *Howe, Piatetskii-Shapiro type packet*.

4.3.1.2. *A Stable Packet.* Also contained in [A3] is a classification of the stable (quasi-)packets of $\text{GSp}(2, \mathbb{A}_F)$. We now describe one which is of interest to us. Let ζ be a nontrivial quadratic character of C_F . Let τ be a ζ -invariant, cuspidal automorphic representation of $\text{GL}(2, \mathbb{A}_F)$. Suppose ζ_v, τ_v are unramified for all v lying outside of some finite set of places S . The stable quasi-packet $\{L(\nu\zeta, \nu^{-1/2}\tau)\}$ is parametrized by the global datum

$$\mathcal{C} \left(\{L(\nu\zeta, \nu^{-1/2}\tau)\}, S, \hat{G} \right) \ni \left[\text{blockdiag} \left(q_v^{1/2} \zeta_v t(\tau_v), q_v^{-1/2} t(\tau_v) \right) \right]_{v \notin S}.$$

In particular, $\{L(\nu\zeta, \nu^{-1/2}\tau)\}$ lifts to the Langlands quotient $J(\nu^{1/2}\tau, \nu^{-1/2}\tau)$ of the induced representation $I_{(2,2)}(\nu^{1/2}\tau, \nu^{-1/2}\tau)$ of $\text{GL}(4, \mathbb{A}_F)$.

If $\omega_\tau = \zeta$, then $\{L(\nu\zeta, \nu^{-1/2}\tau)\}$ is the quasi-packet defined in [F4, V. 10]. It is the restricted tensor product $\otimes_v \{L_v\}$, where $\{L_v\}$ is the local quasi-packet defined as follows:

- If $\zeta_v \neq 1$ and τ_v is cuspidal, $\{L_v\}$ is a singleton consisting of the unique nontempered quotient $L(\nu_v \zeta_v, \nu_v^{-1/2} \tau_v)$ of $\nu_v \zeta_v \rtimes \nu_v^{-1/2} \tau_v$.
- Suppose $\zeta \neq 1$ and τ_v is the induced representation $I(\mu_v, \zeta_v \mu_v)$ for some (possibly trivial) quadratic character μ_v of $F_v^\times$. Then,

$$\{L_v\} = \{L(\nu_v \zeta_v, \zeta_v \rtimes \nu_v^{-1/2} \tau_v), \delta^-(\zeta_v \nu_v^{1/2} \text{St}_{2,v}, \zeta_v \mu_v \nu_v^{-1/2})\}.$$

- Suppose $\zeta_v = 1$, then by the theory of monomial representations τ_v must be induced. Suppose $\tau_v = I(\mu_v, \mu_v^{-1})$, where μ_v is a unitary character of $F_v^\times$. If $\mu_v^2 \neq 1$, $\{L_v\}$ is the singleton consisting of the irreducible representation $\mu_v^{-2} \rtimes \mu_v 1_{2,v}$. If $\mu_v^2 = 1$, $\{L_v\}$ consists of the nontempered subquotients $L(\nu_v, 1_{2,v} \rtimes \nu_v^{-1/2} \mu_v), L(\nu_v^{1/2} \text{St}_{2,v}, \nu_v^{-1/2} \mu_v)$ of $\nu_v \times 1 \rtimes \nu_v^{-1/2} \mu_v$ (see [ST]).

Every member of $\{L(\nu\zeta, \nu^{-1/2}\tau)\}$ occurs with multiplicity one in the discrete spectrum of $\text{GSp}(2)$.

REMARK. In [A3], $\{L(\nu\zeta, \nu^{-1/2}\tau)\}$ is denoted by $\tau \boxtimes \nu(2)$ and is called a *Soudry type packet*.

4.3.1.3. *ε -Invariant Unstable Packets.* For any character χ of C_E , put ${}^\sigma \chi(x) := \chi(\sigma x)$ for all $x \in C_E$.

4.9. LEMMA. Suppose $\{\Pi\}$ is an ε -invariant, unstable, discrete spectrum global (quasi-)packet of $\text{GSp}(2, \mathbb{A}_F)$. Then,

- (1) $\{\Pi\} \neq [\tau, \mu 1_2]$ for any cuspidal automorphic representation τ of the group $\text{GL}(2, \mathbb{A}_F)$, unitary character μ of C_F , such that $\omega_\tau = \mu^2$.
- (2) If $\{\Pi\} = [\tau_1, \tau_2]$ for two distinct cuspidal automorphic representations τ_1, τ_2 of $\text{GL}(2, \mathbb{A}_F)$, then one of the following conditions is satisfied:
 - The representation τ_1 is not E -monomial, and $\tau_2 = \varepsilon \tau_1$.
 - There exist unitary characters ζ, χ of C_E such that τ_1, τ_2 are equal to the E -monomial representations $\pi(\zeta \chi), \pi(\zeta {}^\sigma \chi)$, respectively.

- (3) If $\{\Pi\} = [\mu_1 1_2, \mu_2 1_2]$, where μ_1, μ_2 are distinct unitary characters of C_F , then $\mu_2 = \varepsilon \mu_1$.

PROOF. Suppose $\{\Pi\} = [\tau, \mu 1_2]$ for some cuspidal automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_F)$ and a unitary character μ of C_F such that $\mu^2 = \omega_\tau$. Let Π be an ε -invariant discrete spectrum representation in $\{\Pi\}$. For a sufficiently large finite set of places S such that Π_v is unramified for all $v \notin S$, we have:

$$\mathcal{C}(\Pi, S, \hat{G}) \ni \left[\text{blockdiag}(t(\tau_v), t(\mu_v 1_{2,v})) \right]_{v \notin S}.$$

Since Π is ε -invariant, $\text{blockdiag}(t(\tau_v), t(\mu_v 1_{2,v}))$ must be conjugate in $\hat{G} = \mathrm{GSp}(2, \mathbb{C})$ to $\text{blockdiag}(\varepsilon_v t(\tau_v), \varepsilon_v t(\mu_v 1_{2,v}))$ for each $v \notin S$. In particular, at each place $v \notin S$ such that $\varepsilon_v \neq 1$, τ_v is equivalent to either $\varepsilon_v \tau_v$ or $\varepsilon_v \mu_v 1_{2,v}$. If $\tau_v \cong \varepsilon_v \tau_v$, then μ_v must be equal to $\varepsilon_v \mu_v$, which cannot hold because $\varepsilon_v \neq 1$. Hence, τ_v is equivalent to $\varepsilon_v \mu_v 1_{2,v}$ at infinitely many places v . In particular, τ should be one dimensional, but by assumption τ is cuspidal. The first statement of the lemma follows.

Suppose $\{\Pi\} = [\tau_1, \tau_2]$, where τ_1, τ_2 are two distinct cuspidal automorphic representations of $\mathrm{GL}(2, \mathbb{A}_F)$ whose central characters satisfy $\omega_{\tau_1} = \omega_{\tau_2}$. Let Π be an ε -invariant discrete spectrum representation in $\{\Pi\}$. For a finite set of places S such that Π_v is unramified for all $v \notin S$, we have:

$$\mathcal{C}(\Pi, S, \hat{G}) \ni \left[[t(\tau_{1,v}), t(\tau_{2,v})] \right]_{v \notin S}.$$

View $\mathrm{GSp}(2)$ as a twisted endoscopic group with respect to the triple $(\mathrm{GL}(4), \theta, 1)$ (see [KS, Chap. 2]), where θ is the automorphism $g \mapsto {}^t g^{-1}$ of $\mathrm{GL}(4)$. The F-H classes in $\mathrm{GSp}(2, \mathbb{C})$ parametrizing Π , viewed as classes in $\mathrm{GL}(4, \mathbb{C})$, parametrize the induced representation $I_{(2,2)}(\tau_1, \tau_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

Since Π is an ε -invariant representation, the element $[t(\tau_{1,v}), t(\tau_{2,v})]$ is conjugate to $[\varepsilon_v t(\tau_{1,v}), \varepsilon_v t(\tau_{2,v})]$ in $\mathrm{GSp}(2, \mathbb{C})$, and hence in $\mathrm{GL}(4, \mathbb{C})$, for each $v \notin S$. Consequently, $I_{(2,2)}(\tau_1, \tau_2)$ is equivalent to $I_{(2,2)}(\varepsilon \tau_1, \varepsilon \tau_2)$, which implies that the set $\{\tau_1, \tau_2\}$ is equal to $\{\varepsilon \tau_1, \varepsilon \tau_2\}$ (see [JS1, Thm. 4.4]).

If $\tau_2 \neq \varepsilon \tau_1$, then τ_1 and τ_2 must both be E -monomial. Consequently, there are characters μ_1, μ_2 of C_E such that $\tau_1 = \pi(\mu_1), \tau_2 = \pi(\mu_2)$. The central character of the monomial representation $\pi(\mu_i)$ ($i = 1, 2$) is $\mu_i|_{\mathbb{A}_F^\times} \cdot \varepsilon$; hence, the condition $\omega_{\tau_1} = \omega_{\tau_2}$ implies that $\mu_1|_{\mathbb{A}_F^\times} = \mu_2|_{\mathbb{A}_F^\times}$. Since $\frac{\mu_2}{\mu_1}|_{\mathbb{A}_F^\times}$ is trivial, there exists a character χ of C_E such that $\frac{\mu_2}{\mu_1} = \frac{\sigma \chi}{\chi}$. Let $\zeta = \frac{\mu_1}{\chi}$, then $\tau_1 = \pi(\zeta \chi)$ and $\tau_2 = \pi(\zeta \sigma \chi)$. The second statement of the lemma follows.

Suppose $\{\Pi\} = [\mu_1 1_2, \mu_2 1_2]$, where μ_1, μ_2 are two distinct unitary characters of C_F such that $\mu_1^2 = \mu_2^2$. Let Π be an ε -invariant discrete spectrum representation in $\{\Pi\}$. It lifts to the induced representation $I_{(2,2)}(\mu_1 1_2, \mu_2 1_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$. By the same reasoning used in the second case, $\Pi \cong \varepsilon \Pi$ implies that the induced representations $I_{(2,2)}(\mu_1 1_2, \mu_2 1_2)$ and $I_{(2,2)}(\varepsilon \mu_1 1_2, \varepsilon \mu_2 1_2)$ are equivalent. Hence, $\mu_2 = \varepsilon \mu_1$, and the final statement of the lemma follows. $\square$

4.3.2. Multi-Packets of $\mathbf{H}_1(\mathbb{A}_F)$. Suppose τ is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, and χ is a character of C_E . Recall from Section 3.1.3.3 that we let $\tau \otimes_1 \chi$ denote the global packet of $\mathbf{H}_1(\mathbb{A}_F)$ consisting of those irreducible constituents of $\tau|_{\mathrm{GL}(2, \mathbb{A}_F)^E} \otimes_1 \chi$ which are unramified at almost all places.

The following theorem rests on the assumption that a result of Flicker's ([F4]) extends to an analogous result in an announced but unpublished work of Arthur's ([A]).

4.10. THEOREM. *Let τ be an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. Let χ be a character of C_E . Suppose the packet $\tau \otimes_1 \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to an ε -trace identity. The following holds:*

- *If $\chi \neq {}^\sigma\chi$, and τ is not E -monomial, then the only other contribution to the ε -trace identity from $\mathbf{H}_1$ is $\tau \otimes_1 {}^\sigma\chi$.*
- *If $\chi \neq {}^\sigma\chi$, and τ is the cuspidal monomial representation $\pi(\theta)$ for some character θ of C_E , then the following packets of $\mathbf{H}_1(\mathbb{A}_F)$ contribute to the ε -trace identity provided that they are distinct from $\pi(\theta) \otimes_1 \chi$:*
 - $\pi(\theta) \otimes_1 {}^\sigma\chi$,
 - $\pi(\chi) \otimes_1 \theta$,
 - $\pi(\chi) \otimes_1 {}^\sigma\theta$.
- *If $\chi = {}^\sigma\chi$, then $\tau \otimes_1 \chi$ is the only packet of $\mathbf{H}_1(\mathbb{A}_F)$ which contributes to the ε -trace identity.*

REMARK. The theorem only comments on contribution from $\mathbf{H}_1$, it says nothing regarding contribution from $\mathbf{H}_2$.

SKETCH OF PROOF. The proof requires twisted endoscopic lifting results for the triple $(\mathrm{GL}(4), \theta, 1)$, where θ is the automorphism $g \mapsto {}^t g^{-1}$ of $\mathrm{GL}(4)$ (see [KS], [A3], [F3]). Consider the F -split θ -twisted endoscopic group $\mathbf{C}$ of $\mathrm{GL}(4)$ (see [F3, p. 22–23]). It is equal to $(\mathrm{GL}(2) \times \mathrm{GL}(2))'$, where the prime indicates that the two $\mathrm{GL}(2)$ components have equal determinants. Any automorphic representation of $\mathbf{C}(\mathbb{A}_F)$ has the form

$$\tau_1 \times \tau_2 : (g, h) \mapsto \tau_1(g) \otimes \tau_2(h), \quad \forall (g, h) \in \mathbf{C}(\mathbb{A}_F),$$

where τ_1, τ_2 are automorphic representations of $\mathrm{GL}(2, \mathbb{A}_F)$. Because of the condition on the determinants of the two $\mathrm{GL}(2)$ components, $\tau_1 \times \tau_2$ is the same as $\mu\tau_1 \times \mu^{-1}\tau_2$ for any quasi-character μ of C_F . Suppose $\tau_1 \times \tau_2$ contributes to a θ -twisted trace identity derived from Kottwitz-Shelstad's formula for $(\mathrm{GL}(4), \theta, 1)$. Expecting the results of lifting from $\mathrm{SO}(4)$ to $\mathrm{PGL}(4)$ to be a guide (see proof of [F4, Prop. 4.1]), the only other contribution from $\mathbf{C}$ to the trace identity should be $\tau_2 \times \tau_1$, provided $\tau_1 \neq \tau_2$. Although the work of [F4] has not yet been extended to the case of nontrivial central characters, Arthur has announced in [A3] the lifting results for the triple $(\mathrm{GL}(4), \theta, 1)$. It is likely that a proof of the statement which we assume here will be contained in a complete treatise on the problem. For now, we shall take for granted the above hypothesis regarding the fibers of the lifting from $\mathbf{C}$ to $\mathrm{GL}(4)$.

Recall the hypothesis of our theorem: A global packet $\tau \otimes_1 \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to an ε -trace identity. If $\pi'_1 = \tau' \otimes_1 \chi'$ is another global packet of $\mathbf{H}_1(\mathbb{A}_F)$ which contributes to the ε -trace identity, then the global data parametrizing the representations $\tau' \times \pi(\chi')$ and $\tau \times \pi(\chi)$ of $\mathbf{C}(\mathbb{A}_F)$ lift to the same datum in $\mathrm{GL}(4, \mathbb{C})$. Hence, by our hypothesis $\tau' \times \pi(\chi')$ is equivalent to $\tau \times \pi(\chi)$ and/or $\pi(\chi) \times \tau$.

If two representations $\tau_1 \times \tau_2, \tau'_1 \times \tau'_2$ of $\mathbf{C}(\mathbb{A}_F)$ are equivalent to each other, then there exists a quasi-character μ of C_F such that $\tau'_1 = \mu^{-1}\tau_1$ and $\tau'_2 = \mu\tau_2$. Therefore, if $\tau' \times \pi(\chi')$ is equivalent to $\pi(\chi) \times \tau$, then $\tau' = \mu^{-1}\pi(\chi)$ and $\pi(\chi') = \mu\tau$ for some quasi-character μ of C_F . Since twisting by a character does not change whether a representation of $\mathrm{GL}(2, \mathbb{A}_F)$ is E -monomial, $\tau' \times \pi(\chi')$ cannot be equivalent to $\pi(\chi) \times \tau$ if τ is not E -monomial.

Suppose $\tau' \times \pi(\chi') = \tau \times \pi(\chi)$. If $\pi(\chi') = \mu\pi(\chi)$, then χ' is equal to either $(\mu \circ N_{E/F}) \cdot \chi$ or $(\mu \circ N_{E/F}) \cdot {}^\sigma\chi$ ([L, Chap. 7]). Consequently, if τ is not E -monomial and $\chi \neq {}^\sigma\chi$, we must have $\tau' = \mu^{-1}\tau$, and $\chi' = (\mu \circ N_{E/F}) {}^\sigma\chi$. For any quasi-character μ of C_F , $\mu^{-1}\tau \otimes_1 (\mu \circ N_{E/F}) \chi$ defines the same representation. We conclude that if τ

is non- E -monomial and $\chi \neq {}^\sigma \chi$, then $\tau \otimes_1 {}^\sigma \chi$ is the only other packet of $\mathbf{H}_1(\mathbb{A}_F)$ which contributes to the ε -trace identity.

If τ is non- E -monomial and $\chi = {}^\sigma \chi$, then there is no other contribution from $\mathbf{H}_1(\mathbb{A}_F)$.

Suppose τ is the cuspidal monomial representation $\pi(\theta)$ associated with some character $\theta \neq \chi$ of C_E such that $\theta \neq {}^\sigma \theta$. Then, the representation $\tau' \times \pi(\chi')$ is equivalent to either: (i) $\pi(\theta) \times \pi(\chi)$, or (ii) $\pi(\chi) \times \pi(\theta)$. In the first case, we have $\tau' \otimes_1 \chi' = \pi(\theta) \otimes_1 {}^\sigma \chi$ if $\chi \neq {}^\sigma \chi$. In the second case, $\tau' \otimes_1 \chi'$ must be equal to either $\pi(\chi) \otimes_1 \theta$ or $\pi(\chi) \otimes_1 {}^\sigma \theta$ provided that they are distinct from $\pi(\theta) \otimes_1 \chi$. If $\chi = {}^\sigma \chi = \mu \circ N_{E/F}$ for some character μ of C_F , then $\pi(\chi) = I(\mu, \mu\varepsilon)$. From the discussion in Section 3.1.3.3, neither $I(\mu, \mu\varepsilon) \otimes_1 \theta$ nor $I(\mu, \mu\varepsilon) \otimes_1 {}^\sigma \theta$ contributes to the stable spectrum of $\mathbf{H}_1$.

Contingent upon published results on the θ -twisted endoscopic lifting for $\mathrm{GL}(4)$, the theorem follows. $\square$

Suppose a global packet of $\mathbf{H}_1(\mathbb{A}_F)$ has the form $\tau \otimes_1 \chi$, where τ is a representation of $\mathrm{GL}(2, \mathbb{A}_F)$ and χ is a character of C_E which is not equal to ${}^\sigma \chi$. If $\mathrm{tr}(\tau \otimes_1 \chi)(f_1)$ appears in an ε -trace identity, then so does $\mathrm{tr}(\tau \otimes_1 {}^\sigma \chi)(f_1)$. If an element $(h, x) \in \mathbf{H}_1(\mathbb{A}_F)$ is a norm of some element $g \in \mathrm{GSp}(2, \mathbb{A}_F)$, then $(h, \sigma x)$ is also a norm of g . Therefore, we expect that if $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$ and $f_1 \in C(\mathbf{H}_1(\mathbb{A}_F), \omega)$ are matching functions, then $f_1(h, x) = f_1(h, \sigma x)$ for all $(h, x) \in \mathbf{H}_1(\mathbb{A}_F)$. Consequently, the terms $\mathrm{tr}(\tau \otimes_1 \chi)(f_1)$ and $\mathrm{tr}(\tau \otimes_1 {}^\sigma \chi)(f_1)$ in the trace identity combine into a single term $2 \cdot \mathrm{tr}(\tau \otimes_1 \chi)(f_1)$.

4.3.3. Multi-Packets of $\mathbf{H}_2(\mathbb{A}_F)$. For any representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$, let ${}^\sigma \tau$ denote the $\mathrm{GL}(2, \mathbb{A}_E)$ -module which sends $(g_{ij}) \in \mathrm{GL}(2, \mathbb{A}_E)$ to $\tau((\sigma g_{ij}))$. For any representation $\pi_2 = \tau \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$, where $\omega_\tau = \mu \circ N_{E/F}$, put ${}^\sigma \pi_2 := {}^\sigma \tau \otimes_2 \mu$.

4.11. LEMMA. *Suppose a cuspidal automorphic representation $\pi_2 = \tau \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to an ε -trace identity.*

- *If $\pi_2 \neq {}^\sigma \pi_2$, then the only other cuspidal automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ which contributes to the ε -trace identity is ${}^\sigma \pi_2$.*
- *If $\pi_2 = {}^\sigma \pi_2$, then no other cuspidal automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the ε -trace identity.*

REMARK. The condition “cuspidal” is crucial. The lemma says nothing regarding non-cuspidal contributions from $\mathbf{H}_2(\mathbb{A}_F)$.

Suppose π'_2 is another cuspidal automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ which contributes to the ε -trace identity. Let v be any finite place for which the representations $\pi_{2,v}, \pi'_{2,v}$ are unramified. From Section 4.3, it follows that the F-H classes parametrizing $\pi_{2,v}, \pi'_{2,v}$ lift via $\xi_2 : {}^L H_2 \rightarrow \hat{G}$ to the same F-H class in $\hat{G}$.

To examine what the above statement entails, we need to consider separately the case where the place v is prime in E and the case where v splits in E .

4.12. CLAIM. *Let v be a finite place of F which is unramified, prime in E . If $\pi_{2,v}$ and $\pi'_{2,v}$ are unramified and their Frobenius-Hecke classes lift to the same conjugacy class in $\hat{G}$, then $\pi_{2,v}$ and $\pi'_{2,v}$ are equivalent.*

PROOF. Since π_2 is cuspidal, the unramified representation $\pi_{2,v}$ is an irreducible induced representation $I(\chi_1, \chi_2) \otimes_2 \mu_v$, where μ_v is an unramified character of $F_v^\times$, and χ_1, χ_2 are unramified characters of $E_v^\times$. Since $\mu_v \circ N_{E/F} = \chi_1 \chi_2$, $\pi_{2,v}$ has the form $I\left(\chi_1, \frac{\mu_v \circ N_{E/F}}{\chi_1}\right) \otimes_2 \mu_v$. Since v is prime in E , and χ_1 is unramified, we have $\chi_1 = \chi'_1 \circ N_{E/F}$ for some unramified character χ'_1 of $F_v^\times$. Likewise, $\pi'_{2,v} = I\left(\theta_1, \frac{\mu'_v \circ N_{E/F}}{\theta_1}\right) \otimes_2 \mu'_v$,

where $\theta_1 = \theta'_1 \circ N_{E/F}$ for some unramified character θ'_1 of $F_v^\times$, and μ'_v is an unramified character of $F_v^\times$.

From Section 4.2, the lift of the F-H class parametrizing $\pi_{2,v}$ is represented by

$$\hat{g} = \text{diag} \left(\chi'_1, -\chi'_1, \frac{-\mu_v}{\chi'_1}, \frac{\mu_v}{\chi'_1} \right) \in \text{GSp}(2, \mathbb{C}),$$

while the lift of the class parametrizing $\pi'_{2,v}$ is represented by

$$\hat{g}' = \text{diag} \left(\theta'_1, -\theta'_1, \frac{-\mu'_v}{\theta'_1}, \frac{\mu'_v}{\theta'_1} \right) \in \text{GSp}(2, \mathbb{C}).$$

If $\hat{g}$ and $\hat{g}'$ belong to the same conjugacy class in $\hat{G}$, then there must exist an element w in the Weyl group W of $\hat{G}$ such that $w\hat{g} = \hat{g}'$. Observe that $\mu_v(\varpi_v)$ and $\mu'_v(\varpi_v)$ are the respective similitude factors of $\hat{g}$ and $\hat{g}'$. Since the similitude factor is invariant under conjugation, we conclude that $\mu_v = \mu'_v$. Since E_v/F_v is unramified, $\varepsilon_v(\varpi_v)$ is equal to -1 . By permuting the entries of $\hat{g}$ under W , we conclude that θ' is equal to one of the following characters:

- (1) $\frac{\mu_v}{\chi'_1}$,
- (2) $\varepsilon_v \chi'_1$,
- (3) $\frac{\varepsilon_v \mu_v}{\chi'_1}$,
- (4) χ'_1 .

Since $\varepsilon_v \circ N_{E/F} = 1$, in cases 2 and 4 we have $\theta_1 = \theta'_1 \circ N_{E/F} = \chi'_1 \circ N_{E/F} = \chi_1$. Hence, $\pi'_{2,v} = \pi_{2,v}$. In cases 1 and 3, we have $\pi'_{2,v} = I \left(\frac{\mu_v \circ N_{E/F}}{\chi_1}, \chi_1 \right) \otimes_2 \mu_v$, which is equivalent to $\pi_{2,v}$. Hence, if the F-H classes parametrizing the unramified representations $\pi_{2,v}$ and $\pi'_{2,v}$ lift to the same conjugacy class in $\hat{G}$, then $\pi_{2,v}$ and $\pi'_{2,v}$ are equivalent. $\square$

4.13. CLAIM. *Let v be a finite place of F which splits in E . If $\pi_{2,v}$ and $\pi'_{2,v}$ are unramified, and their Frobenius-Hecke classes lift to the same conjugacy class in $\hat{G}$, then $\pi'_{2,v}$ is equivalent to either $\pi_{2,v}$ or ${}^\sigma \pi_{2,v}$.*

PROOF. If v is a finite place of F which splits in E , then $H_{2,v} = \mathbf{H}_2(F_v)$ has the form

$$[\text{GL}(2, F_v) \times \text{GL}(2, F_v) \times F_v^\times] / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in F_v^\times\}.$$

Suppose v_1, v_2 are the two places of E which lie above v . We have $\pi_{2,v} = \tau_1 \times \tau_2 \times \mu_v$, where $\tau_i = \tau_{v_i}$ ($i = 1, 2$), and ${}^\sigma \pi_{2,v} = \tau_2 \times \tau_1 \times \mu_v$. Since

$$\omega_{\tau_1}(a)\omega_{\tau_2}(b)\mu_v(a)^{-1}\mu_v(b)^{-1} = 1, \quad \forall a, b \in F_v^\times,$$

we must have $\omega_{\tau_1} = \omega_{\tau_2} = \mu_v$. In particular, μ_v is uniquely determined by either τ_1 or τ_2 .

Observe that $H_{2,v}$ is isomorphic via $(g, h, c) \mapsto (g, ch)$ to

$$[\text{GL}(2, F_v) \times \text{GL}(2, F_v)] / \{(zI_2, z^{-1}I_2) : z \in F_v^\times\}.$$

We identify representations of $H_{2,v}$ with representations of the above group, and write $\pi_{2,v} = \tau_1 \times \tau_2$, ${}^\sigma \pi_{2,v} = \tau_2 \times \tau_1$.

If $\pi_{2,v} = \tau_1 \times \tau_2$ is unramified, then $\tau_i = I(\alpha_i, \beta_i)$ ($i = 1, 2$), where α_i, β_i are unramified characters of $F_v^\times$. Likewise, $\pi'_{2,v} = I(\alpha'_1, \beta'_1) \times I(\alpha'_2, \beta'_2)$ for some unramified characters α'_i, β'_i of $F_v^\times$.

The F-H class parametrizing $\pi_{2,v}$ lifts to the conjugacy class in $\hat{G}$ represented by

$$\hat{g} = [\text{diag}(\alpha_1, \beta_1), \text{diag}(\alpha_2, \beta_2)],$$

while the F-H class parametrizing $\pi'_{2,v}$ lifts to the conjugacy class in $\hat{G}$ represented by

$$\hat{g}' = [\text{diag}(\alpha'_1, \beta'_1), \text{diag}(\alpha'_2, \beta'_2)].$$

By assumption, there exists an element $w \in W$ such that $w\hat{g} = \hat{g}'$. If we label the entries of the maximal diagonal torus of $\hat{G}$ by 1, 2, 3, 4, then $W = D_4$ is generated by the permutations (3421) and (23).

Suppose (23) $\hat{g} = \hat{g}'$. The action of (23) sends $\hat{g} = [\text{diag}(\alpha_1, \beta_1), \text{diag}(\alpha_2, \beta_2)]$ to $[\text{diag}(\alpha_1, \beta_1), \text{diag}(\beta_2, \alpha_2)]$. Since $I(\beta_2, \alpha_2)$ is equivalent to $I(\alpha_2, \beta_2)$, $\pi'_{2,v}$ is equivalent to $\pi_{2,v}$.

The action of (3421) takes $\hat{g}$ to $[\text{diag}(\alpha_2, \beta_2), \text{diag}(\beta_1, \alpha_1)]$. Hence, if $\hat{g}' = (3421)\hat{g}$, then $\pi'_{2,v} = \tau_2 \times \tau_1 = {}^\sigma \pi_{2,v}$.

We conclude that if the F-H classes parametrizing the unramified representations $\pi_{2,v}$ and $\pi'_{2,v}$ lift to the same conjugacy class in $\hat{G}$, then $\pi'_{2,v}$ is equivalent to either $\pi_{2,v}$ or ${}^\sigma \pi_{2,v}$. In particular, μ'_v is equal to μ_v . $\square$

From the previous two claims, we see that $\mu_v = \mu'_v$ for almost all v , which implies that $\mu = \mu'$ by the weak approximation theorem for $\mathbb{A}_F$ (The weak approximation theorem says that F is dense in $\prod_{v \in S} F_v$ for any finite set of places S . It follows that $F^\times$ is dense in $\prod_{v \in S} F_v^\times$. Using that μ and μ' are continuous characters of $\mathbb{A}_F^\times$ whose restrictions to $F^\times$ are trivial, our assertion follows).

Suppose $\pi'_2 = \tau' \otimes \mu$, where τ' is a cuspidal automorphic representation of $\text{GL}(2, \mathbb{A}_E)$. To prove Lemma 4.11, it remains to show that:

4.14. CLAIM. τ' is equivalent to either τ or ${}^\sigma \tau$.

Before we prove the claim, we need a digression: Let k be a number field. Let V_k be the set of places of k . Let n, m be positive integers. Let π be an automorphic representation of $\text{GL}(n, \mathbb{A}_k)$ which is unramified at places outside of a finite set $S \subset V_k$. For each $v \notin S$, let $t(\pi_v)$ be a matrix in $\text{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes π_v . Let π' be an automorphic representation of $\text{GL}(m, \mathbb{A}_k)$ which is unramified at places outside of S . Define $t(\pi'_v) \in \text{GL}(m, \mathbb{C})$ likewise for π' .

Consider $t(\pi_v), t(\pi'_v)$ as endomorphisms of $\mathbb{C}^n, \mathbb{C}^m$, respectively. Their tensor product $t(\pi_v) \otimes t(\pi'_v)$, being an endomorphism of $\mathbb{C}^{mn}$, is represented by an $mn \times mn$ complex matrix. Define a partial L -function in $s \in \mathbb{C}$ as follows:

$$L^S(s, \pi \otimes \pi') := \prod_{v \notin S} \det(1 - t_v(\pi) \otimes t_v(\pi') q_v^{-s})^{-1},$$

where q_v is the cardinality of the residue field of k_v . If π and π' are cuspidal, it is known ([JS]) that: (i) $L^S(s, \pi \otimes \pi')$ does not vanish on the line $\text{Re } s = 1$; (ii) $L^S(s, \pi \otimes \pi')$ has a pole of order 1 at $s = 1$ if and only if $n = m$ and π' is equivalent to $\tilde{\pi}$, the contragradient of π .

PROOF OF CLAIM 4.14. Let S be a finite set of places of F such that $\pi_{2,v}, \pi'_{2,v}$ are unramified for all $v \notin S$. We know, for $v \notin S$, that:

$$\pi'_{2,v} = \begin{cases} \pi_{2,v} & \text{if } v \text{ is unramified, prime in } E, \\ \pi_{2,v} \text{ or } {}^\sigma \pi_{2,v} & \text{if } v \text{ splits in } E. \end{cases}$$

Consider τ, τ' as automorphic representations of $\text{GL}(2, \mathbb{A}_E)$, with E as the base field. Let S_E be the set of places of E which lie above the places in S . In particular, S_E is finite. For

any place w of E outside of S_E , τ_w and τ'_w are unramified. We have:

$$\tau'_w = \tau_w \text{ or } ({}^\sigma \tau)_w.$$

Notice the order of operation: We first apply σ to the global τ , and then we take the component at the place w . We have the following equality of products of partial L -functions:

$$L^{S_E}(s, \tau' \otimes \tilde{\tau}) L^{S_E}(s, {}^\sigma \tau' \otimes \tilde{\tau}) = L^{S_E}(s, \tau \otimes \tilde{\tau}) L^{S_E}(s, {}^\sigma \tau \otimes \tilde{\tau}).$$

Since π_2 and π'_2 are cuspidal automorphic representations of $\mathbf{H}_2(\mathbb{A}_F)$, τ_w and τ'_w are cuspidal automorphic representations of $\mathrm{GL}(2, \mathbb{A}_E)$. By [JS] and [Sh], the product on the right has a single pole at $s = 1$; therefore, the product on the left must have a pole, which implies that τ' is equivalent to either τ or ${}^\sigma \tau$. $\square$

Let $1_{\mathrm{GL}(2, \mathbb{A}_E)}$ denote the trivial representation of $\mathrm{GL}(2, \mathbb{A}_E)$.

4.15. LEMMA. *Let χ be a character of C_E , μ be a character of C_F , such that $\chi^2 = \mu \circ N_{E/F}$. Suppose the automorphic representation*

$$\pi_2 = (\chi \circ \det) \otimes_2 \mu = \chi 1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes_2 \mu$$

of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to an ε -trace identity.

- *If $\chi \neq {}^\sigma \chi$, then the only other contribution from $\mathbf{H}_2$ to the ε -trace identity is ${}^\sigma \pi_2 = {}^\sigma \chi 1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes_2 \mu$.*
- *If $\chi = {}^\sigma \chi$, then no other automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the ε -trace identity.*

PROOF. Suppose an automorphic representation $\pi'_2 = \tau \otimes_2 \mu' \neq \pi_2$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity. As in the case where π_2 is cuspidal, we consider the places v , unramified in E , where both $\pi_{2,v}$ and $\pi'_{2,v}$ are unramified. Of these places, we examine separately those which split in E and those which do not.

If v does not split in E , then by the same argument used in the proof of Lemma 4.11, we conclude that $\pi_{2,v} = \pi'_{2,v}$. Suppose v splits into two places v_1, v_2 of E . For $i = 1, 2$, let $\chi_i = \chi_{v_i}$. Since $\chi^2 = \mu \circ N_{E/F}$, we have $\chi_1^2 = \chi_2^2 = \mu_v$. Let ν_v be the normalized absolute value function on F_v . Then, $\pi'_{2,v} = \tau'_{v_1} \times \tau'_{v_2}$ is parametrized by the conjugacy class of one of the following two elements in ${}^L H_2$:

- (1) $\left(\begin{pmatrix} \chi_1 \nu_v^{1/2} & \\ & \chi_1 \nu_v^{-1/2} \end{pmatrix}, \begin{pmatrix} \chi_2 \nu_v^{1/2} & \\ & \chi_2 \nu_v^{-1/2} \end{pmatrix}, \mu_v \right) \rtimes 1,$
- (2) $\left(\begin{pmatrix} \chi_2 \nu_v^{1/2} & \\ & \chi_2 \nu_v^{-1/2} \end{pmatrix}, \begin{pmatrix} \chi_1 \nu_v^{1/2} & \\ & \chi_1 \nu_v^{-1/2} \end{pmatrix}, \mu_v \right) \rtimes 1.$

In the first case, we have $\pi'_{2,v} = \pi_{2,v}$. In the second case, we have $\pi'_{2,v} = {}^\sigma \pi_{2,v}$. Moreover, as in the cuspidal case, we have $\mu_v = \mu'_v$ for almost all v .

Hence, $\pi'_2 = \chi' 1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes_2 \mu$, where χ' is a character of C_E such that χ'_v is equal to χ_v or ${}^\sigma \chi_v$ for almost all v . Applying the partial L -function argument (explained in the proof of Lemma 4.11) to χ as a representation of $\mathrm{GL}(1, \mathbb{A}_E)$, we conclude that χ' is equal to either χ or ${}^\sigma \chi$. The lemma follows. $\square$

4.16. COROLLARY. *Let π_2 be a discrete spectrum automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ which contributes to an ε -trace identity. Then, a discrete spectrum representation $\pi'_2 \neq \pi_2$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity if and only if $\pi_2 \neq {}^\sigma \pi_2$, in which case π'_2 is equal to ${}^\sigma \pi_2$.*

PROOF. The group $\mathbf{H}_2(\mathbb{A}_F)$ is a quotient of $\mathrm{GL}(2, \mathbb{A}_E) \times \mathbb{A}_F^\times$; hence, its discrete spectrum representations are either cuspidal or one dimensional. The corollary follows from Lemmas 4.11 and 4.15. $\square$

If $(h, c) \in \mathbf{H}_2(\mathbb{A}_F)$ is a norm of $g \in \mathbf{G}(\mathbb{A}_F)$, then $(\sigma h, c)$ is also a norm of g . Hence, we expect that if $f \in C(\mathbf{G}(\mathbb{A}_F), \omega)$ and $f_2 \in C(\mathbf{H}_2(\mathbb{A}_F), \omega)$ are matching functions, then $f_2(h, c) = f_2(\sigma h, c)$ for all $(h, c) \in \mathbf{H}_2(\mathbb{A}_F)$. Consequently, if π is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ which is not equal to ${}^\sigma\pi$ and μ is a character of C_F such that $\omega_\pi = \mu \circ \mathrm{N}_{E/F}$, the contributions from $\mathrm{tr}(\pi \otimes_2 \mu)(f_2)$ and $\mathrm{tr}({}^\sigma\pi \otimes_2 \mu)(f_2)$ to an ε -trace identity combine into a single term $2 \cdot \mathrm{tr}(\pi \otimes_2 \mu)(f_2)$.

4.4. Contributions

We now address the question: Under what circumstances are there contributions from both ε -endoscopic groups to an ε -trace identity.

Notation:

- Let Π be an automorphic representation of $\mathbf{G}(\mathbb{A}_F)$. Let f be a function in $C(\mathbf{G}(\mathbb{A}_F), \omega)$. Recall that $\Pi(f \times \varepsilon)$ is defined in Section 3.1. Put

$$\langle \Pi, f \rangle_\varepsilon := \mathrm{tr} \Pi(f \times \varepsilon).$$

If Π is ε -invariant, $\langle \Pi, f \rangle_\varepsilon$ is equal to a product $\prod_v \mathrm{tr} \Pi_v(f_v \times \varepsilon_v)$ of twisted local characters. Put $\langle \Pi_v, f_v \rangle_{\varepsilon_v} := \mathrm{tr} \Pi_v(f_v \times \varepsilon_v)$.

- Let $i = 1$ or 2 . For a representation π_i of $\mathbf{H}_i(\mathbb{A}_F)$ and test function f_i in $C(\mathbf{H}_i(\mathbb{A}_F), \omega)$, put

$$\langle \pi_i, f_i \rangle := \mathrm{tr} \pi_i(f_i).$$

- For any automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_F)$, let $B_{E/F}\tau$ denote the automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ which is obtained by base change from τ (see [L], [F1]).
- Let V be the set of places of F . Let $V^{\mathrm{un}}(E/F)$ denote the set of finite places of F which are unramified in E .
- Let S be a set of places of F . For any adèlic object (automorphic representation, test function, trace, ..., etc.), let subscript S denote the tensor product of local components over the places in S .

Let π be an irreducible admissible representation of a p -adic group H . In [HC1], Harish-Chandra proves the existence of a locally integrable function χ_π on H such that

$$\int_H \chi_\pi(h) f(h) dh = \mathrm{tr} \pi(f), \quad \forall f \in C_c^\infty(H).$$

We call χ_π the **Harish-Chandra character** of π . This notion extends to the ε -twisted characters as follows: Let k be a local field. Let ε_k be a quadratic character of $\mathbf{G}(k)$. Let π be an irreducible, admissible, ε_k -invariant $\mathbf{G}(k)$ -module, i.e. there exists a nontrivial intertwining operator A in $\mathrm{Hom}_{\mathbf{G}(k)}(\pi, \varepsilon_k \pi)$. Note that the space of $\varepsilon_k \pi$ is equal to that of π . The ε_k -**twisted Harish-Chandra character** χ_π^A of π is a function on $\mathbf{G}(k)$ such that

$$\int_{\mathbf{G}(k)} \chi_\pi^A(g) f(g) dg = \mathrm{tr} \pi(f) A, \quad \forall f \in C_c^\infty(\mathbf{G}(k)).$$

It can be shown that χ_π^A is locally constant on the regular set, locally integrable, and unique. It satisfies $\chi_\pi^A(h^{-1}gh) = \varepsilon_k(h) \chi_\pi^A(g)$.

Definition: We say that an irreducible admissible representation of a p -adic group H is **elliptic** if its Harish-Chandra character is not identically zero on the elliptic regular set of H . Likewise, we say that an ε_k -invariant, irreducible, admissible $\mathbf{G}(k)$ -module π is ε_k -**elliptic** if χ_π^A is not identically zero on the elliptic regular set of $\mathbf{G}(k)$. It follows from the

orthogonality relations that each square integrable admissible representation of a connected reductive p -adic group is elliptic. In the twisted case, the ε_k -invariant square integrable representations are similarly elliptic. This too follows from the orthogonality relations of characters.

In order to apply Corollary 4.4 to obtain global lifting results, we fix once and for all two distinct finite places w_1, w_2 of F which are prime in E , and we work with automorphic representations whose local components at w_i ($i = 1, 2$) are elliptic. For $\mathbf{H} = \mathbf{G}$, $\mathbf{H}_1$, or $\mathbf{H}_2$, recall that $E(w_1, w_2, \mathbf{H}(\mathbb{A}_F), \omega)$ denotes the set of functions $f = \otimes_v f_v$ in $C(\mathbf{H}(\mathbb{A}_F), \omega)$ whose local components f_{w_1}, f_{w_2} are elliptic.

For a reductive F -group $\mathbf{H}$ and an automorphic representation π of $\mathbf{H}(\mathbb{A}_F)$, we define the **bad places** for π to be the (finite) set of places of F which is the union of $V - V^{\text{un}}(E/F)$, $\{w_1, w_2\}$, and the set of places v where π_v is ramified (i.e. not unramified).

4.4.1. Base Change. Let τ be a cuspidal non- E -monomial, or one dimensional, automorphic representation of $\text{GL}(2, \mathbb{A}_F)$ such that τ_{w_i} ($i = 1, 2$) is elliptic. Let $\{\Pi\}$ be the unstable (quasi-)packet $[\tau, \varepsilon\tau]$ of $\mathbf{G}(\mathbb{A}_F)$. This is the packet which lifts to the induced representation $I_{(2,2)}(\tau, \varepsilon\tau)$ of $\text{GL}(4, \mathbb{A}_F)$. The table notation was introduced in Section 4.1.

4.17. PROPOSITION. *Let S be the set of bad places for τ . The following table holds for matching test functions with elliptic components at w_1, w_2 and spherical components at all $v \notin S$:*

$$(4.1) \quad \begin{array}{|c|c|c|} \hline G & H_1 & H_2 \\ \hline \{\Pi\} & \tau \otimes_1 1 & B_{E/F}\tau \otimes_2 \omega_\tau \\ \hline \end{array}$$

REMARK. The representation $\tau \otimes_1 \mu \circ N_{E/F}$ is equivalent to $\mu\tau \otimes_1 1$ for any character μ of C_F ; hence, by Hilbert 90, the proposition covers all $\tau \otimes_1 \chi$ where $\chi = {}^\sigma \chi$.

PROOF. For $v \notin S$, let $t(\tau_v)$ be a diagonal matrix in $\text{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes the unramified representation τ_v . From Section 4.2, the global datum $\mathcal{C}(\{\Pi\}, S, \hat{G})$ parametrizing $\{\Pi\}$ is represented by

$$\left[[t(\tau_v), \varepsilon_v(\varpi_v)t(\tau_v)] \right]_{v \notin S} \in \prod_{v \notin S} \hat{G},$$

where ϖ_v is a fixed uniformizer of F_v for each $v \notin S$. By comparing F-H classes, we see that $\tau \otimes_1 1$ and $B_{E/F}\tau \otimes_2 \omega_\tau$ contribute to the ε -trace identity defined by the global datum $\mathcal{C}(\{\Pi\}, S, \hat{G})$. By Theorem 4.10, $\tau \otimes_1 1$ is the only contribution from $\mathbf{H}_1$. By Corollary 4.16, $B_{E/F}\tau \otimes_2 \omega_\tau$ is the only discrete spectrum contribution from $\mathbf{H}_2$. Since the test function f_2 on $\mathbf{H}_2(\mathbb{A}_F)$ has elliptic components, no discretely occurring induced representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity. The proposition follows. $\square$

4.18. COROLLARY. *Let $f \in E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega_\tau)$ and $f_i \in E(w_1, w_2, \mathbf{H}_i, \omega_\tau)$ ($i = 1, 2$) be matching functions whose components at all $v \notin S$ are spherical, the following character identity holds:*

$$(4.2) \quad \sum_{\Pi' \in \{\Pi\}} m(\Pi') \langle \Pi', f \rangle_{\varepsilon, S} = \frac{1}{2} \langle \tau \otimes_1 1, f_1 \rangle_S + \frac{1}{2} \langle B_{E/F}\tau \otimes_2 \omega_\tau, f_2 \rangle_S.$$

Here $m(\Pi')$ is the multiplicity of Π' in the discrete spectrum of $\mathbf{G}(\mathbb{A}_F)$.

REMARK. If $\omega_\tau = 1$, then by [F4, V. 10] the packet $\{\Pi\}$ is the restricted tensor product $\otimes_{v \in V} \{\Pi_v^+, \Pi_v^-\}$, where $\Pi_v^- = 0$ unless τ_v is square integrable. The multiplicity formula is $m(\Pi') = \frac{1}{2}(1 + (-1)^{n(\Pi')})$, where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$.

4.4.2. Exclusive Contributions. We now address under what circumstances do the representations of only one of the ε -endoscopic groups contribute to an ε -trace identity.

4.4.2.1. Case of $\mathbf{H}_1$.

4.19. PROPOSITION. *Let $\pi_1 = \tau \otimes_1 \chi$ be a global (quasi-)packet of representations of $\mathbf{H}_1(\mathbb{A}_F)$ such that:*

- τ is a non- E -monomial automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$;
- τ_{w_i} ($i = 1, 2$) is elliptic;
- $\chi \neq {}^\sigma \chi$.

If π_1 contributes to an ε -trace identity for matching functions f in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and f_i in $E(w_1, w_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$ ($i = 1, 2$), then no representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity.

The proof is given after Lemma 4.23.

Suppose an automorphic representation π_2 of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the ε -trace identity. Let S be the union of the respective sets of bad places for π_1, π_2 . That is, $\pi_{1,v}, \pi_{2,v}$ are unramified for all $v \notin S$. Let $\{\Pi\}$ be the global (quasi-)packet of representations of $\mathbf{G}(\mathbb{A}_F)$ whose global datum is the lift of $\mathcal{C}(\pi_1, S, {}^L H_1)$. Put ${}^\sigma \pi_1 := \tau \otimes_1 {}^\sigma \chi$. By Corollary 4.4, Theorem 4.10, and Corollary 4.16, we have:

$$(4.3) \quad \sum_{\Pi' \in \{\Pi\}} m(\Pi') \langle \Pi', f \rangle_{\varepsilon, S} = \frac{1}{2} \langle \pi_1, f_1 \rangle_S + \frac{1}{2} \langle {}^\sigma \pi_1, f_1 \rangle_S \\ + \frac{1}{2} \langle \pi_2, f_2 \rangle_S + \frac{d}{2} \langle {}^\sigma \pi_2, f_2 \rangle_S,$$

where $m(\Pi')$ is the multiplicity of Π' , and $d = \begin{cases} 1 & \text{if } \pi_2 \neq {}^\sigma \pi_2, \\ 0 & \text{otherwise.} \end{cases}$

4.20. LEMMA. *The (quasi-)packet $\{\Pi\}$ is stable.*

PROOF. Suppose $\{\Pi\}$ is unstable. Then, according to Lemma 4.9, one of the following holds:

- (1) $\{\Pi\} = [\pi, \varepsilon\pi]$, where π is a cuspidal non- E -monomial automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$.
- (2) $\{\Pi\} = [\mu 1_2, \varepsilon \mu 1_2]$ for some character μ of C_E .
- (3) $\{\Pi\} = [\pi(\theta\chi), \pi(\theta {}^\sigma \chi)]$, where θ, χ are characters of C_E such that none of $\theta, \chi, \theta\chi, \theta {}^\sigma \chi$ is fixed by the action of $\mathrm{Gal}(E/F)$.

In the first (resp. second) case, we know from Proposition 4.17 that $\{\Pi\}$ is the lift of the packet $\pi'_1 = \pi \otimes_1 1$ (resp. $\mu 1_2 \otimes_1 1$) of $\mathbf{H}_1(\mathbb{A}_F)$. Since $\chi \neq {}^\sigma \chi$, by Theorem 4.10, π'_1 and π_1 cannot both contribute to the trace identity, whence a contradiction.

In the third case, one can see via a comparison of F-H classes that $\{\Pi\}$ is the lift of $\pi(\theta) \otimes_1 \chi$. Since we assume that τ is not E -monomial, we have a contradiction by Theorem 4.10. $\square$

4.21. COROLLARY. *Each member of $\{\Pi\}$ occurs with multiplicity one in the discrete spectrum of $\mathrm{GSp}(2, \mathbb{A}_F)$.*

Let $\{\{\Pi_v\}\}_{v \in V}$ be the collection of local packets such that $\{\Pi\}$ is the restricted tensor product $\otimes_{v \in V} \{\Pi_v\}$. For each member $\Pi' = \otimes_v \Pi'_v$ of $\{\Pi\}$, $\langle \Pi', f \rangle_\varepsilon$ is equal to a product $\prod_v \langle \Pi'_v, f_v \rangle_{\varepsilon_v}$ of twisted traces. For simplicity, we often drop the subscript v from ε_v if the meaning is clear. By Lemma 4.20, we have:

$$\sum_{\Pi' \in \{\Pi\}} m(\Pi') \langle \Pi', f \rangle_{\varepsilon, S} = \prod_{v \in S} \langle \{\Pi_v\}, f_v \rangle_\varepsilon,$$

where $\langle \{\Pi_v\}, f_v \rangle_\varepsilon := \sum_{\Pi'_v \in \{\Pi'_v\}} \langle \Pi'_v, f_v \rangle_\varepsilon$.

Let $S_0 = S - \{w_1, w_2\}$. Let

$$C_{S_0} = \langle \{\Pi\}, f \rangle_{\varepsilon, S_0}, \quad D_{1, S_0} = \langle \pi_1, f_1 \rangle_{S_0}, \quad D_{2, S_0} = \langle \pi_2, f_2 \rangle_{S_0}.$$

From the matching condition on f, f_1, f_2 , we have $\langle \pi_1, f_1 \rangle = \langle {}^\sigma \pi_1, f_1 \rangle$ and $\langle \pi_2, f_2 \rangle = \langle {}^\sigma \pi_2, f_2 \rangle$. We rewrite (4.3) as follows:

$$(4.4) \quad C_{S_0} \langle \{\Pi_{w_1}\}, f_{w_1} \rangle_\varepsilon \langle \{\Pi_{w_2}\}, f_{w_2} \rangle_\varepsilon \\ = D_{1, S_0} \langle \pi_{1, w_1}, f_{1, w_1} \rangle \langle \pi_{1, w_2}, f_{1, w_2} \rangle + n \cdot D_{2, S_0} \langle \pi_{2, w_1}, f_{2, w_1} \rangle \langle \pi_{2, w_2}, f_{2, w_2} \rangle,$$

where $n = \frac{1}{2}$ or 1, depending on whether π_2 is equal to ${}^\sigma \pi_2$ or not. Here, for each place v , the symbol $\pi_{1, v}$ denotes the local packet of $H_{1, v}$ such that $\pi_1 = \otimes_v \pi_{1, v}$.

4.22. DEFINITION. Let v be a place of F which is prime in E . We say that an elliptic regular element in G_v is of **type** i ($i = 1, 2$) if its norms lie only in $H_{i, v}$. We say that an elliptic function f_v on G_v is of **type** i if the orbital integral of f_v is nonzero only at the elliptic regular elements of type i .

From the norm correspondence among elliptic regular elements (computed in Appendix B), type i ($i = 1, 2$) elements exist. Consequently, type i functions exist.

4.23. LEMMA. *There exists a type 1 elliptic function f_{w_2} on G_{w_2} such that the trace $\langle \{\Pi\}_{w_2}, f_{w_2} \rangle_\varepsilon$ is nonzero.*

PROOF. Let f_{w_1} be a type 1 elliptic function such that $\langle \pi_{1, w_1}, f_{1, w_1} \rangle \neq 0$. This is possible because by hypothesis π_{1, w_1} is elliptic. Then, we may choose f_{2, w_1} to be zero and obtain the following equation:

$$C_{S_0} \langle \{\Pi_{w_1}\}, f_{w_1} \rangle_\varepsilon \langle \{\Pi_{w_2}\}, f_{w_2} \rangle_\varepsilon = D_{1, S_0} \langle \pi_{1, w_1}, f_{1, w_1} \rangle \langle \pi_{1, w_2}, f_{1, w_2} \rangle.$$

For each $v \in S_0$, choose f_v and matching $f_{1, v}$ such that $\langle \pi_{1, v}, f_{1, v} \rangle \neq 0$. Since by assumption π_{1, w_2} is elliptic, there exists a type 1 elliptic f_{w_2} and matching f_{1, w_2} such that $\langle \pi_{1, w_2}, f_{1, w_2} \rangle \neq 0$. The lemma follows. $\square$

PROOF OF PROPOSITION 4.19. Let f_{w_1} be a type 2 elliptic function on G_{w_1} . Let f_{1, w_1} be zero. Then, (4.4) becomes

$$(4.5) \quad C_{S_0} \langle \{\Pi_{w_1}\}, f_{w_1} \rangle_\varepsilon \langle \{\Pi_{w_2}\}, f_{w_2} \rangle_\varepsilon = n D_{2, S_0} \langle \pi_{2, w_1}, f_{2, w_1} \rangle \langle \pi_{2, w_2}, f_{2, w_2} \rangle.$$

Let f_{w_2} be a type 1 function on G_{w_2} such that $\langle \{\Pi\}_{w_2}, f_{w_2} \rangle_\varepsilon$ is nonzero. It exists by Lemma 4.23. Let f_{2, w_2} be zero. Then,

$$C_{S_0} \langle \{\Pi_{w_1}\}, f_{w_1} \rangle_\varepsilon = 0$$

for all elliptic functions f_{w_1} of type 2.

Now let f_{w_2} be an arbitrary test function on G_{w_2} . The equation (4.5), where f_{w_1} is of type 2, implies that

$$(4.6) \quad 0 = C_{S_0} \langle \{\Pi_{w_1}\}, f_{w_1} \rangle_\varepsilon \langle \{\Pi_{w_2}\}, f_{w_2} \rangle_\varepsilon = n D_{2, S_0} \langle \pi_{2, w_1}, f_{2, w_1} \rangle \langle \pi_{2, w_2}, f_{2, w_2} \rangle.$$

There are two possibilities: (i) $\langle \pi_{2,w_2}, f_{2,w_2} \rangle = 0$ for every f_{2,w_2} matching some elliptic function f_{w_2} on G_{w_2} ; (ii) $\langle \pi_{2,w_1}, f_{2,w_1} \rangle = 0$ for every f_{2,w_1} matching some type 2 elliptic function on G_{w_1} . If case (i) holds, we are done. Suppose case (ii) holds. For any $h_2 = (g, c) \in H_{2,w_1}$, put $\sigma h_2 := (\sigma g, c)$. Since f_{2,w_1} matches a function on G_{w_1} , we have $f_{2,w_1}(h_2) = f_{2,w_1}(\sigma h_2)$ for all $h_2 \in H_{2,w_1}$. Hence,

$$0 = \langle \pi_{2,w_1}, f_{2,w_1} \rangle = \int_{\langle \sigma \rangle \backslash H_{2,w_1}} (\chi_{\pi_{2,w_1}}(h_2) + \chi_{\pi_{2,w_1}}(\sigma h_2)) f_{2,w_1}(h_2) dh_2.$$

In particular, $\chi_{\pi_{2,w_1}}(h_2) + \chi_{\pi_{2,w_1}}(\sigma h_2) = 0$ for all elliptic elements h_2 in H_{2,w_1} . It follows that if f_{w_1} is any elliptic function on G_{w_1} , then $\langle \pi_{2,w_1}, f_{2,w_1} \rangle = 0$ for any function f_{2,w_1} on H_{2,w_1} which matches f_{w_1} . The proposition follows. $\square$

4.24. LEMMA. *Let χ be a character of C_E such that $\chi \neq {}^\sigma \chi$. Suppose the one dimensional representation $\pi_1 = 1_2 \otimes_1 \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to an ε -trace identity. Then, no automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity.*

PROOF. Let S be the finite set of bad places for π_1 . By the Chebotarev Density Theorem, there exists a place $v \notin S$ which splits into two places v_1, v_2 of E such that $\chi_{v_1} \neq \chi_{v_2}$. The F-H class in $\hat{H}_1$ parametrizing $\pi_{1,v}$ lifts to the conjugacy class in $\mathrm{GSp}(2, \mathbb{C})$ represented by

$$\hat{g} = \begin{pmatrix} \chi_{v_1} \nu_v^{1/2} & & & \\ & \chi_{v_1} \nu_v^{-1/2} & & \\ & & \chi_{v_2} \nu_v^{1/2} & \\ & & & \chi_{v_2} \nu_v^{-1/2} \end{pmatrix}.$$

Suppose a representation $\pi_2 = \tau \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity, where τ is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ and μ is a character of C_F such that $\omega_\tau = \mu \circ N_{E/F}$. Then, τ_{v_1} must be parametrized by the conjugacy class in $\mathrm{GL}(2, \mathbb{C})$ containing either $\mathrm{diag}(\chi_{v_1} \nu_v^{1/2}, \chi_{v_2} \nu_v^{-1/2})$ or $\mathrm{diag}(\chi_{v_2} \nu_v^{1/2}, \chi_{v_1} \nu_v^{-1/2})$. In either case, since $\chi_{v_1} \neq \chi_{v_2}$, τ_{v_1} is not unitarizable, which is a contradiction because all local components of a unitary automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ are unitarizable. $\square$

Let $R_{E/F}\mathrm{GL}(2)$ be the F -group obtained from $\mathrm{GL}(2)$ on restricting scalars from E to F . It is an ε -twisted endoscopic group of $\mathrm{GL}(4)$, and its group of $\mathbb{A}_F$ -points is $\mathrm{GL}(2, \mathbb{A}_E)$. For an automorphic representation τ_E of $\mathrm{GL}(2, \mathbb{A}_E)$, let $\pi(\tau_E)$ denote the automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$ which is obtained from τ_E via the twisted endoscopic lifting from $R_{E/F}\mathrm{GL}(2)$ to $\mathrm{GL}(4)$ (see [AC, Sect. 3.6]). The representation $\pi(\tau_E)$ is cuspidal if and only if τ_E is cuspidal and $\tau_E \not\cong {}^\sigma \tau_E$.

In the case of $\mathbf{H}_1$, we in general denote a (quasi-)packet by π_1 , i.e. without braces. For a (quasi-)packet π_1 of $\mathbf{H}_1(\mathbb{A}_F)$, let $\xi_1^*(\pi_1)$ denote the (quasi-)packet of $\mathbf{G}(\mathbb{A}_F)$ which is the lift of π_1 .

4.25. LEMMA. *Let τ be a cuspidal or one dimensional automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. Let χ be a character of C_E . Then, the (quasi-)packet $\xi_1^*(\tau \otimes_1 \chi)$ of $\mathbf{G}(\mathbb{A}_F)$ lifts to the automorphic representation $\pi(\chi B_{E/F} \tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.*

PROOF. Let S be the union of the sets of bad places for τ, χ . Hence, τ_v, χ_v are unramified for all $v \notin S$. For each $v \notin S$, let $t(\tau_v) = \mathrm{diag}(a_v, b_v)$ be a diagonal matrix in $\mathrm{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes τ_v . The global datum in $\widehat{\mathrm{GL}(4)} = \mathrm{GL}(4, \mathbb{C})$ parametrizing $\pi(\chi B_{E/F} \tau)$ is as follows:

- Suppose $v \notin S$ is prime in E . Let χ'_v be an unramified character of $F_v^\times$ such that $\chi_v = \chi'_v \circ N_{E/F}$. Then,

$$\mathcal{C}(\pi(\chi B_{E/F}\tau), S, \mathrm{GL}(4, \mathbb{C}))_v \ni \begin{pmatrix} \chi'_v a_v & & & \\ & \chi'_v b_v & & \\ & & -\chi'_v a_v & \\ & & & -\chi'_v b_v \end{pmatrix}.$$

- Suppose $v \notin S$ splits into two places v_1, v_2 of E . Then,

$$\mathcal{C}(\pi(\chi B_{E/F}\tau), S, \mathrm{GL}(4, \mathbb{C}))_v \ni \begin{pmatrix} \chi_{v_1} a_v & & & \\ & \chi_{v_1} b_v & & \\ & & \chi_{v_2} a_v & \\ & & & \chi_{v_2} b_v \end{pmatrix}.$$

The global datum $\mathcal{C}(\xi_1^*(\tau \otimes_1 \chi), S, \mathrm{GSp}(2, \mathbb{C}))$ is described by Corollary 4.6. The proposition follows on comparing the datum $\mathcal{C}(\pi(\chi B_{E/F}\tau), S, \mathrm{GL}(4, \mathbb{C}))$ with the natural image of $\mathcal{C}(\xi_1^*(\tau \otimes_1 \chi), S, \mathrm{GSp}(2, \mathbb{C}))$ in $\prod_{v \notin S} \mathrm{GL}(4, \mathbb{C})$. $\square$

4.26. LEMMA. *Let τ be an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. Let χ be a character of C_E . The automorphic representation $\pi = \chi B_{E/F}\tau$ of $\mathrm{GL}(2, \mathbb{A}_E)$ is invariant under $\mathrm{Gal}(E/F)$ if and only if there exists a (possibly trivial) quadratic character ε' of C_F such that: (i) ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$, and (ii) τ is equivalent to $\varepsilon'\tau$ or $\varepsilon'\varepsilon\tau$.*

PROOF. Suppose $\pi \cong {}^\sigma\pi$. Then, the central character $\omega_\pi = \chi^2 \cdot (\omega_\tau \circ N_{E/F})$ of π must be equal to $\omega_{{}^\sigma\pi} = {}^\sigma\chi^2 \cdot (\omega_\tau \circ N_{E/F})$. Hence, $\chi^2 = {}^\sigma\chi^2$, which by Hilbert 90 implies that ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ for some (possibly trivial) quadratic character ε' of C_F . We have:

$$\chi B_{E/F}(\varepsilon'\tau) = \chi \cdot (\varepsilon' \circ N_{E/F}) B_{E/F}\tau = {}^\sigma\chi B_{E/F}\tau \cong \chi B_{E/F}\tau.$$

By [L], [F1], $B_{E/F}(\varepsilon'\tau) \cong B_{E/F}\tau$ implies that τ is equivalent to $\varepsilon'\tau$ or $\varepsilon'\varepsilon\tau$.

Suppose ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ for some quadratic character ε' of C_F , and τ is equivalent to $\varepsilon'\tau$ or $\varepsilon'\varepsilon\tau$. Then, ${}^\sigma\pi = \chi \cdot (\varepsilon' \circ N_{E/F}) B_{E/F}\tau = \chi B_{E/F}(\varepsilon'\tau)$, which is equivalent to either $\chi B_{E/F}\tau$ or $\chi B_{E/F}(\varepsilon\tau)$. Since $B_{E/F}(\varepsilon\tau) \cong B_{E/F}\tau$, the lemma follows. $\square$

4.27. PROPOSITION. *Let τ be a cuspidal, non- E -monomial, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, χ be a character of C_F , such that: There does not exist a character ε' of C_F such that ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ and $\tau \cong \varepsilon'\tau$. Then, each member of $\{\Pi\} = \xi_1^*(\tau \otimes_1 \chi)$ is a cuspidal automorphic representation of $\mathbf{G}(\mathbb{A}_F)$.*

PROOF. By Lemma 4.25, $\{\Pi\}$ lifts to $\pi(\chi B_{E/F}\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$. Since τ is not E -monomial, $\chi B_{E/F}\tau$ is cuspidal. By Lemma 4.26 and [AC, Sect. 3.6], we conclude that the automorphic representation $\pi(\chi B_{E/F}\tau)$ is cuspidal.

Suppose a representation Π in $\{\Pi\}$ is not cuspidal. Then, Π is an irreducible constituent of a parabolically induced representation I of $\mathbf{G}(\mathbb{A}_F)$. By the results in [F4, Sect. V. 1], I lifts to a parabolically induced representation of $\mathrm{GL}(4, \mathbb{A}_F)$. By the strong multiplicity one (or rigidity) theorem for $\mathrm{GL}(4)$, we conclude that the cuspidal representation $\pi(\chi B_{E/F}\tau)$ is an irreducible constituent of a parabolically induced representation of $\mathrm{GL}(4, \mathbb{A}_F)$, which is a contradiction. $\square$

The case where ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ and $\tau \cong \varepsilon'\tau$ will be addressed by Proposition 4.36 in Section 4.5.

4.4.2.2. *Case of $\mathbf{H}_2$.* For any character θ of C_E , recall that $\pi(\theta)$ denotes the E -monomial representation of $\mathrm{GL}(2, \mathbb{A}_F)$ associated with θ .

4.28. CLAIM. *Let τ be an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ such that $\tau \neq {}^\sigma\tau$ and $\omega_\tau = \mu \circ \mathrm{N}_{E/F}$ for some character μ of C_F . Suppose the representation $\pi_2 = \tau \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to an ε -trace identity. Then, no discrete spectrum packet of $\mathbf{H}_1(\mathbb{A}_F)$ of the form $\pi(\theta) \otimes_1 \chi$, where θ, χ are characters of C_E , contributes to the trace identity.*

PROOF. Let S be the set of bad places for π_2 . For a place $v \notin S$ which does not split in E , μ_v is an unramified character of $F_v^\times$, and τ_v is an induced representation of $\mathrm{GL}(2, E_v)$ of the form $I(\eta_v \circ \mathrm{N}_{E/F}, (\mu_v \eta_v^{-1}) \circ \mathrm{N}_{E/F})$, where η_v is an unramified character of $F_v^\times$. From Section 4.2, we see that the F-H class in ${}^L H_2$ parametrizing $\pi_{2,v}$ lifts to the conjugacy class in $\hat{G}$ represented by

$$\hat{g}_v = \mathrm{diag}(\eta_v, -\eta_v, -\mu_v \eta_v^{-1}, \mu_v \eta_v^{-1}).$$

If $v \notin S$ splits into two places v_1, v_2 of E , then the F-H class parametrizing $\pi_{2,v}$ lifts to the conjugacy class in $\hat{G}$ represented by

$$\hat{g}_v = [t(\tau_{v_1}), t(\tau_{v_2})],$$

where $t(\tau_{v_i})$ ($i = 1, 2$) is a diagonal matrix whose conjugacy class in $\mathrm{GL}(2, \mathbb{C})$ parametrizes the unramified representation τ_{v_i} of $\mathrm{GL}(2, E_{v_i}) = \mathrm{GL}(2, F_v)$.

A lifting of automorphic representations of $\mathrm{GL}(2, \mathbb{A}_E)$ to ε -invariant automorphic representations of $\mathrm{GL}(4, \mathbb{A}_F)$ is described in [AC, Sect. 3.6]. In particular, if an automorphic representation π of $\mathrm{GL}(2, \mathbb{A}_E)$ satisfies $\pi \neq {}^\sigma\pi$, then it lifts to a cuspidal automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$.

Viewing $\mathrm{GSp}(2, \mathbb{C})$ as a subgroup of $\mathrm{GL}(4, \mathbb{C})$, the global datum in $\prod_{v \notin S} \mathrm{GL}(4, \mathbb{C})$ represented by $[\hat{g}_v]_{v \notin S}$ parametrizes the automorphic $\mathrm{GL}(4, \mathbb{A}_F)$ -module Π which is the lift of τ . Since $\tau \neq {}^\sigma\tau$, Π is cuspidal.

On the other hand, $\pi(\theta) \otimes_1 \chi$ lifts to a packet of $\mathbf{G}(\mathbb{A}_F)$ which in turn lifts to the induced representation $I_{(2,2)}(\pi(\theta\chi), \pi(\theta {}^\sigma\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. By the rigidity theorem for $\mathrm{GL}(4)$, the claim follows. $\square$

Recall that we have fixed two finite places w_1, w_2 of F which are prime in E .

4.29. LEMMA. *Let τ be a discrete spectrum representation of $\mathrm{GL}(2, \mathbb{A}_F)$ with the property that τ_{w_1} and τ_{w_2} are elliptic. Suppose the discrete spectrum representation $B_{E/F}\tau \otimes_2 \omega_\tau \varepsilon$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to an ε -trace identity with respect to matching f, f_i ($i = 1, 2$) in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$, $E(w_1, w_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$, respectively. Then, no discrete spectrum packet of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity.*

PROOF. Suppose a packet $\pi_1 = \pi \otimes_1 \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity, where π is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, and χ is a character of C_E . There are two cases to consider: The case where $\chi = {}^\sigma\chi$ and the case where $\chi \neq {}^\sigma\chi$.

Suppose $\chi = {}^\sigma\chi$. Then, $\chi = \mu \circ \mathrm{N}_{E/F}$ for some character μ of C_F , and $\pi_1 = \mu\pi \otimes_1 1$. By Proposition 4.17, the representation $B_{E/F}(\mu\pi) \otimes_2 \omega_{\mu\pi}$ of $\mathbf{H}_2(\mathbb{A}_F)$ must also contribute to the trace identity. By Corollary 4.16, $B_{E/F}\tau \otimes_1 \omega_\tau \varepsilon$ must be equivalent to $B_{E/F}(\mu\pi) \otimes_1 \omega_{\mu\pi}$. The theory of base change for $\mathrm{GL}(2)$ (see [L], [F1]) implies that τ is equivalent to either $\mu\pi$ or $\varepsilon\mu\pi$. In either case, $\omega_\tau \varepsilon$ is equal to $\omega_{\mu\pi} \varepsilon \neq \omega_{\mu\pi}$, whence a contradiction.

Suppose $\chi \neq {}^\sigma\chi$. If $\pi_1 = \pi \otimes_1 \chi$ contributes to the trace identity, then π_{w_1}, π_{w_2} must be elliptic, or else the condition that the functions f_{1,w_1}, f_{1,w_2} are elliptic would imply

that $\langle \pi_1, f_1 \rangle = 0$. If π_{w_1}, π_{w_2} are elliptic, then by Proposition 4.19 and Claim 4.28 no automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ may contribute to the trace identity. The lemma follows. $\square$

4.30. PROPOSITION. *Let μ be a character of C_F . Let τ be a discrete spectrum automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ with the following properties:*

- $\tau \neq {}^\sigma \tau$;
- $\omega_\tau = \mu \circ \mathrm{N}_{E/F}$;
- τ_{w_1} and τ_{w_2} are elliptic.

Suppose the automorphic representation $\pi_2 = \tau \otimes_2 \mu$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to an ε -trace identity with respect to matching test functions with elliptic local components at w_1, w_2 . Then, no discrete spectrum packet of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity.

PROOF. Suppose a packet π_1 of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity. Since $\tau \neq {}^\sigma \tau$, by Claim 4.28 the packet π_1 is not of the form $\pi(\theta) \otimes_1 \chi$ for any characters θ, χ of C_E . By Proposition 4.17 and Lemma 4.11, π_1 is not of the form $\pi \otimes_1 1$, where π is an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$.

Consequently, by Lemma 4.20, the packet $\{\Pi\} = \otimes_v \{\Pi_v\}$ of $\mathbf{G}(\mathbb{A}_F)$ which contributes to the trace identity must be stable; namely, every member of $\{\Pi\}$ occurs with multiplicity one in the discrete spectrum of $\mathbf{G}(\mathbb{A}_F)$.

The argument we then use mirrors completely the one used in the proof of Proposition 4.19. Thus, we shall content ourselves with giving only a sketch: First, we show that there exists a test function f_{w_2} on G_{w_2} such that $\langle \{\Pi_{w_2}\}, f_{w_2} \rangle_\varepsilon$ is nonzero. Then, we show that either π_1 does not exist or π_{1, w_1} is non-elliptic. Since by assumption the test function f_{1, w_1} on H_{1, w_1} is elliptic, the proposition follows. $\square$

4.31. LEMMA. *Let χ be a character of C_E such that $\chi \neq {}^\sigma \chi$ but $\chi^2 = \mu \circ \mathrm{N}_{E/F}$ for some character μ of C_F . Suppose the one dimensional representation $\pi_2 = (\chi \circ \det) \otimes_2 \mu = \chi|_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes_2 \mu$ contributes to an ε -trace identity. Then, no global packet of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity.*

PROOF. Recall from Chapter 2 that

$$\hat{H}_1 = (\mathrm{GL}(2, \mathbb{C}) \times \mathbb{C}^\times \times \mathbb{C}^\times) / \{(\mathrm{diag}(z, z)^{-1}, z, z) : z \in \mathbb{C}^\times\},$$

and ${}^L H_1 = \hat{H}_1 \rtimes \mathrm{Gal}(E/F)$, where the action of σ swaps the two $\mathbb{C}^\times$ -components. Suppose a packet $\pi_1 = \tau \otimes_1 \theta$ of $\mathbf{H}_1(\mathbb{A}_F)$ contributes to the trace identity. By the Chebotarev Density Theorem, there exists a place v of F which splits in E such that $\pi_{1, v}$ and $\pi_{2, v}$ are unramified. Suppose v splits into two places v_1, v_2 of E . Since χ^2 is invariant under the action of $\mathrm{Gal}(E/F)$, the character $\kappa_v := \frac{\chi_{v_1}}{\chi_{v_2}}$ is quadratic. The F-H class parametrizing $\pi_{2, v}$ lifts to the conjugacy class in $\hat{G}$ represented by

$$\begin{aligned} \hat{g}_v &= \mathrm{diag}(\chi_{v_1} \nu_v^{1/2}, \chi_{v_2} \nu_v^{1/2}, \chi_{v_2} \nu_v^{-1/2}, \chi_{v_1} \nu_v^{-1/2}) \\ &= \begin{pmatrix} \chi_{v_1} \nu_v^{1/2} & & & \\ & \chi_{v_2} \nu_v^{1/2} & & \\ & & \nu_v^{-1} \kappa_v \chi_{v_1} \nu_v^{1/2} & \\ & & & \nu_v^{-1} \kappa_v \chi_{v_2} \nu_v^{1/2} \end{pmatrix} \\ &\stackrel{(23)}{\sim} \begin{pmatrix} \chi_{v_1} \nu_v^{1/2} & & & \\ & \chi_{v_2} \nu_v^{-1/2} & & \\ & & \chi_{v_2} \nu_v^{1/2} & \\ & & & \chi_{v_1} \nu_v^{-1/2} \end{pmatrix}. \end{aligned}$$

From Section 4.2, the F-H class in ${}^L H_1$ parametrizing $\pi_{1,v}$ is represented by

$$\hat{h}_v = (t(\tau_v), \theta_{v_1}, \theta_{v_2}) \rtimes 1,$$

where $t(\tau_v)$ is a diagonal matrix in $\mathrm{GL}(2, \mathbb{C})$ whose conjugacy class parametrizes τ_v . Note that the Galois component of $\hat{h}_v$ is trivial because $\mathbf{H}_1$ is split over F_v . Since π_1 contributes to the trace identity, $\hat{h}_v$ lifts via the embedding $\xi_1 : {}^L H_1 \rightarrow \hat{G}$ (see Chapter 2) to the conjugacy class of $\hat{g}_v$ in $\hat{G}$. In other words, the element

$$\xi_1(\hat{h}_v) = \begin{pmatrix} \theta_{v_1} t(\tau_v) & \\ & \theta_{v_2} t(\tau_v) \end{pmatrix} \in \hat{G}$$

is conjugate to $\hat{g}_v$. Hence, $\pi_{1,v}$ is parametrized by the conjugacy class of one of the following types of elements in ${}^L H_1$:

- (1) $(\mathrm{diag}(\chi_{v_1} \nu_v^{1/2}, \chi_{v_2} \nu_v^{1/2}), \eta_1, \eta_2) \rtimes 1$, where the unordered set $\{\eta_1, \eta_2\}$ is equal to $\{1, \kappa_v \nu_v^{-1}\}$ (Notice that there is no minus sign in front of the $1/2$ in $\chi_{v_2} \nu_v^{1/2}$);
- (2) $(\mathrm{diag}(\zeta_1 \nu_v^{1/2}, \zeta_2 \nu_v^{-1/2}), \eta_1, \eta_2) \rtimes 1$, where the unordered set $\{\zeta_1, \zeta_2\}$ is equal to $\{\chi_{v_1}, \chi_{v_2}\}$, and $\{\eta_1, \eta_2\} = \{1, \kappa_v\}$.

In the first case, assume without loss of generality that $\eta_1 = 1$ and $\eta_2 = \kappa_v \nu_v^{-1}$. Then, there exists a quasi-character λ_v of $F_v^\times$ such that $\tau_v = \lambda_v \nu_v^{1/2} I(\chi_{v_1}, \chi_{v_2})$, $\theta_{v_1} = \lambda_v^{-1}$, and $\theta_{v_2} = \lambda_v^{-1} \kappa_v \nu_v^{-1}$. Since τ is a unitary automorphic representation of the group $\mathrm{GL}(2, \mathbb{A}_F)$, τ_v must be unitarizable, which implies that λ_v is non-unitary. Since $\theta_{v_1} = \lambda_v^{-1}$ and θ is unitary, we have a contradiction.

In the second case, assume without loss of generality that $\eta_1 = 1$ and $\eta_2 = \kappa_v$. Then, there exists a quasi-character λ_v of C_{F_v} such that $\tau_v = \lambda_v I(\zeta_1 \nu_v^{1/2}, \zeta_2 \nu_v^{-1/2})$, $\theta_{v_1} = \lambda_v^{-1}$, and $\theta_{v_2} = \lambda_v^{-1} \kappa_v$. The condition $\chi \neq {}^\sigma \chi$ implies that $\zeta_1 \neq \zeta_2$; hence, λ_v must be non-unitary for τ_v to be unitarizable. But then $\theta_{v_1} = \lambda_v^{-1}$ is non-unitary, which is a contradiction. $\square$

For an automorphic representation π_2 of the group $\mathbf{H}_2(\mathbb{A}_F)$, let $\xi_2^*(\pi_2)$ denote the (quasi-)packet of $\mathbf{G}(\mathbb{A}_F)$ which is the lift of π_2 . Recall that, for an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$, $\pi(\tau)$ denotes the automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$ obtained from τ via the twisted endoscopic lifting from $R_{E/F} \mathrm{GL}(2)$ to $\mathrm{GL}(4)$.

4.32. LEMMA. *Let τ be an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$, μ be a character of C_F , such that $\tau \not\cong {}^\sigma \tau$ and $\omega_\tau = \mu \circ N_{E/F}$. The (quasi-)packet $\{\Pi\} = \xi_2^*(\tau \otimes_2 \mu)$ lifts to the automorphic representation $\pi(\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.*

REMARK. The (quasi-)packets $\xi_2^*(\tau \otimes_2 \mu)$ and $\xi_2^*(\tau \otimes_2 \mu\varepsilon)$ are inequivalent, but they both lift to $\pi(\tau)$.

PROOF. Let S be the set of bad places for $\tau \otimes_2 \mu$. At a place $v \notin S$ which is prime in E , τ_v is the induced representation $I(\alpha_v, \beta_v)$ for some unramified characters α_v, β_v of $E_v^\times$. Since α_v, β_v are unramified, there exist characters a_v, b_v of $F_v^\times$ such that $\alpha_v = a_v \circ N_{E/F}$ and $\beta_v = b_v \circ N_{E/F}$. The condition $\omega_\tau = \mu \circ N_{E/F}$ implies that $a_v^2 b_v^2 = \mu_v^2$. Consequently, $a_v b_v(\varpi_v) = \pm \mu_v(\varpi_v)$ (Here, ϖ_v is a fixed uniformizer of F_v). The F-H class $\mathcal{C}(\pi(\tau), S, \mathrm{GL}(4, \mathbb{C}))_v$ parametrizing $\pi(\tau)_v$ is represented by

$$\hat{h}_v = \begin{pmatrix} a_v & & & \\ & b_v & & \\ & & -a_v & \\ & & & -b_v \end{pmatrix} = \begin{pmatrix} a_v & & & \\ & \pm \mu_v / a_v & & \\ & & -a_v & \\ & & & \mp \mu_v / a_v \end{pmatrix} \in \mathrm{GL}(4, \mathbb{C}).$$

From Corollary 4.8, the conjugacy class in $\mathrm{GSp}(2, \mathbb{C})$ parametrizing the unramified component of $\{\Pi_v\}$ is represented by

$$\hat{g}_v = \begin{pmatrix} & \mu_v^{1/2} & & \\ \mu_v^{-1/2} a_v^2 & & & \\ & & \mu_v^{-1/2} b_v^2 & \\ \mu_v^{1/2} & & & \end{pmatrix} \sim \begin{pmatrix} \mu_v/a_v & & & \\ & -\mu_v/a_v & & \\ & & -a_v & \\ & & & a_v \end{pmatrix}.$$

Hence, the image of $\hat{g}_v$ in $\mathrm{GL}(4, \mathbb{C})$ is conjugate to $\hat{h}_v$.

Suppose $v \notin S$ splits into two places v_1, v_2 of E . Then, $\mathrm{GL}(2, E_{v_i}) = \mathrm{GL}(2, F_v)$ ($i = 1, 2$). Let $\tau_i = \tau_{v_i}$. Let $t(\tau_i)$ be a diagonal matrix in $\mathrm{GL}(2, \mathbb{C})$ parametrizing the unramified representation τ_i of $\mathrm{GL}(2, F_v)$. The conjugacy class $\mathcal{C}(\pi(\tau), S, \mathrm{GL}(4, \mathbb{C}))_v$, which parametrizes $\pi(\tau)_v$, is represented by the element $\hat{h}_v = \mathrm{blockdiag}(t(\tau_1), t(\tau_2))$, while the conjugacy class in $\mathrm{GSp}(2, \mathbb{C})$ parametrizing the unramified component of $\{\Pi_v\}$ is represented by $\hat{g}_v = [t(\tau_1), t(\tau_2)]$. The image of $\hat{g}_v$ in $\mathrm{GL}(4, \mathbb{C})$ is conjugate to $\hat{h}_v$.

The lemma follows. $\square$

4.33. PROPOSITION. *Let τ be an automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$, μ be a character of C_F , such that $\tau \not\cong {}^\sigma\tau$ and $\omega_\tau = \mu \circ N_{E/F}$. Then, every automorphic representation in $\{\Pi\} = \xi_2^*(\tau \otimes_2 \mu)$ is cuspidal.*

PROOF. By lemma 4.32, the (quasi-)packet $\{\Pi\}$ lifts to the automorphic representation $\pi(\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$. By the assumption that $\tau \not\cong {}^\sigma\tau$, the representation $\pi(\tau)$ is cuspidal.

Suppose a representation Π in $\{\Pi\}$ is not cuspidal. Then, it is an irreducible constituent of a parabolically induced representation, which by [F4, Sect. V. 1] lifts to a parabolically induced representation of $\mathrm{GL}(4, \mathbb{A}_F)$. By the rigidity theorem for $\mathrm{GL}(4)$, we obtain the contradiction that the cuspidal $\pi(\tau)$ is an irreducible constituent of a parabolically induced representation. $\square$

4.5. Some Global Lifting Results

We now list several additional global lifting results. All test functions are assumed to be matching and have elliptic local components at two fixed places w_1, w_2 which are prime in E .

The stable quasi-packet $\{L(\zeta\nu, \nu^{-1/2}\pi)\}$ is defined in Section 4.3.1, where ζ is a nontrivial quadratic character of C_F , and π is a cuspidal, ζ -invariant, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. Recall that 1_2 denotes the trivial representation of $\mathrm{GL}(2, \mathbb{A}_F)$.

4.34. CLAIM. *Let χ be a character of C_E such that $\chi \neq {}^\sigma\chi$. Let μ be a character of C_F . The following table holds:*

$$(4.1) \quad \begin{array}{|c|c|c|} \hline G & H_1 & H_2 \\ \hline \{L(\varepsilon\nu, \nu^{-1/2}\mu\pi(\chi))\} & \mu 1_2 \otimes_1 \chi & \end{array}$$

PROOF. That there is no contribution from $\mathbf{H}_2$ follows from Lemma 4.24. The rest follows on comparing the lift of the global datum of $\mu 1_2 \otimes_1 \chi$ with the global datum of $\{L(\varepsilon\nu, \nu^{-1/2}\mu\pi(\chi))\}$. $\square$

4.35. CLAIM. *Let χ be a character of C_E such that $\chi \neq {}^\sigma\chi$ but $\chi^2 = {}^\sigma\chi^2$. Then, ${}^\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ for some quadratic character ε' of C_F . Let $\zeta = \varepsilon'$ or $\varepsilon'\varepsilon$. In particular, $\chi^2 = (\chi|_{\mathbb{A}_F^\times} \cdot \zeta) \circ N_{E/F}$. The following table holds:*

$$(4.2) \quad \begin{array}{|c|c|c|} \hline G & H_1 & H_2 \\ \hline \{L(\varepsilon\zeta\nu, \nu^{-1/2}\pi(\chi))\} & & \chi^1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes_2 \chi|_{\mathbb{A}_E^\times} \cdot \zeta \\ \hline \end{array}$$

PROOF. That there is no contribution from $\mathbf{H}_1$ follows from Lemma 4.31. The rest follows from the comparison of global data. $\square$

In what follows, the terms which appear in the ε -trace identity often have nontrivial coefficients (as summarized in Section 3.1.5). So, instead of using tables to describe the global lifting, we record explicitly the ε -trace identities. There is often possible cancellation among the coefficients in these trace identities, but we retain them in order to emphasize how they arise from the trace formula.

Note that if the representations involved do not have elliptic components at w_1, w_2 , then the ε -trace identity is in the trivial form $0 = 0$ due to the restriction on the test functions.

In each of the cases examined, we let S be a finite set of places containing all the bad places for the representations involved. That is, S is a finite set containing the union of $V - V^{\text{un}}(E/F)$ and $\{w_1, w_2\}$ such that the local components of the representations are unramified at all $v \notin S$. The components at $v \notin S$ of the test functions are assumed to be spherical. For any adèlic object (automorphic representation, test function, trace, ..., etc), subscript S denotes the tensor product of local components over the places in S .

First, we consider the cases where the packet π_1 of $\mathbf{H}_1(\mathbb{A}_F)$ has the form $\pi(\theta) \otimes_1 \chi$, where θ, χ are characters of C_E . The following claims follow directly from the results in Section 3.1.5 and Claim 4.28, and we skip their proofs.

4.5.0.3. *Monomial Representations of $\mathbf{H}_1(\mathbb{A}_F)$.* For a character θ of C_E , let $\bar{\theta}$ denote ${}^\sigma\theta$. For the S -part $\{\Pi\}_S$ of a (quasi-)packet $\{\Pi\}$ of $\mathbf{G}(\mathbb{A}_F)$, put $\langle \{\Pi\}_S, f \rangle := \sum_{\Pi' \in \{\Pi\}_S} m(\Pi') \langle \Pi', f \rangle_\varepsilon$, where $m(\Pi')$ is the multiplicity of Π' in the discrete spectrum of $\mathbf{G}$.

Let θ, χ be characters of C_E .

- (1) Suppose $\theta \neq \bar{\theta}$, $\chi \neq \bar{\chi}$, and $\frac{\bar{\theta}}{\theta} \neq \frac{\bar{\chi}}{\chi}, \frac{\chi}{\bar{\chi}}$. Let $\{\Pi\}$ be the packet $[\pi(\theta\chi), \pi(\theta\bar{\chi})]$ of $\mathbf{G}(\mathbb{A}_F)$. It lifts to the induced representation $I_{(2,2)}(\pi(\theta\chi), \pi(\theta\bar{\chi}))$ of $\text{GL}(4, \mathbb{A}_F)$. Those members of $\{\Pi\}$ which are unramified at places outside of S form $\{\Pi\}_S$, which is equal to the tensor product

$$(\otimes_{v \in S} \{\Pi_v^+, \Pi_v^-\}) \otimes (\otimes_{v \notin S} \Pi_v^+).$$

The multiplicity in the discrete spectrum of $\mathbf{G}$ of each $\Pi' \in \{\Pi\}_S$ is given by $m(\Pi') = \frac{1}{2} (1 + (-1)^{n(\Pi')})$, where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$. The following identity holds for matching functions f in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and f_1 in $E(w_1, w_2, \mathbf{H}_1(\mathbb{A}_F), \omega)$ whose components at all $v \notin S$ are spherical:

$$\begin{aligned} (4.3) \quad & \sum_{\Pi' \in \{\Pi\}_S} m(\Pi') \langle \Pi', f \rangle_{\varepsilon, S} \\ &= \frac{1}{2} \cdot \frac{1}{2} \left[\langle \pi(\theta) \otimes_1 \chi, f_1 \rangle_S + \langle \pi(\theta) \otimes_1 \bar{\chi}, f_1 \rangle_S \right. \\ & \quad \left. + \langle \pi(\chi) \otimes_1 \theta, f_1 \rangle_S + \langle \pi(\chi) \otimes_1 \bar{\theta}, f_1 \rangle_S \right] \\ &= \frac{1}{2} \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle_S + \frac{1}{2} \langle \pi(\chi) \otimes_1 \theta, f_1 \rangle_S. \end{aligned}$$

- (2) Suppose $\theta \neq \bar{\theta}$, $\chi \neq \bar{\chi}$, but $\theta\chi = \mu \circ \text{N}_{E/F}$ for some character μ of C_F . Then, $\pi(\theta), \pi(\chi)$ are cuspidal, and

$$\pi(\theta) \otimes_1 \chi = \pi(\chi) \otimes_1 \bar{\theta}, \quad \pi(\chi) \otimes_1 \theta = \pi(\theta) \otimes_1 \bar{\chi}.$$

- (a) Suppose $\chi^2 \neq \bar{\chi}^2$. Then, $\pi(\bar{\chi}/\chi)$ is cuspidal, and the induced representation $\pi(\bar{\chi}/\chi) \rtimes \mu$ is irreducible. The following holds:

$$\begin{aligned}
 (4.4) \quad \frac{1}{2} \left\langle \pi \left(\frac{\bar{\chi}}{\chi} \right) \rtimes \mu, f \right\rangle_{\varepsilon, S} &= \frac{1}{2} \left[\frac{1}{2} \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle_S + \frac{1}{2} \langle \pi(\theta) \otimes_1 \bar{\chi}, f_1 \rangle_S \right] \\
 &= \frac{1}{2} \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle_S.
 \end{aligned}$$

At a place v of F , prime in E , where $\theta_v \neq \bar{\theta}_v$, the representation $\pi(\theta_v)$ is cuspidal. Consequently, the ε_v -twisted Harish-Chandra character of the induced representation $\pi(\bar{\chi}_v/\chi_v) \rtimes \mu_v$ is not identically zero on the elliptic regular set in G_v .

- (b) Suppose $\chi^2 = \bar{\chi}^2$. Then, $\bar{\chi}/\chi = \varepsilon' \circ N_{E/F}$ for some quadratic character ε' of C_F . Since by assumption $\chi \neq \bar{\chi}$, the character ε' is nontrivial and different from ε . Let E' be the quadratic extension of F associated with ε' . The condition $\theta\chi = \mu \circ N_{E/F}$ implies that $\bar{\theta}/\theta = \varepsilon' \circ N_{E/F}$, from which it follows that the E -monomial $\pi(\theta)$ is also E' -monomial ([LL]). Then, $\pi(\theta) \otimes_1 \bar{\chi}$ is equal to $\varepsilon'\pi(\theta) \otimes_1 \chi = \pi(\theta) \otimes_1 \chi$. Hence, if $\pi(\theta) \otimes_1 \chi$ contributes to an ε -trace identity, it is the only contribution from $\mathbf{H}_1$. Since $\pi(\bar{\chi}/\chi)$ is equivalent to the induced representation $I(\varepsilon', \varepsilon'\varepsilon)$, we have:

$$(4.5) \quad \frac{1}{4} \langle \varepsilon' \times \varepsilon'\varepsilon \rtimes \mu, f \rangle_{\varepsilon, S} = \frac{1}{2} \cdot \frac{1}{2} \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle_S.$$

In particular, at a place v of F , prime in E , where $\theta_v \neq \bar{\theta}_v$, the ε_v -twisted Harish-Chandra character of $\varepsilon' \times \varepsilon'\varepsilon \rtimes \mu$ is not identically zero on the elliptic regular set in G_v .

- (3) Suppose $\theta = \mu \circ N_{E/F}$ for some character μ of C_F , and $\chi \neq \bar{\chi}$. Since

$$\pi(\theta) \otimes_1 \chi = \mu\pi(1) \otimes_1 \chi = \pi(1) \otimes_1 (\mu \circ N_{E/F}) \cdot \chi,$$

we assume without loss of generality that $\mu = 1$. The E -monomial representation $\pi(1)$ is equivalent to $I(1, \varepsilon)$. From Section 3.1.3.3, the representations $I(1, \varepsilon) \otimes_1 \chi$ and $I(1, \varepsilon) \otimes_1 \bar{\chi}$ do not contribute to the stable discrete part of the spectral expansion for $\mathbf{H}_1(\mathbb{A}_F)$. The (normalizedly) induced representation $1 \rtimes \pi(\chi)$ of $\mathbf{G}(\mathbb{A}_F)$ is reducible, and its irreducible constituents form the global packet $\{\Pi\} = \otimes_{v \in V} \{\Pi_v^+, \Pi_v^-\}$ (see Section 3.1.2.3). The intertwining operator which appears in the term associated with $1 \rtimes \pi(\chi)$ in the spectral expansion for $\mathbf{H}_1(\mathbb{A}_F)$ acts trivially on Π_v^+ and through the scalar multiplication by -1 on Π_v^- . The following holds:

$$(4.6) \quad \frac{1}{4} \prod_{v \in S} (\langle \Pi_v^+, f_v \rangle_{\varepsilon} - \langle \Pi_v^-, f_v \rangle_{\varepsilon}) = \frac{1}{2} \cdot \frac{1}{2} \langle \pi(\chi) \otimes_1 1, f_1 \rangle_S.$$

- (4) Suppose $\theta = 1$, and $\chi = \mu \circ N_{E/F}$ for some character μ of C_F . Then,

$$\pi(\theta) \otimes_1 \chi = \pi(\chi) \otimes_1 \theta = I(\mu, \mu\varepsilon) \otimes_1 1,$$

which does not contribute to the stable discrete part of the spectral expansion for $\mathbf{H}_1(\mathbb{A}_F)$. If we set up an ε -trace identity using the global datum parametrizing

$\pi(\theta) \otimes_1 \chi$, then there is a nontrivial contribution from $\mathbf{H}_2$, namely, the discretely occurring representation

$$B_{E/F} \pi(\chi) \otimes_2 \mu^2 \varepsilon = (\mu \circ N_{E/F}) I(1, 1) \otimes_2 \mu^2 \varepsilon.$$

The following holds for matching functions:

$$(4.7) \quad \frac{1}{8} \langle 1 \times \varepsilon \rtimes \mu, f \rangle_{\varepsilon, S} = \frac{1}{2} \cdot \frac{1}{4} \langle (\mu \circ N_{E/F}) I(1, 1) \otimes_2 \mu^2 \varepsilon, f_2 \rangle_S.$$

The identity (4.7) cannot be proven using the global techniques employed thus far, for the right hand side comes from an induced representation, and our test functions are required to have two elliptic components. Rather, (4.7) follows from the local character identities deduced in Section 5.1.

4.5.0.4. A Special Case of Non- E -Monomial Representations. Let E' be a quadratic extension of F different from E . Let EE' be the compositum of E and E' . It is a bi-quadratic extension of F . Let σ, σ' be the generators of $\text{Gal}(EE'/F)$, such that E is the fixed field of σ' , E' is the fixed field of σ , and $\text{Gal}(E/F) = \langle \sigma \rangle$, $\text{Gal}(E'/F) = \langle \sigma' \rangle$. Let ε' be the quadratic character of C_F associated with E'/F via class field theory. Let θ be a character of $C_{E'}$ such that $\theta \neq \sigma' \theta$. Let $\pi_{E'}(\theta)$ be the E' -monomial representation of $\text{GL}(2, \mathbb{A}_F)$ associated with θ . Let χ be a character of C_E such that $\chi \neq \sigma \chi$ and $\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$.

Let μ be a character of $C_{E'}$ which satisfies $\mu \circ N_{EE'/E'} = \chi \circ N_{EE'/E}$. Then, $\sigma' \mu / \mu = \varepsilon \circ N_{E'/F}$. Suppose $\mu \theta \neq \sigma'(\mu \theta)$. This implies in particular that $\sigma' \theta / \theta \neq \varepsilon \circ N_{E'/F}$. Let $\pi_{E'}(\mu \theta)$ be the cuspidal E' -monomial representation of $\text{GL}(2, \mathbb{A}_F)$ associated with $\mu \theta$.

Let $\{\Pi\}$ be the global packet of $\mathbf{G}(\mathbb{A}_F)$ which consists of the irreducible constituents of $\varepsilon' \varepsilon \rtimes \pi_{E'}(\mu \theta)$. For any place v , let Π_v^+, Π_v^- denote the two constituents of Π_v if it is reducible (This occurs if and only if $\varepsilon'_v \varepsilon_v = 1$ and $\pi_{E'}(\mu \theta)_v$ is square integrable). If Π_v is irreducible, put $\Pi_v^+ := \Pi_v$ and $\Pi_v^- := 0$. Then, $\{\Pi\} = \otimes_v \{\Pi_v^+, \Pi_v^-\}$.

Let $\{\pi_{E'}(\theta) \otimes_1 \chi\}$ denote the multi-packet of $\mathbf{H}_1(\mathbb{A}_F)$ containing $\pi_{E'}(\theta) \otimes_1 \chi$.

4.36. PROPOSITION. *The following table holds:*

$$(4.8) \quad \begin{array}{|c|c|c|} \hline G & H_1 & H_2 \\ \hline \{\Pi\} & \{\pi_{E'}(\theta) \otimes_1 \chi\} & \\ \hline \end{array}$$

PROOF. We first show that $\pi_{E'}(\theta) \otimes_1 \chi$ contributes to the ε -trace identity by comparing F-H classes. Then, we show that no automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the trace identity.

We consider the all but finitely many odd finite places v of F such that: (i) the place v is unramified in both E and E' ; (ii) the characters θ_v, μ_v, χ_v are unramified. For any such place v , the local packet $\{\Pi_v\}$ is a singleton consisting of a fully induced unramified representation, which we denote by Π_v^+ .

Since any p -adic local field of odd residual characteristic has only one unramified quadratic extension, there are four cases to consider:

(1) $\varepsilon_v \neq 1$.

(a) $\varepsilon'_v = \varepsilon_v$, i.e. $E_v = E'_v$, and $(EE')_v = E_v \oplus E_v$.

We have $\Pi_v^+ = 1 \rtimes \pi_E(\mu_v \theta_v)$. Since θ_v factors through $N_{E'/F}$ via some character θ'_v of $F_v^\times$, the local packet $\pi(\theta_v) \otimes_1 \chi_v$ is equal to $I(\theta'_v, \theta'_v \varepsilon_v) \otimes_1 \chi_v$, which lifts to $1 \rtimes \theta'_v \pi(\chi_v) = 1 \rtimes \pi(\theta_v \chi_v)$. The condition $\chi \circ N_{EE'/E} = \mu \circ N_{EE'/E'}$ implies that, for any $(z_1, z_2) \in (EE')_v^\times$, $\chi_v(z_1 z_2)$ is equal to $\mu_v(z_1 z_2)$. Hence, $\chi_v = \mu_v$, and $\pi_{1,v}$ lifts to Π_v^+ .

- (b) $\varepsilon'_v = 1$, i.e. $E'_v = F_v \oplus F_v$, and $(EE')_v = E_v \oplus E_v$.

In this case, v splits into two places v_1, v_2 of E' . For $i = 1, 2$, let μ_i, θ_i denote μ_{v_i}, θ_{v_i} , respectively. Then, $\Pi_v^+ = \varepsilon_v \rtimes I(\mu_1\theta_1, \mu_2\theta_2)$.

Let χ'_v be an unramified character of $F_v^\times$ such that $\chi_v = \chi'_v \circ N_{E/F}$. Then, $\pi_{1,v} = I(\theta_1, \theta_2) \otimes_1 \chi_v$ lifts to $\varepsilon_v \rtimes I(\chi'_v\theta_1, \varepsilon_v\chi'_v\theta_2)$.

The condition $\chi \circ N_{EE'/E} = \mu \circ N_{EE'/E'}$ implies that, for all (z_1, z_2) in $(EE')_v^\times$, we have the equality

$$\chi'_v(N_{E/F}z_1)\chi'_v(N_{E/F}z_2) = \mu_1(N_{E/F}z_1)\mu_2(N_{E/F}z_2);$$

hence, $\mu_i \circ N_{E/F} = \chi'_v \circ N_{E/F}$ for $i = 1, 2$. The condition $\sigma'\mu/\mu = \varepsilon \circ N_{E'/F}$ implies that $\mu_1/\mu_2 = \varepsilon_v$. Consequently, the set $\{\mu_1, \mu_2\}$ is equal to $\{\chi'_v, \varepsilon_v\chi'_v\}$. Thus, $\pi_{1,v}$ lifts to $\varepsilon_v \rtimes I(\mu_1\theta_1, \mu_2\theta_2)$ or $\varepsilon_v \rtimes I(\mu_2\theta_1, \mu_1\theta_2)$. The two representations are equivalent because $\mu_2 = \varepsilon_v\mu_1$, and the representation $\varepsilon_v \rtimes I(\chi'_v\theta_1, \varepsilon_v\chi'_v\theta_2)$ is equivalent to

$$\varepsilon_v^{-1} \rtimes \varepsilon_v I(\chi'_v\theta_1, \varepsilon_v\chi'_v\theta_2) = \varepsilon_v \rtimes I(\varepsilon_v\chi'_v\theta_1, \chi'_v\theta_2).$$

Thus, $\pi_{1,v}$ lifts to Π_v^+ .

- (2) $\varepsilon_v = 1$, i.e. $E_v = F_v \oplus F_v$.

- (a) $\varepsilon'_v \neq 1$, i.e. $[E'_v, F_v] = 2$, and $(EE')_v = E'_v \oplus E'_v$.

We have $\Pi_v^+ = \varepsilon_v \rtimes I(\mu'_v\theta'_v, \mu'_v\theta'_v\varepsilon'_v)$, where μ'_v, θ'_v are unramified characters of $F_v^\times$ such that $\mu_v = \mu'_v \circ N_{E'/F}$ and $\theta_v = \theta'_v \circ N_{E'/F}$.

By assumption, v splits into two places v_1, v_2 of E_v , and $\pi_{1,v}$ is equal to $I(\theta'_v, \theta'_v\varepsilon'_v) \otimes_1 (\chi_1 \otimes \chi_2)$, where χ_i ($i = 1, 2$) denotes χ_{v_i} . The unramified representation $\pi_{1,v}$ lifts to $\varepsilon'_v \rtimes \theta'_v I(\chi_1, \chi_2)$ of G_v .

The condition $\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ implies that

$$\frac{\chi_1}{\chi_2}(x) \frac{\chi_2}{\chi_1}(y) = \varepsilon'_v(xy), \quad \forall (x, y) \in E_v^\times = F_v^\times \oplus F_v^\times;$$

hence, $\chi_1/\chi_2 = \varepsilon'_v$.

By the condition $\mu \circ N_{EE'/E} = \chi \circ N_{EE'/E}$, we have

$$\chi_1(N_{E'/F}z_1)\chi_2(N_{E'/F}z_2) = \mu_v(z_1z_2),$$

$$\forall (z_1, z_2) \in (EE')_v^\times = E_v'^\times \oplus E_v'^\times.$$

Consequently, $\chi_1 \circ N_{E'/F} = \chi_2 \circ N_{E'/F} = \mu_v = \mu'_v \circ N_{E'/F}$, and $\{\chi_1, \chi_2\} = \{\mu'_v, \varepsilon'_v\mu'_v\}$. It follows that $\pi_{1,v}$ lifts to Π_v^+ .

- (b) $\varepsilon'_v = 1$, i.e. $E'_v = E_v = F_v \oplus F_v$, and $(EE')_v = F_v \oplus F_v \oplus F_v \oplus F_v$.

Suppose v splits into v_1, v_2 in E and into v'_1, v'_2 in E' . For $i = 1, 2$, let μ_i, θ_i, χ_i denote $\mu_{v_i}, \theta_{v_i}, \chi_{v_i}$, respectively. Then, Π_v^+ is equal to $1 \rtimes I(\mu_1\theta_1, \mu_2\theta_2)$. The representation $\pi_{1,v} = I(\theta_1, \theta_2) \otimes_1 (\chi_1 \otimes \chi_2)$ lifts to $\chi_1/\chi_2 \rtimes \chi_2 I(\theta_1, \theta_2)$. The condition $\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ implies that $\chi_1 = \chi_2$. Likewise, $\mu_1 = \mu_2$.

The condition $\mu \circ N_{EE'/E} = \chi \circ N_{EE'/E}$ implies that

$$\mu_1(xy)\mu_2(zx) = \chi_1(xz)\chi_2(yt), \quad \forall (x, y, z, t) \in (EE')_v^\times;$$

hence, $\mu_1 = \mu_2 = \chi_1 = \chi_2$. We conclude that $\pi_{1,v}$ lifts to Π_v^+ .

The packet $\{\Pi\}$ lifts to $I(\pi_{E'}(\mu\theta), \varepsilon\pi_{E'}(\mu\theta))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. By the same argument used in the proof of Claim 4.28, we conclude that if an automorphic representation $\pi_2 = \pi \otimes_2 \eta$ of $\mathbf{H}_2(\mathbb{A}_F)$ contributes to the ε -trace identity, then $\pi = {}^\sigma\pi$. Consequently, $\pi = B_{E/F}\tau$ for some automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_F)$, and η is equal to either ω_τ or $\varepsilon\omega_\tau$.

In the first case, we know from Proposition 4.17 that $\tau \otimes_1 1$ must contribute to the trace identity, which is a contradiction by Theorem 4.10. The second case is ruled out by Lemma 4.29. The proposition follows. $\square$

4.37. CLAIM. *The following holds for matching functions:*

$$(4.9) \quad \frac{1}{2} \prod_{v \in S} (\langle \Pi_v^+, f_v \rangle - \langle \Pi_v^-, f_v \rangle_\varepsilon) = \frac{1}{2} \langle \pi_{E'}(\theta) \otimes_1 \chi, f_1 \rangle_S.$$

PROOF. This follows from Proposition 4.36 and the equation (3.7). Since ${}^\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$, the packet $\pi_{E'}(\theta) \otimes_1 {}^\sigma \chi$ of $\mathbf{H}_1(\mathbb{A}_F)$ is equal to $\varepsilon' \pi_{E'}(\theta) \otimes_1 \chi = \pi_{E'}(\theta) \otimes_1 \chi$. Consequently, $\pi_{E'}(\theta) \otimes_1 \chi$ is the only contribution from $\mathbf{H}_1$ to the trace identity. Since by assumption $\theta \neq {}^{\sigma'} \theta$, the cuspidal E' -monomial representation $\pi_{E'}(\mu\theta)$ is not E -monomial, whence the coefficient $1/2$. $\square$

4.5.0.5. *Induced Automorphic Representations of $\mathrm{GSp}(2, \mathbb{A}_F)$.* Let τ be a cuspidal non- E -monomial, or one dimensional, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. Let Π be the (normalizedly) induced representation $\varepsilon \rtimes \tau$ of $\mathbf{G}(\mathbb{A}_F)$. For any place v of F , let Π_v^+, Π_v^- denote the two constituents of Π_v if it is reducible. If Π_v is irreducible, put $\Pi_v^+ := \Pi_v$ and $\Pi_v^- := 0$. From Section 3.1.2.3, the intertwining operator $M_{P_\beta}(s_{2\alpha+\beta}, 0)_v$ acts trivially on Π_v^+ and as the scalar multiplication by -1 on Π_v^- .

4.38. CLAIM. *The following twisted trace identity holds for matching functions f in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and f_2 in $E(w_1, w_2, \mathbf{H}_2(\mathbb{A}_F), \omega)$:*

$$(4.10) \quad \frac{1}{2} \prod_{v \in S} (\langle \Pi_v^+, f_v \rangle_\varepsilon - \langle \Pi_v^-, f_v \rangle_\varepsilon) = \frac{1}{2} \langle B_{E/F} \tau \otimes_1 \omega_\tau \varepsilon, f_2 \rangle_S.$$

PROOF. This follows from Lemma 4.29 and a comparison of F-H classes. $\square$

The following global character identities follow from local identities which we shall prove in Section 5.1. We record them here for the completeness of the global picture. These global identities hold for all matching test functions, being products of local identities.

For matching functions,

- (1) $\langle \varepsilon' \varepsilon \rtimes \pi(\chi), f \rangle_\varepsilon = \left\langle \chi I(1, 1) \otimes_2 \chi|_{\mathbb{A}_F^\times} \cdot \varepsilon', f_2 \right\rangle,$
where χ is a character of C_E such that ${}^\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$ for some nontrivial quadratic character ε' of C_F ;
- (2) $\langle \varepsilon \times \varepsilon \rtimes \mu, f \rangle_\varepsilon = \langle \mu I(1, 1) \otimes_1 1, f_1 \rangle = \langle (\mu \circ N_{E/F}) I(1, 1) \otimes_2 \mu^2, f_2 \rangle,$
where ε, μ are charaters of C_F ;
- (3) $\langle \varepsilon \rtimes \pi(\chi), f \rangle_\varepsilon = \langle I(1, 1) \otimes_1 \chi, f_1 \rangle,$
where χ is a character of C_E .

4.6. Final Words

Let σ be the generator of $\mathrm{Gal}(E/F)$. The Galois group $\mathrm{Gal}(E/F)$ acts on the representations τ of $\mathrm{GL}(2, \mathbb{A}_E)$ via ${}^\sigma \tau(g) = \tau(\sigma g)$ for all $g \in \mathrm{GL}(2, \mathbb{A}_E)$.

For an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A}_E)$, let $\pi(\tau)$ denote the automorphic representation of $\mathrm{GL}(4, \mathbb{A}_F)$ which is obtained from τ via the twisted endoscopic lifting from $\mathrm{R}_{E/F} \mathrm{GL}(2)$ to $\mathrm{GL}(4)$.

For an automorphic representation π of $\mathrm{GL}(2, \mathbb{A}_F)$, let $B_{E/F} \pi$ denote the automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$ obtained from π via base change ([L], [F1]).

For an automorphic representation/packet π_i of $\mathbf{H}_i(\mathbb{A}_F)$ ($i = 1, 2$), let $\xi_i^*(\pi_i)$ denote the (quasi-)packet of $\mathrm{GSp}(2, \mathbb{A}_F)$ which is the lift of π_i .

We say that a global (quasi-)packet is **discrete spectrum** if it contains a discrete spectrum automorphic representation. A discrete spectrum (quasi-)packet of the group $\mathrm{GSp}(2, \mathbb{A}_F)$ is said to be ε -**invariant** if it contains a *discrete spectrum*, ε -invariant, automorphic representation.

We now give the list of (quasi-)packets which comprise the image of the ε -endoscopic lifting in the discrete spectrum of $\mathrm{GSp}(2)$. These packets are necessarily ε -invariant.

(1) **Unstable, cuspidal:**

- (a) $[\pi, \varepsilon\pi] = \xi_1^*(\pi \otimes 1) = \xi_2^*(B_{E/F}\pi \otimes \omega_\pi)$,
where π is a cuspidal, non- E -monomial, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$. The packet lifts to the induced representation $I_{(2,2)}(\pi, \varepsilon\pi)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.
- (b) $[\pi(\theta\chi), \pi(\theta^\sigma\chi)] = \xi_1^*(\pi(\theta) \otimes 1)$,
where θ, χ are characters of C_E such that none of $\theta, \chi, \theta\chi, \theta^\sigma\chi$ is invariant under the action of σ . The packet lifts to the induced representation $I_{(2,2)}(\pi(\theta\chi), \pi(\theta^\sigma\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

(2) **Unstable, residual:**

- $[\mu 1_2, \varepsilon\mu 1_2] = \xi_1^*(\mu 1_2 \otimes 1) = \xi_2^*((\mu \circ N_{E/F}) 1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes \mu^2)$,
where μ is a character of C_F , and 1_2 denotes the trivial representation of $\mathrm{GL}(2, \mathbb{A}_F)$. The packet lifts to $I_{(2,2)}(\mu 1_2, \varepsilon\mu 1_2)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

(3) **Stable, cuspidal:**

- (a) $\xi_1^*(\tau \otimes 1)$,
where τ is a cuspidal, non- E -monomial, automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$, χ a character of C_E , such that there is no quadratic character (trivial or otherwise) ε' of C_F for which $\sigma\chi/\chi = \varepsilon' \circ N_{E/F}$ and τ is equivalent to $\varepsilon'\tau$ or $\varepsilon'\varepsilon\tau$.
The packet lifts to the ε -invariant, cuspidal, automorphic representation $\pi(\chi B_{E/F}\tau)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.
- (b) $\xi_2^*(\tau_E \otimes 2)$,
where τ_E is a cuspidal automorphic representation of $\mathrm{GL}(2, \mathbb{A}_E)$, μ a character of C_F , such that $\tau_E \neq \sigma\tau_E$ and $\omega_{\tau_E} = \mu \circ N_{E/F}$.
The packet lifts to the ε -invariant, cuspidal, automorphic representation $\pi(\tau_E)$ of $\mathrm{GL}(4, \mathbb{A}_F)$.

(4) **Stable, residual:**

- (a) $\{L(\varepsilon\nu, \nu^{-1/2}\pi(\chi))\} = \xi_1^*(1_2 \otimes 1)$,
where χ is a character of C_E not fixed by $\mathrm{Gal}(E/F)$. The quasi-packet lifts to $J(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$, the Langlands quotient of the induced representation $I_{(2,2)}(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$.
- (b) $\{L(\varepsilon\zeta\nu, \nu^{-1/2}\pi(\chi))\} = \xi_2^*(\chi 1_{\mathrm{GL}(2, \mathbb{A}_E)} \otimes 2 \chi|_{\mathbb{A}_F^\times} \cdot \zeta)$,
where $\zeta \neq \varepsilon$ is a nontrivial quadratic character of C_F , χ a character of C_E , such that $\sigma\chi/\chi = \zeta \circ N_{E/F}$. The quasi-packet lifts to the Langlands quotient $J(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. Note that the packet $\{L(\zeta\nu, \nu^{-1/2}\pi(\chi))\}$ is disjoint from $\{L(\varepsilon\zeta\nu, \nu^{-1/2}\pi(\chi))\}$, but they both lift to $J(\nu^{1/2}\pi(\chi), \nu^{-1/2}\pi(\chi))$.

4.6.0.6. *Surjectivity of the Twisted Endoscopic Lifting.* Recall the definition of the operator $\rho_\omega(\varepsilon)$, or $\rho(\varepsilon)$, on $L(\mathbf{G}(\mathbb{A}_F), \omega)$:

$$\rho_\omega(\varepsilon) : \phi \mapsto \varepsilon\phi, \quad \forall \phi \in L(\mathbf{G}(\mathbb{A}_F), \omega).$$

4.39. LEMMA. *If an irreducible, discrete spectrum, automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$, with central character ω , is ε -invariant, then the distribution $f \mapsto \text{tr } \Pi(f)\rho_\omega(\varepsilon)$ on $C(\mathbf{G}(\mathbb{A}_F), \omega)$ is nonzero.*

PROOF. Let $\tilde{G} = \mathbf{G}(\mathbb{A}_F) \times \{\pm 1\} \rtimes \langle \varepsilon \rangle$, where $\varepsilon^2 = 1$, and $(1, 1, \varepsilon)(g, 1, 1) = (g, \varepsilon(g), \varepsilon)$.

By assumption, Π is a subrepresentation of the right-regular representation ρ_ω on $L(\mathbf{G}(\mathbb{A}_F), \omega)$. Hence, $\rho_\omega(\varepsilon)\Pi$ is a subrepresentation and consequently belongs to the discrete spectrum. By the assumption that the multiplicity one theorem holds for $\mathbf{G}$, the operator $\rho_\omega(\varepsilon)$ maps the space of Π to itself, with

$$(4.1) \quad \rho_\omega(\varepsilon)^{-1}\Pi(g)\rho_\omega(\varepsilon) = \varepsilon(g)\Pi(g), \quad \forall g \in \mathbf{G}(\mathbb{A}_F).$$

We extend Π to a representation Π_1 of $\tilde{G}$, on the space of Π , as follows:

$$\Pi_1(g, 1, 1) := \Pi(g), \quad \Pi_1(1, \pm 1, 1) := \pm 1, \quad \Pi_1(1, 1, \varepsilon) := \rho_\omega(\varepsilon).$$

For all $g \in \mathbf{G}(\mathbb{A}_F)$, we have:

$$\Pi_1(\underbrace{(1, 1, \varepsilon)(g, 1, 1)}_{=(g, \varepsilon(g), \varepsilon)}) = \rho_\omega(\varepsilon)\pi(g), \quad \Pi_1((g, \varepsilon(g), \varepsilon)) = \varepsilon(g)\pi(g)\rho_\omega(\varepsilon).$$

It follows from (4.1) that Π_1 is indeed a representation of $\tilde{G}$.

We extend Π also to a representation Π_2 of $\tilde{G}$ such that Π_2 is the same as Π_1 on the subgroup $\mathbf{G}(\mathbb{A}_F) \times \{\pm 1\}$ of $\tilde{G}$, but $\Pi_2(1, 1, \varepsilon) := -\rho_\omega(\varepsilon)$. Since Π is irreducible, it follows from Schur's lemma that Π_1 and Π_2 are inequivalent.

The center of $\tilde{G}$ is $\tilde{Z} = \{(z, 1, \varepsilon) : z \in \mathbf{Z}(\mathbb{A}_F)\}$, where $\mathbf{Z}$ is the center of $\mathbf{G}$. For $i = 1, 2$, the character of Π_i is $\langle \Pi_i, \tilde{f} \rangle = \text{tr} \int_{\tilde{Z} \backslash \tilde{G}} \Pi_i(\tilde{g}) \tilde{f}(\tilde{g}) d\tilde{g}$, which is equal to

$$\text{tr} \int_{\{\tilde{g}=(g, \pm 1, 1)\}} \pm \Pi(g) \tilde{f}(\tilde{g}) d\tilde{g} + \text{tr} \int_{\{\tilde{g}=(g, \pm 1, \varepsilon)\}} \pm \Pi(g) A_i \tilde{f}(\tilde{g}) d\tilde{g},$$

where $A_1 = \rho_\omega(\varepsilon)$ and $A_2 = -\rho_\omega(\varepsilon)$. Hence, the difference $\langle \Pi_1, \tilde{f} \rangle - \langle \Pi_2, \tilde{f} \rangle$ is equal to

$$\begin{aligned} & 2 \text{tr} \int_{\{\tilde{g}=(g, \pm 1, \varepsilon)\}} \pm \Pi(g) \rho_\omega(\varepsilon) \tilde{f}(\tilde{g}) d\tilde{g} \\ &= 2 \text{tr} \int_{\{(g, 1, \varepsilon)\}} \Pi(g) \rho_\omega(\varepsilon) \tilde{f}((g, 1, \varepsilon)) d\tilde{g} - 2 \text{tr} \int_{\{(g, -1, \varepsilon)\}} \Pi(g) \rho_\omega(\varepsilon) \tilde{f}((g, -1, \varepsilon)) d\tilde{g}. \end{aligned}$$

Suppose the twisted character $\langle \Pi, f \rangle_{\rho_\omega(\varepsilon)}$ is identically zero. Then, both terms in the sum above are zero, which implies that $\langle \Pi_1, \tilde{f} \rangle - \langle \Pi_2, \tilde{f} \rangle = 0$ for all test functions $\tilde{f}$ on $\tilde{G}$. Since $\Pi_1 \not\cong \Pi_2$, we have a contradiction by the linear independence of characters. $\square$

4.40. LEMMA. *Let $\{\Pi_1, \Pi_2, \dots, \Pi_n\}$ be a finite set of inequivalent automorphic representations of $\mathbf{G}(\mathbb{A}_F)$ which satisfy the hypothesis of Lemma 4.39. The twisted characters $\{\langle \Pi_i, f \rangle_{\rho_\omega(\varepsilon)}\}_{1 \leq i \leq n}$ are linearly independent.*

PROOF. For each $1 \leq i \leq n$, let $\Pi_{i,1}, \Pi_{i,2}$ be the representations of $\tilde{G}$ defined as in the proof of Lemma 4.39. Suppose there exist complex numbers $c_1, c_2, \dots, c_n$, not all zero, such that

$$\sum_{i=1}^n c_i \langle \Pi_i, f \rangle_{\rho_\omega(\varepsilon)} = 0.$$

Then, by the same argument used in the proof of Lemma 4.39, we have:

$$\sum_{i=1}^n c_i \left(\langle \Pi_{i,1}, \tilde{f} \rangle - \langle \Pi_{i,2}, \tilde{f} \rangle \right) = 0$$

for all test functions $\tilde{f}$ on $\tilde{G}$. Hence, we have a contradiction by the linear independence of the characters $\{\langle \Pi_{i,j}, \tilde{f} \rangle\}_{\substack{1 \leq i \leq n \\ j=1,2}}$. $\square$

4.41. COROLLARY. *Each ε -invariant, discrete spectrum, automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ with two elliptic components belongs to a packet listed in this section.*

PROOF. Suppose a discrete spectrum automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$, with central character ω and two elliptic components, is ε -invariant. If we set up an ε -trace identity using the global datum of Π , by Lemma 4.40 the $\mathbf{G}$ side of the identity is nonzero. Consequently, there must be nonzero contribution from the endoscopic groups side of the trace identity. The image of the lifting of the discrete spectrum automorphic representations of the endoscopic groups consists precisely of those representations which appear in the list, whence the claim. $\square$

CHAPTER 5

The Local Picture

Let k be a p -adic field with ring of integers $\mathcal{O}$.

For an algebraic k -group $\mathbf{H}$, let $H = \mathbf{H}(k)$. Let H^{reg} denote the set of regular elements in H . Let $Z_0(H)$ be the maximal k -split component of the center of H . Let $\bar{H}$ denote the quotient $H/Z_0(H)$. Let $C(H)$ denote the space of smooth, compactly supported modulo $Z_0(H)$ functions on H . For a character ω of $Z_0(H)$, let $C(H, \omega)$ denote the space of functions f in $C(H)$ which satisfy

$$f(zh) = \omega(z)^{-1}f(h), \quad \forall z \in Z_0(H), h \in H.$$

Let $J = \begin{pmatrix} & & & 1 \\ & & 1 & \\ & -1 & & \\ -1 & & & \end{pmatrix} \in \text{GL}(4)$. Let $\mathbf{G}$ be the reductive k -group

$$\text{GSp}(2) = \{g \in \text{GL}(4) : {}^t g J g = \lambda(g) J \text{ for some } \lambda(g) \in \mathbb{G}_m\}.$$

We call $\lambda(g)$ the similitude factor of g . The character $\lambda \in \text{Hom}(\mathbf{G}, \mathbb{G}_m)$ descends to a character in $\text{Hom}(G, k^\times)$. Let $\mathbf{Z} = \{\text{diag}(z, z, z, z) : z \in \mathbb{G}_m\}$ be the center of $\mathbf{G}$. Since $\mathbf{Z}$ is k -split, the group $Z_0(G)$ is equal to $\mathbf{Z}$.

Let ε be a quadratic character of $k^\times$. We let ε denote also the character

$$g \mapsto \varepsilon(\lambda(g))$$

on G . The similitude factor of any $\text{diag}(z, z, z, z) \in \mathbf{Z}$ is z^2 ; hence, ε is trivial on the center of G .

Notation:

- For a representation π of G , let $\varepsilon\pi$ denote the representation of G on the space of π defined by

$$\varepsilon\pi = \varepsilon \otimes \pi : g \mapsto \varepsilon(g)\pi(g), \quad \forall g \in G.$$

- For $g \in \text{GL}(2, k)$, put $\varepsilon(g) := \varepsilon(\det g)$. For a representation τ of $\text{GL}(2, k)$, let $\varepsilon\tau$ denote the $\text{GL}(2, k)$ -module on the space of τ where g in $\text{GL}(2, k)$ acts by $\varepsilon(g)\tau(g)$.

Let G' be either $\text{GSp}(2, k)$ or $\text{GL}(2, k)$. For an admissible ([BZ]) representation (π, V) of G' , where V is the complex vector space on which G' acts via π , put

$$\text{Hom}_{G'}(\pi, \varepsilon\pi) := \{A \in \text{Hom}_{\mathbb{C}}(V, V) : \varepsilon(g)\pi(g)A = A\pi(g), \forall g \in G'\}.$$

We call $A \in \text{Hom}_G(\pi, \varepsilon\pi)$ an **intertwining operator**.

5.1. DEFINITION. We say that an admissible representation π of G' is **ε -invariant** if π is equivalent to $\varepsilon\pi$, i.e. there exists a nontrivial intertwining operator A in $\text{Hom}_{G'}(\pi, \varepsilon\pi)$.

Let (π, V) be an irreducible, admissible, ε -invariant representation of G . Let A be a nontrivial operator in $\text{Hom}_G(\pi, \varepsilon\pi)$. Since ε is quadratic, A^2 intertwines the irreducible π with itself. Hence, by Schur's lemma A^2 is a scalar multiplication on V . Multiplying A by the scalar $(A^2)^{-1/2}$ if necessary, we assume $A^2 = 1$. Then, A is unique up to a sign.

Fix once and for all a Haar measure ([BZ]) dg on $\bar{G} = G/Z$. Let ω_π denote the central character of π . For $f \in C(G, \omega_\pi)$, define the convolution operator $\pi(f)A$ on V as follows:

$$\pi(f)Av := \int_{\bar{G}} f(g)\pi(g)Av \, dg, \quad \forall v \in V.$$

Since f is locally constant, compactly supported modulo Z , and π is admissible, $\pi(f)A$ has finite rank and its trace $\text{tr } \pi(f)A$ is finite. We often drop the ω_π from $C(G, \omega_\pi)$ and write simply $C(G)$. It should be understood implicitly that the test function f transforms under Z via ω_π^{-1} .

Recall from chapter 2 the definitions of the ε -endoscopic groups $\mathbf{H}_1, \mathbf{H}_2$ of $\mathbf{G}$ over k . From local class field theory, there is a quadratic extension $\mathcal{K}$ of k which corresponds to ε . In particular, ε is the unique nontrivial quadratic character of $k^\times$ whose restriction to $N_{\mathcal{K}/k}\mathcal{K}^\times$ is trivial. The group of k -points of $\mathbf{H}_i$ ($i = 1, 2$) is as follows:

- (1) $H_1 = (\text{GL}(2, k) \times \mathcal{K}^\times)' := \{(g, x) \in \text{GL}(2, k) \times \mathcal{K}^\times : \det g = N_{\mathcal{K}/k}x\}$;
- (2) $H_2 = (\text{GL}(2, \mathcal{K}) \times k^\times) / \mathcal{K}^\times$,
where $\mathcal{K}^\times$ embeds into $\text{GL}(2, \mathcal{K}) \times k^\times$ via $x \mapsto (\text{diag}(x, x), N_{\mathcal{K}/k}x^{-1})$.

We have:

$$\begin{aligned} Z_0(H_1) &= \{(\text{diag}(z, z), z) : z \in k^\times\}, \\ Z_0(H_2) &= \{(\text{diag}(1, 1), z)_* : z \in k^\times\}, \end{aligned}$$

where the lower $*$ signifies the image in H_2 of an element in $\text{GL}(2, \mathcal{K}) \times k^\times$.

Let $i = 1, 2$. Fix once and for all a Haar measure dh_i on $\bar{H}_i = H_i/Z_0(H_i)$. Let π_i be an irreducible admissible representation of H_i . For $z \in Z_0(H_i)$, $\pi_i(z)$ acts as a scalar $\omega_{\pi_i}(z)$ on the space of π_i . We call ω_{π_i} the **split central character** of π_i . For $f_i \in C(H_i, \omega_{\pi_i})$, let $\pi(f_i)$ denote the convolution operator

$$\int_{\bar{H}_i} f_i(h)\pi_i(h) \, dh_i.$$

It has finite rank; hence, its trace $\text{tr } \pi_i(f_i)$ is finite. As in the case of G , we often write $C(H_i)$ instead of $C(H_i, \omega_{\pi_i})$. It should be understood that the test function f_i transforms under $Z_0(H_i)$ via $\omega_{\pi_i}^{-1}$.

Notation/Terminology:

- Put $\text{GL}(2, k)^\mathcal{K} := \{g \in \text{GL}(2, k) : \det g \in N_{\mathcal{K}/k}\mathcal{K}^\times\}$.
- For any quadratic extension $\mathcal{L}$ of k , we say that an admissible representation π of $\text{GL}(2, k)$ is **$\mathcal{L}$ -monomial** if it is the monomial representation $\pi(\theta)$ associated with a character θ of $\mathcal{L}^\times$ (see [JL, Thm. 4.6], [K]).
- We say that a representation is **fully induced** if it is an irreducible parabolically induced representation.
- For a representation τ of $\text{GL}(2, k)$ and a character χ of $\mathcal{K}^\times$, let $\tau \otimes_1 \chi$ denote the following representation of H_1 on the space of τ :

$$\tau \otimes_1 \chi : (g, x) \mapsto \chi(x)\tau(g), \quad \forall (g, x) \in H_1.$$

Note that $\tau \otimes_1 \chi$ is reducible of length two if τ is $\mathcal{K}$ -monomial ([LL]).

- For a representation π of $\text{GL}(2, \mathcal{K})$, a character μ of $k^\times$, such that $\omega_\pi = \mu \circ N_{\mathcal{K}/k}$, let $\pi \otimes_2 \mu$ denote the following representation of H_2 on the space of π :

$$\pi \otimes_2 \mu : (g, c)_* \mapsto \mu(c)\pi(g), \quad \forall (g, c)_* \in H_2.$$

The lower $*$ signifies the image of $(g, c) \in \text{GL}(2, \mathcal{K}) \times k^\times$ in H_2 . Note that $\pi \otimes_2 \mu$ is well defined because of the condition on ω_π and μ .

- For an admissible ε -invariant G -module π , a nontrivial intertwining operator $A \in \text{Hom}_G(\pi, \varepsilon\pi)$, and a function f in $C(G)$, put

$$\langle \pi, f \rangle_A := \text{tr } \pi(f)A.$$

It depends implicitly on the fixed Haar measure dg on $\bar{G}$.

- Let $i = 1$ or 2 . For an admissible H_i -module π_i and a function $f_i \in C(H_i)$, put

$$\langle \pi_i, f_i \rangle := \text{tr } \pi_i(f_i).$$

It depends on the fixed Haar measure dh_i on $\bar{H}_i$.

- Let σ be the generator of $\text{Gal}(\mathcal{K}/k)$. For $\gamma \in \mathcal{K}$, put $\bar{\gamma} := \sigma\gamma$. For a character χ of $\mathcal{K}^\times$, put ${}^\sigma\chi(\gamma) := \chi(\bar{\gamma})$ for all $\gamma \in \mathcal{K}^\times$.

For a representation π of $\text{GL}(2, \mathcal{K})$, let ${}^\sigma\pi$ denote the representation

$${}^\sigma\pi((g_{ij})) = \pi((\sigma g_{ij})), \quad \forall (g_{ij}) \in \text{GL}(2, \mathcal{K}).$$

5.1. Induced Representations

5.1.1. The Intertwining Operator. Let $P = MN$ be the upper triangular Heisenberg parabolic subgroup of G with Levi component

$$M = \left\{ \begin{pmatrix} a & & \\ & g_2 & \\ & & \frac{\det g_2}{a} \end{pmatrix} : a \in k^\times, g_2 \in \text{GL}(2, k) \right\}$$

and unipotent component

$$N = \left\{ \begin{pmatrix} 1 & * & * & * \\ & 1 & 0 & * \\ & & 1 & * \\ & & & 1 \end{pmatrix} \right\}.$$

We identify M with $k^\times \times \text{GL}(2, k)$ via $\begin{pmatrix} a & & \\ & g_2 & \\ & & (\det g_2)/a \end{pmatrix} \mapsto (a, g_2)$.

Let μ be a character of $k^\times$. Let χ be a character of $\mathcal{K}^\times$. Let (τ_2, V_2) be the monomial representation $\pi(\chi)$ of $\text{GL}(2, k)$ associated with χ . Let $(\mu \otimes \tau_2, V_2)$ be the representation of P defined by

$$\mu \otimes \tau_2 : (a, g_2)n \mapsto \mu(a)\tau_2(g_2), \quad \forall (a, g_2) \in M, n \in N.$$

Let $(\pi, V) = \mu \rtimes \tau_2 := I_P^G(\mu \otimes \tau_2)$ be the representation of G normalizedly induced ([BZ]) from $\mu \otimes \tau_2$.

Let $\tau = \mu \otimes \tau_2$. For $m \in M$, let $\delta_M(m) = |\det(\text{Ad } m|_{\mathfrak{n}})|$, where $\mathfrak{n}$ denotes the Lie algebra of N . By the definition of normalizedly induced representations, the vector space V consists of the smooth ([BZ]) functions $\varphi : G \rightarrow V_2$ which satisfy

$$\varphi(mng) = (\delta_M^{1/2}\tau)(m)\varphi(g), \quad \forall m \in M, n \in N, g \in G.$$

The group G acts on V via right translation $(\pi(g)\varphi)(h) = \varphi(hg)$.

The similitude factor of $(a, g_2) \in M$ is equal to $\det g_2$; hence, $\varepsilon \otimes (\mu \rtimes \tau_2) = \mu \rtimes \varepsilon\tau_2$. Since $\tau_2 = \pi(\chi)$ is ε -invariant (in the context of $\text{GL}(2, k)$), the induced representation $\pi = \mu \rtimes \tau_2$ is ε -invariant (in the context of $\text{GSp}(2, k)$). In what follows, we define a nontrivial operator A in $\text{Hom}_G(\pi, \varepsilon\pi)$, and then we compute explicitly the twisted character $\langle \pi, f \rangle_A$ for $f \in C(G)$.

Let $\text{GL}(2, k)^\mathcal{K} = \{g \in \text{GL}(2, k) : \det g \in \text{N}_{\mathcal{K}/k}\mathcal{K}^\times\}$. It is a subgroup of index 2 in $\text{GL}(2, k)$. From [LL, Sect. 2], there exist subspaces V_2^+, V_2^- in the space V_2 of τ_2 such that $V_2 = V_2^+ \oplus V_2^-$ as a $\text{GL}(2, k)^\mathcal{K}$ -module. Moreover, V_2^- is equal to $\tau_2(w)V_2^+$ for any element $w \in \text{GL}(2, k) - \text{GL}(2, k)^\mathcal{K}$. Let A_2 be the operator on V_2 defined by

$$A_2(v^+ + v^-) := v^+ - v^-, \quad \forall v^+ \in V_2^+, v^- \in V_2^-.$$

It is an intertwining operator in $\text{Hom}_{\text{GL}(2,k)}(\tau_2, \varepsilon\tau_2)$ whose square is the identity. Note that by the symmetry between V_2^+ and V_2^- , a choice is involved in the definition of A_2 .

Define a map A from V to the set of smooth V_2 -valued functions on G as follows:

$$(5.1) \quad (A\varphi)(g) := \varepsilon(g)A_2(\varphi(g)), \quad \forall g \in G, \varphi \in V.$$

5.2. CLAIM. *The map A is an automorphism of V such that $A^2 = 1$, and*

$$\varepsilon\pi(g)A = A\pi(g), \quad g \in G.$$

In other words, A is a nontrivial intertwining operator in $\text{Hom}_G(\pi, \varepsilon\pi)$.

PROOF. First, we show that A is an endomorphism of V . It then follows trivially that $A^2 = 1$ and hence A is invertible.

For every $\varphi \in V$, we need to show that $(A\varphi)(mng) = (\delta_M^{1/2}\tau)(m)\varphi(g)$ for all $m = (a, g_2) \in M, n \in N, g \in G$. The restriction of ε to N is trivial, for the similitude factor of a unipotent element is 1. We have:

$$\begin{aligned} (A\varphi)(mng) &= \varepsilon(m)\varepsilon(n)\varepsilon(g)A_2(\varphi(mng)) \\ &= \varepsilon(\det g_2)\varepsilon(g)A_2\left(\delta_M^{1/2}(m)\mu(a)\tau_2(g_2)\varphi(g)\right) \\ &= \delta_M^{1/2}(m)\mu(a)\varepsilon(\det g_2)\varepsilon(\det g_2)\tau_2(g_2)\varepsilon(g)A_2(\varphi(g)) \\ &= (\delta_M^{1/2}\tau)(m)\varepsilon(g)A_2(\varphi(g)) \\ &= (\delta_M^{1/2}\tau)(m)(A\varphi)(g). \end{aligned}$$

Hence, $A\varphi$ lies in V .

Next, we show that

$$(5.2) \quad \varepsilon(g)\pi(g)A\varphi = A\pi(g)\varphi, \quad \forall g \in G, \varphi \in V.$$

For all $h \in G$, on the left-hand side of (5.2) we have:

$$(\varepsilon(g)\pi(g)A\varphi)(h) = \varepsilon(g)(A\varphi)(hg) = \varepsilon(g)\varepsilon(hg)A_2(\varphi(hg)) = \varepsilon(h)A_2(\varphi(hg)).$$

On the right-hand side, we have:

$$(A\pi(g)\varphi)(h) = \varepsilon(h)A_2((\pi(g)\varphi)(h)) = \varepsilon(h)A_2(\varphi(hg)).$$

Hence, left-hand side equals right-hand side, which completes the proof. $\square$

5.1.2. A Trace Identity between G and H_1 . Suppose the field $\mathcal{K}$ is equal to $k(\sqrt{B})$ for some $B \in k^\times - k^{\times 2}$. We define an embedding $\phi : \mathcal{K}^\times \hookrightarrow \text{GL}(2, k)$ as follows:

$$\phi : a + b\sqrt{B} \mapsto \begin{pmatrix} a & bB \\ b & a \end{pmatrix}, \quad \forall a, b \in k; (a, b) \neq (0, 0).$$

Note that, for $\gamma \in \mathcal{K}^\times$, the (possibly non-distinct) eigenvalues of $\phi(\gamma)$ are $\gamma, \bar{\gamma}$, and $\det \phi(\gamma) = N_{\mathcal{K}/k}\gamma$. Let

$$T_{\mathcal{K}} = \{\phi(\gamma) : \gamma \in \mathcal{K}^\times\}.$$

For an element $t_2 \in \text{GL}(2, k)$ which is conjugate to $\phi(\gamma)$ for some $\gamma \in \mathcal{K}^\times$, put

$$D_{T_{\mathcal{K}}}(t_2) := \frac{|(\gamma - \bar{\gamma})^2|}{|N_{\mathcal{K}/k}\gamma|}.$$

Recall that $\tau_2 = \pi(\chi)$ is the $\mathcal{K}$ -monomial representation associated with a character χ of $\mathcal{K}^\times$. We have defined in the previous section an intertwining operator A_2 in $\text{Hom}_{\text{GL}(2,k)}(\tau_2, \varepsilon\tau_2)$. Let $\chi_{\tau_2}^\varepsilon$ be the ε -twisted Harish-Chandra character associated with

the distribution $f \mapsto \text{tr } \tau_2(f) A_2$ on $C(\text{GL}(2, k))$. In particular, for $g, h \in \text{GL}(2, k)$, we have $\chi_{\tau_2}^\varepsilon(h^{-1}gh) = \varepsilon(h)\chi_{\tau_2}^\varepsilon(g)$.

Let t_2 be a regular element in $\text{GL}(2, k)$. If $\chi_{\tau_2}^\varepsilon(t_2) \neq 0$, then the centralizer of t_2 must be a torus which lies in the kernel of ε . All such tori are conjugate to $T_{\mathcal{K}}$; hence, $\chi_{\tau_2}^\varepsilon(t_2)$ is zero unless t_2 is conjugate to an element in $T_{\mathcal{K}}$. By [L, Lemma 7.19], for each regular $t_2 = \phi(\gamma) \in T_{\mathcal{K}}$, $\gamma \in \mathcal{K}^\times - k^\times$, we have:

$$(5.3) \quad D_{T_{\mathcal{K}}}^{1/2}(t_2)\chi_{\pi(\chi)}^\varepsilon(t_2) = d \cdot \lambda(\mathcal{K}/k, \psi) \cdot \varepsilon(\gamma - \bar{\gamma}) [\chi(\gamma) + \chi(\bar{\gamma})].$$

Here, $\lambda(\mathcal{K}/k, \psi)$ is a constant independent of t_2 , and d is ± 1 depending on the choice of sign involved in the definition of A_2 . For simplicity, we assume A_2 is chosen such that $d = 1$.

Let $T = \left\{ \begin{pmatrix} a & t_2 \\ & (\det t_2)/a \end{pmatrix} : a \in k^\times, t_2 \in T_{\mathcal{K}} \right\}$. It is an elliptic maximal torus in M . Observe that, for $\gamma \in \mathcal{K}^\times$,

$$\varepsilon \left(\begin{pmatrix} a & \phi(\gamma) \\ & (\det \phi(\gamma))/a \end{pmatrix} \right) = \varepsilon(\det \phi(\gamma)) = \varepsilon(N_{\mathcal{K}/k}\gamma) = 1;$$

hence, T lies in the kernel of ε .

Fix a Haar measure dt on $\bar{T} = T/Z$. Let $d\bar{g} = dg/dt$. For $t \in T$ and $f \in C(G)$, put

$$O_G^\varepsilon(f, t) := \int_{T \backslash G} \varepsilon(\bar{g}) f(\bar{g}^{-1}t\bar{g}) d\bar{g}.$$

Note that $\varepsilon(\bar{g})$ is well defined because $T \subset \ker \varepsilon$.

Recall that we identify (a, g_2) in $k^\times \times \text{GL}(2, k)$ with $\text{blockdiag}(a, g_2, (\det g_2)/a)$ in M .

5.3. LEMMA. *Let μ be a character of $k^\times$. Let χ be a character of $\mathcal{K}^\times$. Let A in $\text{Hom}_G(\mu \rtimes \pi(\chi), \mu \rtimes \varepsilon\pi(\chi))$ be the intertwining operator defined in Section 5.1.1. Then,*

$$(5.4) \quad \langle \mu \rtimes \pi(\chi), f \rangle_A = \frac{1}{4} \int_{t=(a, t_2) \in \bar{T}^{\text{reg}}} \left[\mu(a) + \mu \left(\frac{N_{\mathcal{K}/k}\gamma}{a} \right) \right] [\chi(\gamma) + \chi(\bar{\gamma})] \\ \cdot \lambda(\mathcal{K}/k, \psi) \varepsilon(\gamma - \bar{\gamma}) D_{T \backslash G}^{1/2}(t) O_G^\varepsilon(f, t) dt.$$

Here, $t_2 = \phi(\gamma) \in T_{\mathcal{K}}$, $\gamma \in \mathcal{K}^\times$.

PROOF. We keep the same notation used in the previous section, i.e. $\tau_2 = \pi(\chi)$, $(\pi, V) = \mu \rtimes \pi(\chi)$, $\tau = \mu \otimes \tau_2, \dots$, etc. Let K be the hyperspecial maximal compact subgroup $\text{GSp}(2, \mathcal{O})$ of G . We have the Iwasawa decomposition $G = PK = NMK$; hence, $\bar{G} = N\bar{M}K$, where $\bar{M} = M/Z$. We also have the corresponding measure decomposition $dg = \delta_M^{-1}(m) dndmdk$, where dm is a measure on $\bar{M}$. Recall that $\varepsilon|_N = 1$. For

$h \in G, \varphi \in V,$

$$\begin{aligned}
(\pi(f)A\varphi)(h) &= \int_{\bar{G}} f(g)(\pi(g)(A\varphi))(h) dg = \int_{\bar{G}} f(g)(A\varphi)(hg) dg \\
&= \int_G f(h^{-1}g)(A\varphi)(g) dg = \int_G f(h^{-1}g)\varepsilon(g)A_2(\varphi(g)) dg \\
&= \int_N \int_{\bar{M}} \int_K f(h^{-1}n_1mk)\varepsilon(n_1mk)A_2(\varphi(n_1mk))\delta_M^{-1}(m) dk dm dn_1 \\
&= \int_N \int_{\bar{M}} \int_K f(h^{-1}n_1mk)\varepsilon(mk)\delta_M^{-1/2}(m)A_2(\tau(m)\varphi(k)) dk dm dn_1.
\end{aligned}$$

Writing a general $m \in M$ in the form (a, g_2) , where $a \in k^\times$ and $g_2 \in \mathrm{GL}(2, k)$, the above expression is equal to

$$\begin{aligned}
&\int_N \int_{\bar{M}} \int_K f(h^{-1}n_1mk)\varepsilon(g_2)\varepsilon(k)\delta_M^{-1/2}(m)\mu(a)A_2\tau_2(g_2)\varphi(k) dk dm dn_1 \\
&= \int_N \int_{\bar{M}} \int_K f(h^{-1}n_1mk)\varepsilon(g_2)\varepsilon(k)\delta_M^{-1/2}(m)\mu(a)\varepsilon(g_2)\tau_2(g_2)A_2\varphi(k) dk dm dn_1 \\
&= \int_N \int_{\bar{M}} \int_K f(h^{-1}n_1mk)\varepsilon(k)\delta_M^{-1/2}(m)\mu(a)\tau_2(g_2)A_2\varphi(k) dk dm dn_1.
\end{aligned}$$

For each $m \in \bar{M}$, we change variables $n_1 \mapsto n$, where n is defined by $n^{-1}mnm^{-1} = n_1$. The resulting Jacobian is $|\det(1 - \mathrm{Ad} m)|_{\mathfrak{n}}$. Let $D_{M \setminus G}(m) = |\det(1 - \mathrm{Ad} m)|_{\mathfrak{g}/\mathfrak{m}}$, where lower case gothic type denotes the Lie algebra of a group. Since

$$D_{M \setminus G}^{1/2}(m) = \delta_M^{-1/2}(m) |\det(1 - \mathrm{Ad} m)|_{\mathfrak{n}},$$

the trace $(\pi(f)A\varphi)(h)$ is equal to

$$\int_N \int_{\bar{M}} \int_K f(h^{-1}n^{-1}mnk)\varepsilon(k)D_{M \setminus G}^{1/2}(m)\mu(a)\pi(g_2)A_2\varphi(k) dk dm dn.$$

Since $G = PK$ and functions in V transform under P via τ , we may identify V with a space of functions on K . The operator $\varphi \mapsto \pi(f)A\varphi$ on V is therefore an integral operator with kernel

$$\begin{aligned}
(h, k) \in K \times K &\mapsto \\
&\int_N \int_{\bar{M}} f(h^{-1}n^{-1}mnk)\varepsilon(k)D_{M \setminus G}^{1/2}(m)\mu(a)\pi(g_2)A_2 dm dn.
\end{aligned}$$

This kernel is an operator on the vector space V_2 of τ_2 , for each $(h, k) \in K \times K$.

We compute $\langle \pi, f \rangle_A = \mathrm{tr} \pi(f)A$ by integrating the character of this kernel (as an operator on V_2) over the diagonal $\{(k, k) : k \in K\}$. We have:

$$\begin{aligned}
\langle \pi, f \rangle_A &= \int_K \int_N \int_{\bar{M}} f(k^{-1}n^{-1}mnk)\varepsilon(k)D_{M \setminus G}^{1/2}(m)\mu(a)\chi_{\tau_2}^\varepsilon(g_2) dm dn dk \\
&= \int_{\bar{M}} D_{M \setminus G}^{1/2}(m)\mu(a)\chi_{\tau_2}^\varepsilon(g_2) \int_K \int_N f(k^{-1}n^{-1}mnk)\varepsilon(k) dn dk dm,
\end{aligned}$$

Let $N_G(M)$ be the normalizer of M in G . Let $W_{M \setminus G} = N_G(M)/M$. Then, $|W_{M \setminus G}| = 2$, and the nontrivial element of $W_{M \setminus G}$ sends (a, g_2) to $((\det g_2)/a, g_2)$. We

have:

$$\begin{aligned} \langle \pi, f \rangle_A &= \frac{1}{|W_{M \setminus G}|} \int_{\bar{M}} \left(\mu(a) + \mu \left(\frac{\det g_2}{a} \right) \right) D_{M \setminus G}^{1/2}(m) \chi_{\tau_2}^\varepsilon(g_2) \\ &\quad \cdot \left[\int_K \int_N f(k^{-1} n^{-1} m n k) \varepsilon(k) \, dn \, dk \right] dm. \end{aligned}$$

Let t_2 be a regular element in $\mathrm{GL}(2, k)$. For any $g_2 \in \mathrm{GL}(2, k)$, put $t_2^{g_2} := g_2^{-1} t_2 g_2$. Recall that $\chi_{\tau_2}^\varepsilon(t_2)$ is zero unless t_2 is conjugate to an element in T_K , and $\chi_{\tau_2}^\varepsilon(t_2^{g_2}) = \varepsilon(g_2) \chi_{\tau_2}^\varepsilon(t_2)$ for all $g_2 \in \mathrm{GL}(2, k)$. By the Weyl integration formula, if a function f on a group H is zero outside of the conjugacy class of a maximal torus T , then

$$(5.5) \quad \int_H f(h) \, dh = \frac{1}{|W_{T \setminus H}|} \int_{T^{\mathrm{reg}}} D_{T \setminus H}(t) \int_{T \setminus H} f(\bar{h}^{-1} t \bar{h}) \, d\bar{h} \, dt.$$

Here, $D_{T \setminus H}(t) := |\det(1 - \mathrm{Ad} \, t)|_{\mathfrak{h}/\mathfrak{t}}|$. Viewing $\langle \pi, f \rangle_A$ as an integral over $\bar{M}$, the integrand vanishes outside of the union of the tori conjugate to T in M . Applying (5.5), we obtain: $\langle \pi, f \rangle_A =$

$$\begin{aligned} &\frac{1}{|W_{T \setminus M}|} \frac{1}{|W_{M \setminus G}|} \int_{t=(a, t_2) \in \bar{T}^{\mathrm{reg}}} D_{T \setminus M}(t) \\ &\quad \cdot \int_{\bar{m}=(b, g_2) \in T \setminus M} D_{M \setminus G}^{1/2}(\bar{m}^{-1} t \bar{m}) \left[\mu(a^b) + \mu \left(\frac{\det t_2^{g_2}}{a^b} \right) \right] \chi_{\tau_2}^\varepsilon(t_2^{g_2}) \\ &\quad \cdot \int_K \int_N f(k^{-1} n^{-1} \bar{m}^{-1} t \bar{m} n k) \varepsilon(k) \, dn \, dk \, d\bar{m} \, dt \\ &= \frac{1}{|W_{T \setminus G}|} \int_{t=(a, t_2) \in \bar{T}^{\mathrm{reg}}} D_{T \setminus M}(t) D_{M \setminus G}^{1/2}(t) \\ &\quad \int_{\bar{m}=(b, g_2) \in T \setminus M} \left[(\mu(a^b) + \mu \left(\frac{\det t_2^{g_2}}{a^b} \right)) \right] \chi_{\tau_2}^\varepsilon(t_2^{g_2}) \\ &\quad \int_K \int_N f(k^{-1} n^{-1} \bar{m}^{-1} t \bar{m} n k) \varepsilon(k) \, dn \, dk \, d\bar{m} \, dt. \end{aligned}$$

Here, $W_{T \setminus G}$ denotes the Weyl group of T in G .

For $m = (b, g_2) \in M$, we have $\chi_{\tau_2}^\varepsilon(t_2^{g_2}) = \varepsilon(g_2) \chi_{\tau_2}^\varepsilon(t_2) = \varepsilon(m) \chi_{\tau_2}^\varepsilon(t_2)$. Inserting $\varepsilon(n) = 1$ ($n \in N$) at an appropriate place, we express $\langle \pi, f \rangle_A$ as follows:

$$\begin{aligned} &\frac{1}{|W_{T \setminus G}|} \int_{t=(a, t_2) \in \bar{T}^{\mathrm{reg}}} D_{T \setminus M}(t) D_{M \setminus G}^{1/2}(t) \left[\mu(a) + \mu \left(\frac{\det t_2}{a} \right) \right] \chi_{\tau_2}^\varepsilon(t_2) \\ &\quad \cdot \int_{T \setminus M} \varepsilon(\bar{m}) \int_K \int_N f(k^{-1} n^{-1} \bar{m}^{-1} t \bar{m} n k) \varepsilon(n k) \, dn \, dk \, d\bar{m} \, dt \\ &= \frac{1}{|W_{T \setminus G}|} \int_{\bar{T}^{\mathrm{reg}}} D_{T \setminus M}(t) D_{M \setminus G}^{1/2}(t) \left[\mu(a) + \mu \left(\frac{\det t_2}{a} \right) \right] \chi_{\tau_2}^\varepsilon(t_2) \\ &\quad \cdot \int_{T \setminus M} \int_K \int_N \varepsilon(\bar{m} n k) f(k^{-1} n^{-1} \bar{m}^{-1} t \bar{m} n k) \, dn \, d\bar{m} \, dt \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{|W_{T \setminus G}|} \int_{\bar{T}^{\text{reg}}} \left[\mu(a) + \mu\left(\frac{\det t_2}{a}\right) \right] D_{T \setminus M}^{1/2}(t) \chi_{\tau_2}^\varepsilon(t_2) \\
&\quad \cdot D_{T \setminus M}^{1/2}(t) D_{M \setminus G}^{1/2}(t) \int_{T \setminus G} \varepsilon(\bar{g}) f(\bar{g}^{-1} t \bar{g}) d\bar{g} dt.
\end{aligned}$$

Since $M \cong k^\times \times \text{GL}(2, k)$, $D_{T \setminus M}((a, t_2))$ is equal to $D_{T_K}(t_2) := D_{T_K \setminus \text{GL}(2, k)}(t_2)$. Moreover, $D_{T \setminus M} D_{M \setminus G} = D_{T \setminus G}$. Hence,

$$\begin{aligned}
(5.6) \quad \langle \pi, f \rangle_A &= \frac{1}{|W_{T \setminus G}|} \int_{t=(a, t_2) \in \bar{T}^{\text{reg}}} \left[\mu(a) + \mu\left(\frac{\det t_2}{a}\right) \right] D_{T_K}^{1/2}(t_2) \chi_{\tau_2}^\varepsilon(t_2) \\
&\quad \cdot D_{T \setminus G}^{1/2}(t) O_G^\varepsilon(f, t) dt.
\end{aligned}$$

By (5.3), the above expression is equal to

$$\begin{aligned}
&\frac{1}{|W_{T \setminus G}|} \int_{t=(a, \phi(\gamma)) \in \bar{T}^{\text{reg}}} \left[\mu(a) + \mu\left(\frac{N_{K/k} \gamma}{a}\right) \right] \cdot \left[\chi(\gamma) + \chi(\bar{\gamma}) \right] \\
&\quad \cdot \lambda(K/k, \psi) \varepsilon(\gamma - \bar{\gamma}) D_{T \setminus G}^{1/2}(t) O_G^\varepsilon(f, t) dt.
\end{aligned}$$

Since $|W_{T \setminus G}| = 4$, the lemma follows. $\square$

Let P_1 be the upper triangular parabolic subgroup of H_1 . Its Levi component is the maximal diagonal torus

$$T_1 = \{(\text{diag}(a, b), x) : a, b \in k^\times, x \in K^\times, ab = N_{K/k} x\}.$$

For characters μ_1, μ_2 of $k^\times$ and a character χ of $K^\times$, let $\mu_1 \otimes \mu_2 \otimes_1 \chi$ denote the representation of P_1 defined as follows:

$$\mu_1 \otimes \mu_2 \otimes_1 \chi : \left(\begin{pmatrix} a & * \\ 0 & b \end{pmatrix}, x \right) \mapsto \mu_1(a) \mu_2(b) \chi(x).$$

Let $I(\mu_1, \mu_2) \otimes_1 \chi$ denote the representation of H_1 normalizedly induced from $\mu_1 \otimes \mu_2 \otimes_1 \chi$.

Due to the condition that each element $(\text{diag}(a, b), x) \in T_1$ satisfies $ab = N_{K/k} x$, the H_1 -module $I(\mu_1, \mu_2) \otimes_1 \chi$ is equal to $\eta^{-1} I(\mu_1, \mu_2) \otimes_1 (\eta \circ N_{K/k}) \cdot \chi$ for any character η of $k^\times$. In particular, we have:

$$I(\mu_1, \mu_2) \otimes_1 \chi = I(\mu_1 \mu_2^{-1}, 1) \otimes_1 (\mu_2 \circ N_{K/k}) \cdot \chi.$$

Hence, any H_1 -module induced from P_1 has the form $I(\mu, 1) \otimes_1 \chi$, where μ, χ are characters of $k^\times, K^\times$, respectively.

Let $\bar{T}_1 = T_1/Z_0(H_1)$. For $a \in k^\times, \gamma \in K^\times$, recall that $(a, \phi(\gamma))$ denotes the element $\text{diag}(a, \phi(\gamma), (N_{K/k} \gamma)/a) \in T \subset G$. The map

$$\lambda_1 : (a, \phi(\gamma)) \mapsto \left(\begin{pmatrix} a & N_{K/k} \gamma \\ 0 & a \end{pmatrix}, \gamma \right)$$

defines an isomorphism from $\bar{T}$ to $\bar{T}_1$. Let dt_1 be the Haar measure on $\bar{T}_1$ which is compatible with the measure dt on $\bar{T}$ via λ_1 . That is, for each measurable subspace $\bar{S} \subset \bar{T}$, we have $\int_{\lambda_1(\bar{S})} dt_1 = \int_{\bar{S}} dt$.

For $t_1 \in T_1$ and $f_1 \in C(H_1)$, let

$$O_{H_1}(f_1, t_1) = \int_{T_1 \setminus H_1} f_1(\bar{h}_1^{-1} t_1 \bar{h}_1) d\bar{h}_1.$$

Let μ, χ be the characters of $k^\times, K^\times$, respectively. The split central character of $I(\mu, 1) \otimes_1 \chi$ is $\chi|_{k^\times} \cdot \mu$.

5.4. LEMMA. For $f_1 \in C(H_1, \chi|_{k^\times} \cdot \mu)$, the following holds:

$$(5.7) \quad \langle I(\mu, 1) \otimes_1 \chi, f_1 \rangle = \frac{1}{4} \int_{t_1 = (\text{diag}(a, b), \gamma) \in \bar{T}_1^{\text{reg}}} [\mu(a) + \mu(b)] [\chi(\gamma) + \chi(\bar{\gamma})] \cdot D_{T_1 \setminus H_1}^{1/2}(t_1) O_{H_1}(f_1, t_1) dt_1.$$

PROOF. We skip the proof, which is similar to that of Lemma 5.3 and is simpler because there is no twisting by ε . $\square$

Recall the embedding $\phi : \mathcal{K}^\times \hookrightarrow \text{GL}(2, k)$ which sends $a + b\sqrt{B}$ to $\begin{pmatrix} a & bB \\ b & a \end{pmatrix}$. Note that $\det \phi(\gamma) = N_{\mathcal{K}/k} \gamma$ for all $\gamma \in \mathcal{K}^\times$. Given $a, b \in k^\times, \gamma \in \mathcal{K}^\times - k^\times$, such that $ab = N_{\mathcal{K}/k} \gamma$ and $a \neq b$, the element $t = \text{blockdiag}(a, \phi(\gamma), b) \in M$ is regular. Every norm of t in H_1 is conjugate to either $t_1 = (\text{diag}(a, b), \gamma)$ or $\bar{t}_1 = (\text{diag}(a, b), \bar{\gamma})$. In particular, t_1 and $\bar{t}_1$ are $\mathbf{G}$ -regular. We make the following choice of transfer factors for the pairs (t_1, t) , $(\bar{t}_1, t)$:

$$\Delta(t_1, t) = \Delta(\bar{t}_1, t) = \lambda(\mathcal{K}/k, \psi) \varepsilon(a) \varepsilon(\gamma - \bar{\gamma}) \left[\frac{D_{T \setminus G}(t)}{D_{T_1 \setminus H_1}(t_1)} \right]^{1/2}.$$

5.5. DEFINITION. We say that $f \in C(G)$ and $f \in C(H_1)$ are **weakly matching** if

$$O_{H_1}(f_1, t_1) = \Delta(t_1, t) O_G^\varepsilon(f, t)$$

for every $(t_1, t) \in T_1 \times T^{\text{reg}}$ such that t_1 is a norm of t .

The assumption is that there exists a choice of transfer factors for $H_1 \times G$, compatible with $\Delta(t_1, t)$, such that matching implies weakly matching.

5.6. PROPOSITION. Let μ be a character of $k^\times$. Let χ be a character of $\mathcal{K}^\times$. For weakly matching functions, the following character identity holds:

$$\langle \mu \rtimes \pi(\chi), f \rangle_A = \langle I(\varepsilon\mu, 1) \otimes_1 \chi, f_1 \rangle.$$

PROOF. The elements in H_1^{reg} which are not $\mathbf{G}$ -regular form a set of measure zero. Hence, by Lemma 5.4 and our choice of dt_1 , $\langle I(\varepsilon\mu, 1) \otimes_1 \chi, f_1 \rangle$ is equal to

$$(5.8) \quad \begin{aligned} & \frac{1}{4} \int_{t=(a, \phi(\gamma)) \in \bar{T}^{\text{reg}}} \left[\varepsilon\mu(a) + \varepsilon\mu\left(\frac{N_{\mathcal{K}/k}\gamma}{a}\right) \right] [\chi(\gamma) + \chi(\bar{\gamma})] \\ & \quad \cdot D_{T_1 \setminus H_1}^{1/2}(\lambda_1(t)) O_{H_1}(f_1, \lambda_1(t)) dt \\ & = \frac{1}{4} \int_{t=(a, \phi(\gamma)) \in \bar{T}^{\text{reg}}} \varepsilon(a) \left[\mu(a) + \mu\left(\frac{N_{\mathcal{K}/k}\gamma}{a}\right) \right] [\chi(\gamma) + \chi(\bar{\gamma})] \\ & \quad \cdot D_{T_1 \setminus H_1}^{1/2}(\lambda_1(t)) O_{H_1}(f_1, \lambda_1(t)) dt. \end{aligned}$$

Since the functions $f_1 \in C(H_1)$ and $f \in C(G)$ are weakly matching, we have:

$$O_{H_1}(f_1, \lambda_1(t)) = \lambda(\mathcal{K}/k, \psi) \varepsilon(a) \varepsilon(\gamma - \bar{\gamma}) \left[\frac{D_{T \setminus G}(t)}{D_{T_1 \setminus H_1}(\lambda_1(t))} \right]^{1/2} O_G^\varepsilon(f, t)$$

for all regular $t = (a, \phi(\gamma)) \in T$. The expression (5.8) is therefore equal to

$$(5.9) \quad \begin{aligned} & \frac{1}{4} \int_{t=(a, \phi(\gamma)) \in \bar{T}^{\text{reg}}} \left[\mu(a) + \mu\left(\frac{N_{\mathcal{K}/k}\gamma}{a}\right) \right] [\chi(\gamma) + \chi(\bar{\gamma})] \\ & \quad \cdot \lambda(\mathcal{K}/k, \psi) \varepsilon(\gamma - \bar{\gamma}) D_{T \setminus G}^{1/2}(t) O_G^\varepsilon(f, t) dt. \end{aligned}$$

By Lemma 5.3, the proposition follows. $\square$

5.1.3. A Trace Identity between G and H_2 . Let $T_2 = \{(\text{diag}(\alpha, \beta), c)_* : \alpha, \beta \in \mathcal{K}^\times, c \in k^\times\}$ be the maximal diagonal torus of H_2 . Here, lower $*$ denotes the image of $(\text{diag}(\alpha, \beta), c)$ in H_2 . Let $\bar{T}_2 = T_2/Z_0(H_2)$.

Recall that $T = \{(a, \phi(\gamma)) : a \in k^\times, \gamma \in \mathcal{K}^\times\} \subset G$, where $(a, \phi(\gamma))$ denotes the element $\text{blockdiag}(a, \phi(\gamma), (\det \phi(\gamma))/a)$. Any element in T has the form

$$t = (cN_{\mathcal{K}/k}\alpha, c\phi(\alpha\bar{\beta}))$$

for some $c \in k^\times, \alpha, \beta \in \mathcal{K}^\times$.

The norm correspondence between G and H_2 gives the following isomorphism from $\bar{T}$ to $\bar{T}_2$:

$$\lambda_2 : (cN_{\mathcal{K}/k}\alpha, c\phi(\alpha\bar{\beta})) \mapsto (\text{diag}(\alpha, \beta), c)_*, \quad \forall \alpha, \beta \in \mathcal{K}^\times, c \in k^\times.$$

Let dt_2 be the Haar measure on $\bar{T}_2$ which is compatible with the measure dt on $\bar{T}$ via λ_2 .

For $t_2 \in T_2$ and $f_2 \in C(H_2)$, let

$$O_{H_2}(f_2, t_2) = \int_{T_2 \setminus H_2} f_2(\bar{h}_2^{-1}t_2\bar{h}_2) d\bar{h}_2.$$

Let c be an element in $k^\times$. Let α, β be elements in $\mathcal{K}^\times$. Suppose $\alpha\bar{\beta} \notin k^\times$, then the element $t = (cN_{\mathcal{K}/k}\alpha, c\phi(\alpha\bar{\beta})) \in T$ is regular. A norm of t in H_2 is conjugate to either $t_2 = (\text{diag}(\alpha, \beta), c)_*$ or $\bar{t}_2 = (\text{diag}(\bar{\alpha}, \bar{\beta}), c)_*$. Extend the character ε of $k^\times$ to $\mathcal{K}^\times$. We make the following choice of transfer factors for the pairs (t_2, t) , $(\bar{t}_2, t)$:

$$\begin{aligned} \Delta(t_2, t) &= \Delta(\bar{t}_2, t) = \lambda(\mathcal{K}/k, \psi)\varepsilon(cN_{\mathcal{K}/k}\alpha)\varepsilon(c\alpha\bar{\beta} - c\bar{\alpha}\beta) \left[\frac{D_{T \setminus G}(t)}{D_{T_2 \setminus H_2}(t_2)} \right]^{1/2} \\ &= \lambda(\mathcal{K}/k, \psi)\varepsilon(\alpha\bar{\beta} - \bar{\alpha}\beta) \left[\frac{D_{T \setminus G}(t)}{D_{T_2 \setminus H_2}(t_2)} \right]^{1/2}. \end{aligned}$$

5.7. DEFINITION. We say that $f \in C(G)$ and $f_2 \in C(H_2)$ are **weakly matching** if

$$O_{H_2}(f_2, t_2) = \Delta(t_2, t)O_G^\varepsilon(f, t)$$

for every $(t_2, t) \in T_2 \times T^{\text{reg}}$ such that t_2 is a norm of t .

5.8. PROPOSITION. Let μ, χ be characters of $k^\times, \mathcal{K}^\times$, respectively. For weakly matching functions f in $C(G)$ and f_2 in $C(H_2)$, the following character identity holds:

$$\left\langle \frac{\varepsilon\mu}{\chi|_{k^\times}} \rtimes \pi(\chi), f \right\rangle_A = \left\langle I\left(\frac{\mu \circ N}{\chi}, \chi\right) \otimes_2 \mu, f_2 \right\rangle.$$

Here, A is the intertwining operator defined in Section 5.1.1.

PROOF. For simplicity, let N denote $N_{\mathcal{K}/k}$. By Lemma 5.3, $\left\langle \frac{\varepsilon\mu}{\chi|_{k^\times}} \rtimes \pi(\chi), f \right\rangle_A$ is equal to

$$\begin{aligned} (5.10) \quad & \frac{1}{4} \int_{t=(cN\alpha, c\phi(\alpha\bar{\beta})) \in \bar{T}^{\text{reg}}} \varepsilon\mu(c) \\ & \cdot \left[\mu(N\alpha) \left(\chi\left(\frac{\beta}{\alpha}\right) + \chi\left(\frac{\bar{\beta}}{\bar{\alpha}}\right) \right) + \mu(N\beta) \left(\chi\left(\frac{\alpha}{\beta}\right) + \chi\left(\frac{\bar{\alpha}}{\bar{\beta}}\right) \right) \right] \\ & \cdot \lambda(\mathcal{K}/k, \psi)\varepsilon(c\alpha\bar{\beta} - c\bar{\alpha}\beta) D_{T \setminus G}^{1/2}(t) O_G^\varepsilon(f, t) dt. \end{aligned}$$

Employing the same technique used in the proof of Lemma 5.3, we obtain:

$$(5.11) \quad \left\langle I\left(\frac{\mu \circ N}{\chi}, \chi\right) \otimes_2 \mu, f_2 \right\rangle = \frac{1}{|\text{Gal}(\mathcal{K}/k)|} \cdot \frac{1}{|W_{T_2 \setminus H_2}|} \cdot \int_{t_2 = (\text{diag}(\alpha, \beta), c)_* \in \bar{T}_2^{\text{reg}}} \left[\mu(N\alpha) \left(\chi\left(\frac{\beta}{\alpha}\right) + \chi\left(\frac{\bar{\beta}}{\alpha}\right) \right) + \mu(N\beta) \left(\chi\left(\frac{\alpha}{\beta}\right) + \chi\left(\frac{\bar{\alpha}}{\beta}\right) \right) \right] \cdot \mu(c) D_{T_2 \setminus H_2}^{1/2}(t_2) O_{H_2}(f_2, t_2) dt_2.$$

By our choice of dt_2 , and the fact that $|\text{Gal}(\mathcal{K}, k)| = |W_{T_2 \setminus H_2}| = 2$, the above expression is equal to

$$\frac{1}{4} \int_{t = (cN\alpha, c\phi(\alpha\bar{\beta})) \in \bar{T}^{\text{reg}}} \left[\mu(N\alpha) \left(\chi\left(\frac{\beta}{\alpha}\right) + \chi\left(\frac{\bar{\beta}}{\alpha}\right) \right) + \mu(N\beta) \left(\chi\left(\frac{\alpha}{\beta}\right) + \chi\left(\frac{\bar{\alpha}}{\beta}\right) \right) \right] \cdot \mu(c) D_{T_2 \setminus H_2}^{1/2}(\lambda_2(t)) O_{H_2}(f_2, \lambda_2(t)) dt.$$

The proposition now follows from the weakly matching condition on f, f_2 . $\square$

REMARK. By the Weyl integration formula and the Iwasawa decomposition for real reductive groups, it is clear that the archimedean analogues of Propositions 5.6, 5.8 also hold. That is: If $k = \mathbb{R}$, $\mathcal{K} = \mathbb{C}$, and μ, χ are characters of $\mathbb{R}^\times, \mathbb{C}^\times$, respectively, then

$$\begin{aligned} \langle \mu \rtimes \pi(\chi), f \rangle_A &= \langle I(\varepsilon\mu, 1) \otimes_1 \chi, f_1 \rangle, \\ \left\langle \frac{\varepsilon\mu}{\chi|_{k^\times}} \rtimes \pi(\chi), f \right\rangle_A &= \left\langle I\left(\frac{\mu \circ N}{\chi}, \chi\right) \otimes_2 \mu, f_2 \right\rangle \end{aligned}$$

for matching functions on $\mathbf{G}(\mathbb{R}), \mathbf{H}_1(\mathbb{R}), \mathbf{H}_2(\mathbb{R})$. Here, $\varepsilon(x)$ is the sign of x for all $x \in \mathbb{R}$.

5.1.4. Examples. For an admissible representation τ of $\text{GL}(2, k)$, let $B_{\mathcal{K}/k}\tau$ denote the representation of $\text{GL}(2, \mathcal{K})$ which is obtained from τ via base change ([L], [F1]). In particular, for characters μ_1, μ_2 of $k^\times$, $B_{\mathcal{K}/k}I(\mu_1, \mu_2)$ is the induced representation $I(\mu_1 \circ N_{\mathcal{K}/k}, \mu_2 \circ N_{\mathcal{K}/k})$.

For $x \in k$, let $\text{ord } x$ denote the p -adic valuation of x , with $\text{ord } 0 := \infty$. Let q be the cardinality of the residue field of k . Let $\nu : x \mapsto q^{-\text{ord } x}$ be the normalized absolute value function on k .

5.9. COROLLARY. *Let θ be a character of $\mathcal{K}^\times$. The following holds for weakly matching functions $f \in C(G)$ and $f_1 \in C(H_1)$:*

$$\left\langle \varepsilon\nu \rtimes \nu^{-1/2}\pi(\theta), f \right\rangle_A = \left\langle I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 \theta, f_1 \right\rangle.$$

PROOF. Noting that $I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 \theta = I(\nu, 1) \otimes_1 (\nu^{-1/2} \circ N_{\mathcal{K}/k}) \cdot \theta$, the corollary follows from Proposition 5.6. $\square$

5.10. COROLLARY. *Let η be a character of $k^\times$. For weakly matching functions f, f_1, f_2 on G, H_1, H_2 , respectively, the following quantities are equal to one another:*

- $\left\langle \varepsilon\nu \rtimes \nu^{-1/2}\eta, f \right\rangle_A,$
- $\left\langle \eta I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 1, f_1 \right\rangle,$
- $\left\langle B_{\mathcal{K}/k}I(\eta\nu^{1/2}, \eta\nu^{-1/2}) \otimes_2 \eta^2, f_2 \right\rangle.$

PROOF. The representation $\varepsilon\nu \times \varepsilon \rtimes \nu^{-1/2}\eta$ is equal to $\varepsilon\nu \rtimes \pi((\nu^{-1/2}\eta) \circ N_{\mathcal{K}/k})$. The representation $\eta I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 1$ of H_1 is equal to $I(\nu, 1) \otimes_1 (\nu^{-1/2}\eta) \circ N_{\mathcal{K}/k}$. The corollary follows from Corollary 5.9 and Proposition 5.8. $\square$

Let $\nu_{\mathcal{K}}$ be the normalized absolute value function on $\mathcal{K}$.

5.11. COROLLARY. *Suppose the extension $\mathcal{K}/k$ is unramified. Let θ be a character of $\mathcal{K}^\times$ which satisfies ${}^\sigma\theta/\theta = \varepsilon' \circ N_{\mathcal{K}/k}$ for some quadratic character ε' of $k^\times$. The following holds for weakly matching functions $f \in C(G)$ and $f_2 \in C(H_2)$:*

$$\left\langle \varepsilon\varepsilon'\nu \rtimes \nu^{-1/2}\pi(\theta), f \right\rangle_A = \left\langle \theta I(\nu_{\mathcal{K}}^{1/2}, \nu_{\mathcal{K}}^{-1/2}) \otimes_2 \theta|_{k^\times} \cdot \varepsilon', f_2 \right\rangle.$$

PROOF. Since $\mathcal{K}/k$ is unramified, $\nu_{\mathcal{K}}$ is equal to $\nu \circ N_{\mathcal{K}/k}$. The corollary then follows from Proposition 5.8. $\square$

5.2. Induced Representations—Split Case

Let F be a number field. Let C_F denote the idèle class group of F . Let $\mathcal{E}$ be a nontrivial quadratic character of C_F . Let E be the quadratic extension of F which corresponds to $\mathcal{E}$ via global class field theory.

Regard $\mathbf{G} = \mathrm{GSp}(2)$ as an F -group. Let $\mathbf{H}_1$ and $\mathbf{H}_2$ be the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let v be a (possibly archimedean) place of F which splits in E . Then, $E_v := E \otimes_F F_v = F_v \oplus F_v$. We have:

$$\begin{aligned} \mathbf{H}_1(F_v) &= (\mathrm{GL}(2, F_v) \times F_v^\times \times F_v^\times)' \\ &:= \{(g, a, b) \in \mathrm{GL}(2, F_v) \times F_v^\times \times F_v^\times : \det g = ab\}, \end{aligned}$$

$$\mathbf{H}_2(F_v) = (\mathrm{GL}(2, F_v) \times \mathrm{GL}(2, F_v) \times F_v^\times) / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in F_v^\times\}.$$

Let $G_v = \mathbf{G}(F_v)$, $H_{i,v} = \mathbf{H}_i(F_v)$ ($i = 1, 2$).

Let M_v denote the Levi component of the Heisenberg parabolic subgroup of G_v . The norm correspondence between $\mathbf{G}$ and $\mathbf{H}_1$ gives the following isomorphism from M_v to $H_{1,v}$:

$$\lambda_1 : (a, g_2) \mapsto (g_2, a, (\det g_2)/a), \quad \forall a \in k^\times, g_2 \in \mathrm{GL}(2, k).$$

Here, (a, g_2) denotes the element $\mathrm{blockdiag}(a, g_2, (\det g_2)/a)$ in M_v . In particular, λ_1 defines a one to one correspondence between the conjugacy classes of the maximal tori in M_v and those of the maximal tori in $H_{1,v}$.

Each maximal torus in M_v has the form

$$T_v = \{(a, t') : a \in F_v^\times, t' \in T'_v\},$$

where T'_v is a maximal torus in $\mathrm{GL}(2, F_v)$.

A norm in $H_{1,v}$ of a regular element $t = (a, t') \in M_v$ is conjugate to either $t_1 = (t', a, (\det t')/a)$ or $\bar{t}_1 = (t', (\det t')/a, a)$. Let $T_v, T_{1,v}$ be the maximal tori which contain t, t_1 , respectively. We define the transfer factors for the pairs $(t_1, t), (\bar{t}_1, t)$ as follows:

$$\Delta(t_1, t) = \Delta(\bar{t}_1, t) = \left[\frac{D_{T_v \setminus G_v}(t)}{D_{T_{1,v} \setminus H_{1,v}}(t_1)} \right]^{1/2}.$$

Let Z_v be the center of G_v . Let $C(G_v)$ denote the space of functions on G_v which are smooth and compactly supported modulo Z_v . Let $Z'(H_{1,v})$ denote the subgroup

$$\{(\mathrm{diag}(z, z), z, z) : z \in F_v^\times\} \subset H_{1,v}.$$

It is the group of F_v -points of the maximal F -split component of the center of $\mathbf{H}_1$. Let $C(H_{1,v})$ denote the space of functions on $H_{1,v}$ which are smooth and compactly supported modulo $Z'(H_{1,v})$.

For a maximal torus $T_{1,v}$ in $H_{1,v}$, put $\bar{T}_{1,v} := T_{1,v}/Z'(H_{1,v})$. Put $\bar{G}_v := G_v/Z_v$, $\bar{H}_{1,v} := H_{1,v}/Z'(H_{1,v})$. We fix a Haar measure dt on each maximal torus $\bar{T}_v$ in $\bar{G}_v$. It induces via λ_1 a measure dt_1 on the maximal torus $\bar{T}_{1,v} = \lambda_1(\bar{T}_v)$ in $\bar{H}_{1,v}$.

We say that $f_v \in C(G_v)$ and $f_{1,v} \in C(H_{1,v})$ are **weakly matching** if

$$O_{H_{1,v}}(f_{1,v}, t_1) = \Delta(t_1, t) O_{G_v}(f_v, t)$$

for all $(t_1, t) \in H_{1,v} \times M_v^{\text{reg}}$ such that t_1 is a norm of t .

5.12. LEMMA. *Let τ be an admissible representation of $\text{GL}(2, F_v)$. Let μ_1, μ_2 be characters of $F_v^\times$. Then,*

$$\left\langle \frac{\mu_1}{\mu_2} \rtimes \mu_2 \tau, f_v \right\rangle = \langle \tau \otimes_1 (\mu_1 \otimes \mu_2), f_{1,v} \rangle$$

for weakly matching functions.

PROOF. The proof is similar to that of Proposition 5.6.

By the Weyl integration formula, $\left\langle \frac{\mu_1}{\mu_2} \rtimes \mu_2 \tau, f \right\rangle$ is equal to

$$\begin{aligned} & \sum_{T_v} \frac{1}{|W_{T_v \setminus M_v}|} \int_{t=(a, t') \in \bar{T}_v^{\text{reg}}} D_{T_v \setminus M_v}(t) \frac{\mu_1}{\mu_2}(a) \chi_{\mu_2 \tau}(t') D_{M_v \setminus G_v}^{1/2}(t) O_{G_v}(f_v, t) dt \\ &= \sum_{T_v} \frac{1}{|W_{T_v \setminus M_v}|} \\ & \quad \cdot \int_{t=(a, t') \in \bar{T}_v^{\text{reg}}} D_{T_v \setminus M_v}^{1/2}(t) \mu_1(a) \mu_2\left(\frac{\det t'}{a}\right) \chi_\tau(t') D_{T_v \setminus G_v}^{1/2}(t) O_{G_v}(f_v, t) dt. \end{aligned}$$

Here, T_v ranges over a set of representatives of the conjugacy classes of the maximal tori in M_v , and $W_{T_v \setminus M_v}$ is the Weyl group of T_v in M_v . The last equality holds because $D_{T_v \setminus M_v} D_{M_v \setminus G_v} = D_{T_v \setminus G_v}$.

The trace $\langle \tau \otimes_1 (\mu_1 \otimes \mu_2), f_{1,v} \rangle$ is equal to

$$\begin{aligned} & \sum_{T_{1,v}} \frac{1}{|W_{T_{1,v} \setminus H_{1,v}}|} \int_{t_1=(t', a, (\det t')/a) \in \bar{T}_{1,v}^{\text{reg}}} D_{T_{1,v} \setminus H_{1,v}}^{1/2}(t_1) \mu_1(a) \mu_2\left(\frac{\det t'}{a}\right) \chi_\tau(t') \\ & \quad \cdot D_{T_{1,v} \setminus H_{1,v}}^{1/2}(t_1) O_{H_{1,v}}(f_{1,v}, t_1) dt_1 \\ &= \sum_{\lambda_1(T_v)} \frac{1}{|W_{\lambda_1(T_v) \setminus H_{1,v}}|} \int_{t=(a, t') \in \bar{T}_v^{\text{reg}}} D_{T_{1,v} \setminus H_{1,v}}^{1/2}(\lambda_1(t)) \mu_1(a) \mu_2\left(\frac{\det t'}{a}\right) \chi_\tau(t') \\ & \quad \cdot D_{T_{1,v} \setminus H_{1,v}}^{1/2}(\lambda_1(t)) O_{H_{1,v}}(f_{1,v}, \lambda_1(t)) dt. \end{aligned}$$

Here, $T_{1,v}, T_v$ range over sets of representatives of the conjugacy classes of the maximal tori in $H_{1,v}, G_v$, respectively.

By assumption, $f_v, f_{1,v}$ are weakly matching functions; hence,

$$D_{T_v \setminus G_v}^{1/2}(t) O_{G_v}(f_v, t) = D_{T_{1,v} \setminus H_{1,v}}^{1/2}(\lambda_1(t)) O_{H_{1,v}}(f_{1,v}, \lambda_1(t))$$

for all regular t in M_v . Moreover, $|W_{T_v \setminus M_v}| = |W_{\lambda_1(T_v) \setminus H_{1,v}}| = 2$ for all maximal tori T_v in M_v . The lemma follows. $\square$

Let $T_{2,v}$ be the maximal diagonal torus in $H_{2,v}$. Let T_v be the maximal diagonal torus in G_v . Any element in T_v has the form $t = \text{diag}(ac, ad, bc, bd)$, $a, b, c, d \in F_v^\times$. The norm correspondence between $\mathbf{G}$ and $\mathbf{H}_2$ gives the following isomorphism from T_v to $T_{2,v}$:

$$\lambda_2 : \text{diag}(ac, ad, bc, bd) \mapsto (\text{diag}(a, b), \text{diag}(c, d), 1)_*.$$

Here, lower $*$ denotes image in $H_{2,v}$.

A norm in $H_{2,v}$ of a regular element $t = \text{diag}(ac, ad, bc, bd) \in T_v$ is conjugate to either $t_2 = (\text{diag}(a, b), \text{diag}(c, d), 1)_*$ or $\bar{t}_2 = (\text{diag}(c, d), \text{diag}(a, b), 1)_*$. We define the transfer factors for the pairs (t_2, t) , $(\bar{t}_2, t)$ as follows:

$$\Delta(t_2, t) = \Delta(\bar{t}_2, t) = \left[\frac{D_{T_v \setminus G_v}(t)}{D_{T_{2,v} \setminus H_{2,v}}(t_{2,v})} \right]^{1/2}.$$

Let $Z'(H_{2,v}) = \{(\text{diag}(1, 1), \text{diag}(1, 1), z)_* : z \in F_v^\times\} \subset H_{2,v}$. It is the group of F_v -points of the maximal F -split component of the center of $\mathbf{H}_2$. Let $C(H_{2,v})$ denote the space of smooth, compactly supported modulo $Z'(H_{2,v})$ functions on $H_{2,v}$.

Put $\bar{T}_{2,v} := T_{2,v}/Z'(H_{2,v})$. As usual, the Haar measure dt_2 on $\bar{T}_{2,v}$ is chosen such that it is compatible with the measure dt on $\bar{T}_v$ via λ_2 . We say that the functions $f_v \in C(G_v)$ and $f_{2,v} \in C(H_{2,v})$ are **weakly matching** if

$$O_{H_{2,v}}(f_{2,v}, t_2) = \Delta(t_2, t) O_{G_v}(f_v, t)$$

for every $(t_2, t) \in T_{2,v} \times T_v^{\text{reg}}$ such that t_2 is a norm of t .

5.13. LEMMA. *Let $\alpha_1, \alpha_2, \beta_1, \beta_2$ be characters of $F_v^\times$ such that $\alpha_1\beta_1 = \alpha_2\beta_2$. Then,*

$$\langle \beta_1^{-1} I(\alpha_2, \beta_2) \rtimes \beta_1, f_v \rangle = \langle I(\alpha_1, \beta_1) \times I(\alpha_2, \beta_2), f_{2,v} \rangle$$

for weakly matching functions $f_v \in C(G_v)$, $f_{2,v} \in C(H_{2,v})$.

PROOF. The induced representations are supported only on the conjugacy classes of the maximal diagonal tori. As in the proof of Lemma 5.12, the character identity follows from the Weyl integration formula and the weakly matching condition on the test functions. $\square$

5.3. ε -Invariant Packets

Recall that k is our fixed p -adic field, ε is a nontrivial quadratic character of $k^\times$, and $\mathcal{K}$ is the quadratic extension of k corresponding to ε . Let $G = \mathbf{G}(k)$, $H_i = \mathbf{H}_i(k)$ ($i = 1, 2$).

Let $\bar{G} = \text{PGSp}(2, k)$. Let $C_0 = \text{PGL}(2, k) \times \text{PGL}(2, k)$. For representations τ_1, τ_2 of $\text{PGL}(2, k)$, let $\tau_1 \times \tau_2$ denote the C_0 -module where $(g, h) \in C_0$ acts by $\tau_1(g) \otimes \tau_2(h)$ on the tensor product of the spaces of τ_1, τ_2 .

Let τ be a cuspidal or one dimensional representation of $\text{GL}(2, k)$ with trivial central character. Write $\dagger$ for $+$ if τ is cuspidal, or $\times$ if τ is one dimensional. Following the same notation as in [F4], we let $\{\pi^\dagger, \pi^-\}$ denote the local (quasi-)packet of $\bar{G}$ which lifts to the representation $I_{(2,2)}(\tau, \varepsilon\tau)$ of $\text{PGL}(4, k)$. In the terminology of [F4], which we also adopt, $\{\pi^+, \pi^-\}$ is a packet, while $\{\pi^\times, \pi^-\}$ is a quasi-packet.

5.14. LEMMA. *The representations $\pi^\dagger$ and π^- are ε -invariant.*

PROOF. Let f, f_4, f_{C_0} be matching functions on $\bar{G}$, $\text{PGL}(4, k)$, C_0 , respectively. Let $\zeta = 1$ or ε . By [F4, Prop. V. 5] and [F4, Prop. V. 8.6], we have:

$$\begin{aligned} \langle \zeta \pi^\dagger, f \rangle &= \frac{1}{2} \left(\langle \zeta I_{(2,2)}(\tau, \varepsilon\tau), f_4 \rangle + \langle \zeta \tau \times \zeta \varepsilon\tau, f_{C_0} \rangle \right) \\ \langle \zeta \pi^-, f \rangle &= \frac{1}{2} \left(\langle \zeta I_{(2,2)}(\tau, \varepsilon\tau), f_4 \rangle - \langle \zeta \tau \times \zeta \varepsilon\tau, f_{C_0} \rangle \right). \end{aligned}$$

The representation $I_{(2,2)}(\tau, \varepsilon\tau)$ is ε -invariant. By the norm correspondence between $\tilde{G}$ and C_0 , and the matching condition on the test functions, the convolution operator $(\tau \times \varepsilon\tau)(f_{C_0})$ is equal to $(\varepsilon\tau \times \tau)(f_{C_0})$. Hence, the right-hand sides of the above equations are independent of the choice of ζ . Consequently,

$$\begin{aligned}\langle \pi^\dagger, f \rangle &= \langle \varepsilon\pi^\dagger, f \rangle, \\ \langle \pi^-, f \rangle &= \langle \varepsilon\pi^-, f \rangle.\end{aligned}$$

By the linear independence of characters, we conclude that $\pi^\dagger$ and π^- are ε -invariant. $\square$

5.15. COROLLARY. *Let F be a totally imaginary number field. Let $\mathcal{E}$ be a nontrivial quadratic character of the idèle class group of F . Let T be a cuspidal or one dimensional automorphic representation of $\mathrm{GL}(2, \mathbb{A}_F)$ with trivial central character. Let $\{\Pi\}$ be the unstable (quasi)-packet $[T, \mathcal{E}T]$ of $\mathbf{G}(\mathbb{A}_F)$ (see Section 4.3.1). Then, every member of $\{\Pi\}$ is $\mathcal{E}$ -invariant.*

PROOF. Let Π be a member of $\{\Pi\}$. By Lemma 5.14, Π_v is $\mathcal{E}_v$ -invariant for each finite place v .

Let E be the quadratic extension of F corresponding to $\mathcal{E}$. Let v be an archimedean place of F . Since F is totally imaginary, F_v is algebraically closed. Thus, v splits in E , which implies $\mathcal{E}_v = 1$. $\square$

5.4. Character Identities for Unstable Packets

Notation

- For a number field F , let C_F denote the idèle class group of F .
- Suppose S is a set of places of a number field F . For any adèlic object (representation, test function, trace, ..., etc) over F , let subscript S denote the tensor product of local components over the places in S . For example, if Π denotes an irreducible representation of $\mathbf{G}(\mathbb{A}_F)$, we put $\Pi_S := \otimes_{v \in S} \Pi_v$.
- For an algebraic group $\mathbf{H}$ over a number field F , put $H_v := \mathbf{H}(F_v)$ for each place v of F .

5.4.1. One Dimensional Representations. Let $1_{2,l}$ ($l = k, \mathcal{K}$) denote the trivial representation of $\mathrm{GL}(2, l)$. It is nontempered. Let $\mathrm{St}_{2,l}$ denote the Steinberg representation of $\mathrm{GL}(2, l)$. It is square integrable.

Let ξ be a character of $k^\times$ with $\xi^2 = 1$. Let $\{\pi^\times, \pi^-\}$ be the local quasi-packet of G which lifts to the representation $I_{(2,2)}(\xi 1_{2,k}, \varepsilon \xi 1_{2,k})$ of $\mathrm{PGL}(4, k)$ according to [F4, Prop. V. 8.5]. In other words, $\pi^\times$ is the nontempered subquotient $L(\nu\varepsilon, \varepsilon \rtimes \nu^{-1/2}\xi)$ of the induced representation $\nu\varepsilon \times \varepsilon \rtimes \nu^{-1/2}\xi$ (see [ST]), and $\pi^- = \delta^-(\varepsilon\nu^{1/2}\mathrm{St}_{2,k}, \nu^{-1/2}\xi)$, the cuspidal member of the local packet which contains the unique square integrable subrepresentation $\delta(\varepsilon\nu^{1/2}\mathrm{St}_{2,k}, \nu^{-1/2}\xi)$ of $\nu\varepsilon \times \varepsilon \rtimes \nu^{-1/2}\xi$ (see [F4, Sect. V. 8]). By Lemma 5.14, $\pi^\times$ and π^- are ε -invariant representations.

5.16. PROPOSITION. *Suppose ε is unramified. There exist nontrivial intertwining operators $A^\times \in \mathrm{Hom}_G(\pi^\times, \varepsilon\pi^\times)$, $A^- \in \mathrm{Hom}_G(\pi^-, \varepsilon\pi^-)$ such that the following identities hold for matching functions:*

$$\begin{aligned}\langle \pi^\times, f \rangle_{A^\times} + \langle \pi^-, f \rangle_{A^-} &= \langle \xi 1_{2,k} \otimes 1, f_1 \rangle, \\ \langle \pi^\times, f \rangle_{A^\times} - \langle \pi^-, f \rangle_{A^-} &= \langle (\xi \circ N_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes 2, f_2 \rangle.\end{aligned}$$

REMARK. It is likely that the proposition still holds if we remove the condition that ε is unramified; however, the author is unable to prove it.

The proof will be given after Claim 5.19.

Using Krasner's lemma and the strong approximation theorem (see [PR]), we construct a totally imaginary number field F and a quadratic extension E of F such that:

- There exists a finite place w of F such that $F_w = k$;
- the place w is unramified, prime in E ;
- $E_w = \mathcal{K}$.

Here, w denotes both the place of F and the unique place of E which lies above it. Let V be the set of places of F . Let $V^{\text{un}}(E/F)$ be the set of places of F which are unramified in E .

Let $\mathcal{E}$ be the quadratic character of C_F which corresponds to the extension E/F via global class field theory. In particular, $\mathcal{E}_w = \varepsilon$.

Consider $\mathbf{G}$ as an F -group. Let $\mathbf{H}_1, \mathbf{H}_2$ be the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F . For $i = 1, 2$, we have $\mathbf{H}_i(F_w) = H_i$. We fix Tamagawa measures ([PR]) $dg = \otimes_v dg_v$ on $\mathbf{G}(\mathbb{A}_F)$, and $dh_i = \otimes_v dh_{i,v}$ on $\mathbf{H}_i(\mathbb{A}_F)$, such that dg_w and $dh_{i,w}$ coincide with the local Haar measures we have chosen for G and H_i , respectively.

Let 1_2 be the trivial representation of $\text{GL}(2, \mathbb{A}_F)$. Let $1_{2,E}$ be the trivial representation of $\text{GL}(2, \mathbb{A}_E)$. Let $\mathcal{Z}$ be a character of C_F such that $\mathcal{Z}^2 = 1$ and $\mathcal{Z}_w = \xi$. Let π_1 be the one dimensional representation $\mathcal{Z}1_2 \otimes 1$ of $\mathbf{H}_1(\mathbb{A}_F)$. Let π_2 be the one dimensional representation $(\mathcal{Z} \circ N_{E/F})1_{2,E} \otimes 1$ of $\mathbf{H}_2(\mathbb{A}_F)$. Let

$$\{\Pi\} = \bigotimes_{v \in V} \{\Pi_v^\times, \Pi_v^-\}$$

be the unstable quasi-packet $[\mathcal{Z}1_2, \mathcal{E}\mathcal{Z}1_2]$ of $\mathbf{G}(\mathbb{A}_F)$ (see Section 4.3.1). It lifts to the induced representation $I_{(2,2)}(\mathcal{Z}1_2, \mathcal{E}\mathcal{Z}1_2)$ of $\text{GL}(4, \mathbb{A}_F)$. From [F4, Sect. V. 10.6], $\Pi_v^\times = L(\varepsilon_v \nu_v, \varepsilon_v \rtimes \nu_v^{-1/2} \xi_v)$ for each place v , and

$$\Pi_v^- = \begin{cases} \delta^-(\varepsilon_v \nu_v^{1/2} \text{St}_{2,v}, \nu_v^{-1/2} \xi_v) & \text{if } \varepsilon_v \neq 1, \\ L(\nu_v^{1/2} \text{St}_{2,v}, \nu_v^{-1/2} \xi_v) & \text{if } \varepsilon_v = 1. \end{cases}$$

Here, ν_v is the normalized absolute value function on F_v , and $\text{St}_{2,v}$ denotes the Steinberg representation of $\text{GL}(2, F_v)$. In particular, $\Pi_u^\times = \pi_u^\times$ and $\Pi_u^- = \pi_u^-$.

Fix a place u of F which splits in E . Then, $\mathcal{E}_u$ is trivial, and

$$H_{1,u} = \mathbf{H}_1(F_u) = \{(g, a, b) \in \text{GL}(2, F_u) \times F_u^\times \times F_u^\times : \det g = ab\}.$$

We have $\pi_{1,u} = \mathcal{Z}_u 1_{2,u} \otimes 1$ (1 $\otimes$ 1). Here, $1_{2,u} := 1_{\text{GL}(2, F_u)}$, the trivial representation of $\text{GL}(2, F_u)$. The representations

$$\Pi_u^\times = L(\nu_u, 1 \rtimes \nu_u^{-1/2} \xi_u), \quad \Pi_u^- = L(\nu_u^{1/2} \text{St}_{2,u}, \nu_u^{-1/2} \xi_u)$$

are the two inequivalent nontempered subquotients of $1 \rtimes \mathcal{Z}_u 1_{2,u}$ (see [ST, Lemma 3.8]).

5.17. LEMMA. *For matching functions f_u on G_u and $f_{1,u}$ on $H_{1,u}$, the following holds:*

$$\langle \Pi_u^\times, f_u \rangle + \langle \Pi_u^-, f_u \rangle = \langle 1 \rtimes \mathcal{Z}_u 1_{2,u}, f_u \rangle = \langle \pi_{1,u}, f_{1,u} \rangle.$$

PROOF. The first equality is clear. The second follows from Lemma 5.12. □

The group of F_u -points of $\mathbf{H}_2$ is

$$H_{2,u} = (\text{GL}(2, F_u) \times \text{GL}(2, F_u) \times F_u^\times) / \{(aI_2, bI_2, (ab)^{-1}) : a, b \in F_u^\times\}.$$

For a character ω_u of the center Z_u of G_u , let $C(G_u, \omega_u)$ denote the space of smooth functions on G_u which are compactly supported modulo Z_u and transform under Z_u via ω_u^{-1} . If $\omega_u = 1$, then each $f_u \in C(G_u, 1)$ defines a test function $\bar{f}_u$ on $\mathrm{PGSp}(2, F_u)$.

The map $\mathrm{diag}(z, z, z, z) \mapsto (\mathrm{diag}(1, 1), \mathrm{diag}(1, 1), z)_*$, $z \in F_u^\times$, defines an isomorphism from Z_u to

$$Z'(H_{2,u}) := \{(\mathrm{diag}(1, 1), \mathrm{diag}(1, 1), z)_* : z \in F_u^\times\} \subset H_{2,u}.$$

Here, lower $*$ denotes image in $H_{2,u}$. For a character $\omega_{2,u}$ of the group $Z'(H_{2,u})$, let $C(H_{2,u}, \omega_{2,u})$ denote the space of smooth, compactly supported mod $Z'(H_{2,u})$ functions on $H_{2,u}$ which transform under $Z'(H_{2,u})$ via $\omega_{2,u}^{-1}$. If a function $f_{2,u}$ in $C(H_{2,u}, \omega_{2,u})$ matches some $f_u \in C(G_u, 1)$, then $\omega_{2,u}$ is necessarily trivial.

The map $(g, h, c)_* \mapsto (g, ch)$ defines an isomorphism from $H_{2,u}$ to the group

$$(\mathrm{GL}(2, F_u) \times \mathrm{GL}(2, F_u)) / \{(zI_2, z^{-1}I_2) : z \in F_u^\times\}.$$

Hence, the representations of $H_{2,u}$ with trivial central characters may be identified with the representations of $\mathrm{PGL}(2, F_u) \times \mathrm{PGL}(2, F_u)$, which is the group of F_u -points of the F -group $\mathbf{C}_0 = \mathrm{PGL}(2) \times \mathrm{PGL}(2)$. Moreover, if a function $f_{2,u}$ in $C(H_{2,u}, 1)$ matches a function f_u in $C(G_u, 1)$, then $f_{2,u}$ defines a function $\bar{f}_{2,u}$ on $C_{0,u}$ which matches $\bar{f}_u$ in the context of twisted endoscopy for $\mathrm{PGL}(4)$ (see [F4]).

5.18. LEMMA. *The following holds for matching functions $f_u \in C(G_u, 1)$ and $f_{2,u} \in C(H_{2,u}, 1)$:*

$$\langle \Pi_u^\times, f_u \rangle - \langle \Pi_u^-, f_u \rangle = \langle \mathcal{Z}_u 1_{2,u} \times \mathcal{Z}_u 1_{2,u}, \bar{f}_{2,u} \rangle = \langle \pi_{2,u}, f_{2,u} \rangle.$$

PROOF. The first equality follows from [F4, Prop. V. 8.6]. Let u_1, u_2 be the distinct places of E which lie above u . For $i = 1, 2$, we have $E_{u_i} = F_u$, $1_{2,E_{u_i}} = 1_{2,u}$, and $(\mathcal{Z} \circ \mathrm{N}_{E/F})_{u_i} = \mathcal{Z}_u$. Hence, $\pi_{2,u} = (\mathcal{Z}_u 1_{2,u} \times \mathcal{Z}_u 1_{2,u}) \otimes_2 1$, which may be identified with the representation $\mathcal{Z}_u 1_{2,u} \times \mathcal{Z}_u 1_{2,u}$ of $C_{0,u}$. The second equality follows. $\square$

In summary, the following local identities hold for matching functions:

$$(5.1) \quad \begin{aligned} \langle \Pi_u^\times, f_u \rangle + \langle \Pi_u^-, f_u \rangle &= \langle \pi_{1,u}, f_{1,u} \rangle, \\ \langle \Pi_u^\times, f_u \rangle - \langle \Pi_u^-, f_u \rangle &= \langle \pi_{2,u}, f_{2,u} \rangle. \end{aligned}$$

Let S' be the finite set of bad places for $\mathcal{Z}$, i.e. it is the union of $V - V^{\mathrm{un}}(E/F)$ and the set of places v where $\mathcal{Z}_v$ is ramified. In particular, the fixed places w, u lie outside of S' . For each place $v \notin S'$, $\pi_{1,v}$ and $\pi_{2,v}$ are unramified representations.

Fix two places $w_1, w_2 \notin S' \cup \{w\}$ which are prime in E . For a character ω of the center of $\mathbf{G}(\mathbb{A}_F)$, recall the definition of $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ (resp. $E(w_1, w_2, \mathbf{H}_i(\mathbb{A}_F), \omega)$): It denotes the space of functions in $C(\mathbf{G}(\mathbb{A}_F), \omega)$ (resp. $C(\mathbf{H}_i(\mathbb{A}_F), \omega)$) whose local components at w_1, w_2 are elliptic.

Let $S_0 = S' \cup \{w_1, w_2\}$. Let S be a finite set of places of F containing S_0 .

By Corollary 5.15, each representation Π' in $\{\Pi\}$ is $\mathcal{E}$ -invariant; hence, the twisted character $\langle \Pi', f \rangle_{\mathcal{E}}$ is equal to a product $\prod_v \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters. Here, $\langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ denotes the trace of $\Pi'_v(f_v) \rho(\mathcal{E})_v$, where $\rho(\mathcal{E})_v$ is the operator on the space of Π'_v restricted from the operator $\rho(\mathcal{E}) : \phi \mapsto \mathcal{E} \phi$ on $L^2(\mathbf{G}(F) \backslash \mathbf{G}(\mathbb{A}_F))$. Put $\langle \Pi', f \rangle_{\mathcal{E}, S} := \prod_{v \in S} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$.

Let $f = \otimes_v f_v$ be a function in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), 1)$ such that f_v is spherical for all $v \notin S$. For $i = 1, 2$, let $f_i = \otimes_v f_{i,v}$ be a function in $E(w_1, w_2, \mathbf{H}_i(\mathbb{A}_F), 1)$ which

matches f and has spherical local components at all $v \notin S$. The following $\mathcal{E}$ -trace identity follows from Proposition 4.17:

$$(5.2) \quad \sum_{\Pi' \in \{\Pi\}} m(\Pi') \langle \Pi', f \rangle_{\mathcal{E}, S} = \frac{1}{2} \langle \pi_1, f_1 \rangle_S + \frac{1}{2} \langle \pi_2, f_2 \rangle_S.$$

Here, $m(\Pi')$ is the multiplicity of Π' in the discrete spectrum of $\mathbf{G}$. From [F4, Sect. V. 10.6], we have:

$$m(\Pi') = \frac{1}{2} \left(1 + (-1)^{n(\Pi')} \right),$$

where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$. In other words, $\Pi' \in \{\Pi\}$ appears (with multiplicity one) in the discrete spectrum if and only if $\Pi'_v = \Pi_v^-$ for an even number of places v .

Let $P^+(S)$ denote the set of representations Π' in $\{\Pi\}$ such that Π'_v is unramified for all $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an *even* number of places v in S .

Let $P^-(S)$ denote the set of Π' in $\{\Pi\}$ such that Π'_v is unramified for all $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an *odd* number of places v in S .

By the multiplicity formula for $\{\Pi\}$, the equation (5.2) is equivalent to

$$(5.3) \quad 2 \sum_{\Pi' \in P^+(S)} \langle \Pi', f \rangle_{\mathcal{E}, S} = \langle \pi_1, f_1 \rangle_S + \langle \pi_2, f_2 \rangle_S.$$

5.19. CLAIM. *The following identities hold for matching functions:*

$$\begin{aligned} 2 \sum_{\Pi' \in P^+(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} &= \langle \pi_1, f_1 \rangle_{S_0} + \langle \pi_2, f_2 \rangle_{S_0}, \\ 2 \sum_{\Pi' \in P^-(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} &= \langle \pi_1, f_1 \rangle_{S_0} - \langle \pi_2, f_2 \rangle_{S_0}. \end{aligned}$$

PROOF. Recall that we have fixed a place u which splits in E . The identity (5.3), when applied to $S = S_0 \cup \{u\}$, is equivalent to

$$(5.4) \quad 2 \sum_{\Pi' \in P^+(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} \langle \Pi_u^\times, f_u \rangle + 2 \sum_{\Pi' \in P^-(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} \langle \Pi_u^-, f_u \rangle \\ = \langle \pi_1, f_1 \rangle_{S_0} \langle \pi_{1,u}, f_{1,u} \rangle + \langle \pi_2, f_2 \rangle_{S_0} \langle \pi_{2,u}, f_{2,u} \rangle.$$

By (5.1), the right-hand side of the above equation is equal to

$$(\langle \pi_1, f_1 \rangle_{S_0} + \langle \pi_2, f_2 \rangle_{S_0}) \langle \Pi_u^\times, f_u \rangle + (\langle \pi_1, f_1 \rangle_{S_0} - \langle \pi_2, f_2 \rangle_{S_0}) \langle \Pi_u^-, f_u \rangle.$$

Since the representations $\Pi_u^\times, \Pi_u^-$ are inequivalent, the claim follows from the linear independence of characters. $\square$

PROOF OF PROPOSITION 5.16. Recall that we have fixed a place w of F such that the local components of the global objects (fields, groups, representations, ..., etc.) at w coincide with the desired local objects over k . Applying (5.3) to $S = S_0 \cup \{w\}$, we obtain the following identity for matching functions whose local components at all $v \notin S$ are spherical:

$$(5.5) \quad 2 \sum_{\Pi' \in P^+(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} \langle \Pi_w^\times, f_w \rangle_{\mathcal{E}_w} + 2 \sum_{\Pi' \in P^-(S_0)} \langle \Pi', f \rangle_{\mathcal{E}, S_0} \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} \\ = \langle \pi_1, f_1 \rangle_{S_0} \langle \pi_{1,w}, f_{1,w} \rangle + \langle \pi_2, f_2 \rangle_{S_0} \langle \pi_{2,w}, f_{2,w} \rangle.$$

By Claim 5.19, the left-hand side of the above equation is equal to

$$\langle \pi_1, f_1 \rangle_{S_0} \langle \Pi_w^\times + \Pi_w^-, f_w \rangle_{\mathcal{E}_w} + \langle \pi_2, f_2 \rangle_{S_0} \langle \Pi_w^\times - \Pi_w^-, f_w \rangle_{\mathcal{E}_w}.$$

Rearranging terms, (5.5) is equivalent to

$$(5.6) \quad \langle \pi_1, f_1 \rangle_{S_0} \left[\langle \Pi_w^\times + \Pi_w^-, f_w \rangle_{\mathcal{E}_w} - \langle \pi_{1,w}, f_{1,w} \rangle \right] \\ = - \langle \pi_2, f_2 \rangle_{S_0} \left[\langle \Pi_w^\times - \Pi_w^-, f_w \rangle_{\mathcal{E}_w} - \langle \pi_{2,w}, f_{2,w} \rangle \right].$$

Pick a place $v \in S_0$ which does not split in E . Recall that an elliptic regular element in G_v is of type i ($i = 1, 2$) if its norms lie only in $\mathbf{H}_i(F_v)$. Choose f_v to be a function on G_v whose $\mathcal{E}_v$ -twisted orbital integral is supported only on the set of elliptic regular elements of type 1. We may then set $f_2 = 0$ without violating the matching condition on the test functions. For such choices of test functions, (5.6) becomes

$$\langle \pi_1, f_1 \rangle_{S_0} \left[\langle \Pi_w^\times + \Pi_w^-, f_w \rangle_{\mathcal{E}_w} - \langle \pi_{1,w}, f_{1,w} \rangle \right] = 0.$$

Since there exists $f_1 \in E(w_1, w_2, \mathbf{H}_1(\mathbb{A}_F), 1)$ such that $\langle \pi_1, f_1 \rangle_{S_0} \neq 0$, we obtain:

$$(5.7) \quad \langle \Pi_w^\times, f_w \rangle_{\mathcal{E}_w} + \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{1,w}, f_{1,w} \rangle.$$

We may likewise pick a place v in S_0 , and choose the function f , such that the twisted orbital integral of f_v is supported only on the elliptic regular set of type 2. It then follows that

$$(5.8) \quad \langle \Pi_w^\times, f_w \rangle_{\mathcal{E}_w} - \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{2,w}, f_{2,w} \rangle.$$

Let f, f_1, f_2 now be matching test functions on G, H_1, H_2 , respectively. For $* = \times, -$, we have by construction $\Pi_w^* = \pi^*$. Let $A^* \in \text{Hom}_G(\pi^*, \varepsilon \pi^*)$ be the intertwining operator defined by $\rho(\mathcal{E})_w$. By (5.7) and (5.8), we have:

$$\langle \pi^\times, f \rangle_{A^\times} + \langle \pi^-, f \rangle_{A^-} = \langle \xi 1_{2,k} \otimes 1, f_1 \rangle, \\ \langle \pi^\times, f \rangle_{A^\times} - \langle \pi^-, f \rangle_{A^-} = \langle (\xi \circ N_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes 2, f_2 \rangle.$$

If one of $A^\times, A^-$ is zero, then

$$\langle \xi 1_{2,k} \otimes 1, f_1 \rangle = \pm \langle (\xi \circ N_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes 2, f_2 \rangle.$$

Let f be an arbitrary test function on G whose ε -twisted orbital integral is supported only on the elliptic regular set of type 1. Let $f_2 = 0$. Then, $\langle \xi 1_{2,k} \otimes 1, f_1 \rangle$ is zero for each elliptic function f_1 on H_1 which matches f , which is a contradiction because $1_{2,k} \otimes 1$ is supported on elliptic elements. Thus, neither $A^\times$ nor A^- is trivial. The proposition follows. $\square$

REMARK. Our proof of Proposition 5.16 relies on the multiplicity formula deduced in [F4] for the global quasi-packet $[\mathcal{Z}1_2, \mathcal{E}\mathcal{Z}1_2]$. Said multiplicity formula is proven only for the case of trivial central character, hence the assumption that the character ξ in the hypothesis of the proposition satisfies $\xi^2 = 1$. If the same multiplicity formula holds in general, then it is clear that we may remove the condition $\xi^2 = 1$.

5.4.1.1. *Steinberg Representations.* Suppose the character ε of $k^\times$ is unramified, i.e. $\mathcal{K}/k$ is an unramified field extension.

Let $\nu, \nu_{\mathcal{K}}$ be the normalized absolute value functions on $k, \mathcal{K}$, respectively. In particular, since $\mathcal{K}/k$ is unramified, $\nu_{\mathcal{K}} = \nu \circ N_{\mathcal{K}/k}$. Recall that $\text{St}_{2,l}$ ($l = k, \mathcal{K}$) denotes the Steinberg square integrable representation of $\text{GL}(2, l)$.

Let ξ be a character of $k^\times$ with $\xi^2 = 1$. By [ST, Lemma 3.6], the induced representation $I = \varepsilon\nu \times \varepsilon \rtimes \nu^{-1/2}\xi$ of G is reducible, with 4 distinct subquotients:

$$I = \delta(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi) + L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi) \\ + L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi) + L(\nu\varepsilon, \varepsilon \rtimes \nu^{1/2}\xi).$$

Here, $\delta(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ is the unique square integrable subrepresentation of I . For $\zeta = \xi, \varepsilon\xi$, the representation $L(\nu^{1/2}\varepsilon\text{St}_{\text{GL}(2,k)}, \nu^{-1/2}\zeta)$ is the unique nontempered quotient of $\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\zeta$. The representation $L(\nu\varepsilon, \varepsilon \rtimes \nu^{-1/2}\xi)$ is a nontempered quotient of $\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\xi$.

Since the representation $L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ is the unique nontempered quotient of $\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\xi$, and $L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi)$ that of

$$\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\varepsilon\xi = \varepsilon \otimes (\nu^{1/2}\varepsilon\text{St}_{2,k} \rtimes \nu^{-1/2}\xi),$$

we have $L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi) \cong \varepsilon L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$. Since the subquotients of I are distinct, neither $L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ nor $L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi)$ is ε -invariant.

Recall that the representation $\delta(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$ belongs to a local packet of size two. The other member of this local packet is a cuspidal representation, denoted by $\delta^-(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$.

Let $\delta = \delta(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$, $\pi^- = \delta^-(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)$, and let $\pi^\times$ denote $L(\nu\varepsilon, \varepsilon \rtimes \nu^{-1/2}\xi)$.

5.20. LEMMA. *There exist nontrivial intertwining operators $A^\delta \in \text{Hom}_G(\delta, \varepsilon\delta)$, $A^- \in \text{Hom}_G(\pi^-, \varepsilon\pi^-)$, such that the following twisted character identities hold for matching functions:*

$$(5.9) \quad \begin{aligned} \langle \delta, f \rangle_{A^\delta} - \langle \pi^-, f \rangle_{A^-} &= \langle \xi \text{St}_{2,k} \otimes 1, f_1 \rangle, \\ \langle \delta, f \rangle_{A^\delta} + \langle \pi^-, f \rangle_{A^-} &= \langle (\xi \circ N_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes 2, f_2 \rangle. \end{aligned}$$

PROOF. The representation $I = \varepsilon\nu \times \varepsilon \rtimes \nu^{-1/2}\xi$ is equal to $\varepsilon\nu \rtimes I(\nu^{-1/2}\xi, \varepsilon\nu^{-1/2}\xi)$, and $I(\nu^{-1/2}\xi, \varepsilon\nu^{-1/2}\xi)$ is equal to the monomial representation $\pi((\nu^{-1/2}\xi) \circ N_{\mathcal{K}/k})$ associated with the character $(\nu^{-1/2}\xi) \circ N_{\mathcal{K}/k}$ of $\mathcal{K}^\times$. Thus,

$$I = \varepsilon\nu \rtimes \pi((\nu^{-1/2}\xi) \circ N_{\mathcal{K}/k}).$$

Let $A \in \text{Hom}_G(I, \varepsilon I)$ be the operator defined in Section 5.1.1. It intertwines each subquotient L of I to a representation which is equivalent to εL . The subquotients

$$L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi), \quad L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi)$$

are obtained from each other by tensoring with ε . Since they are inequivalent, A swaps their spaces, which implies that

$$\text{tr } L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\xi)(f)A = \text{tr } L(\nu^{1/2}\varepsilon\text{St}_{2,k}, \nu^{-1/2}\varepsilon\xi)(f)A = 0.$$

Since δ is the unique square integrable subrepresentation of I , the operator A maps the space of δ to itself. We have determined the image under A of three of the four subquotients of I ; hence, A must map the space of the unique irreducible quotient $\pi^\times$ of I to itself.

Let A^δ denote the restriction of A to δ . Let A' denote the operator on $\pi^\times$ induced from A . Then, A^δ, A' are nontrivial operators in $\text{Hom}_G(\delta, \varepsilon\delta), \text{Hom}_G(\pi^\times, \varepsilon\pi^\times)$, respectively. Moreover, we have:

$$(5.10) \quad \langle I, f \rangle_A = \langle \delta, f \rangle_{A'} + \langle \pi^\times, f \rangle_{A^\delta}.$$

For the ε -endoscopic groups, we have:

$$(5.11) \quad \begin{aligned} \xi I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 1 &= \xi 1_{2,k} \otimes_1 1 + \xi \text{St}_{2,k} \otimes_1 1, \\ (\xi \circ N_{\mathcal{K}/k}) I(\nu_{\mathcal{K}}^{1/2}, \nu_{\mathcal{K}}^{-1/2}) \otimes_2 1 &= (\xi \circ N_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes_2 1 + (\xi \circ N_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes_2 1. \end{aligned}$$

By Corollary 5.10, the following characters are equal to one another:

- $\langle \varepsilon\nu \times \varepsilon \rtimes \nu^{-1/2} \xi, f \rangle_{A'}$,
- $\langle \xi I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 1, f_1 \rangle$,
- $\langle B_{\mathcal{K}/k} (\xi I(\nu^{1/2}, \nu^{-1/2})) \otimes_2 1, f_2 \rangle = \langle (\xi \circ N_{\mathcal{K}/k}) I(\nu_{\mathcal{K}}^{1/2}, \nu_{\mathcal{K}}^{-1/2}) \otimes_2 1, f_2 \rangle$.

By Proposition 5.16, there exist nontrivial intertwining operators $A^\times, A^-$ such that

$$(5.12) \quad \begin{aligned} \langle \pi^\times, f \rangle_{A^\times} + \langle \pi^-, f \rangle_{A^-} &= \langle \xi 1_{2,k} \otimes_1 1, f_1 \rangle, \\ \langle \pi^\times, f \rangle_{A^\times} - \langle \pi^-, f \rangle_{A^-} &= \langle (\xi \circ N_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes_2 1, f_2 \rangle. \end{aligned}$$

Recall that $A^\times$ is restricted from a global operator on $L^2(\mathbf{G}(F) \backslash \mathbf{G}(\mathbb{A}_F))$ (see proof of Proposition 5.16). The representation $\pi^\times$ is irreducible. Since both A' and $A^\times$ have the property that their squares are the identity, by Schur's lemma we have $A^\times = d \cdot A'$, where $d = \pm 1$. It follows from (5.11), (5.12), and (5.10) that

$$\langle \xi \text{St}_{2,k} \otimes_1 1, f_1 \rangle = -d \langle \pi^\times, f \rangle_{A'} - \langle \pi^-, f \rangle_{A^-} + \langle \delta, f \rangle_{A^\delta} + \langle \pi^\times, f \rangle_{A'}.$$

The representation $\pi^\times$ is nontempered, while the rest of the representations in the above equation are square integrable. The central exponents of square integrable representations all decay ([Be, Prop. 29]). Invoking the linear independence of central exponents (see [FK, Sect. 21]), we conclude that $-d \langle \pi^\times, f \rangle_{A'} + \langle \pi^\times, f \rangle_{A'} = 0$. Hence, $d = 1$, and

$$\langle \xi \text{St}_{2,k} \otimes_1 1, f_1 \rangle = \langle \delta, f \rangle_{A^\delta} - \langle \pi^-, f \rangle_{A^-}.$$

Applying the same argument to $\langle (\xi \circ N_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes_2 1, f_2 \rangle$, we obtain:

$$\langle (\xi \circ N_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes_2 1, f_2 \rangle = \langle \delta, f \rangle_{A^\delta} + \langle \pi^-, f \rangle_{A^-}.$$

The lemma follows. $\square$

5.4.2. Cuspidal Representations. In this section, the character ε is not assumed to be unramified.

In what follows, we prove local character identities by constructing global objects whose local components at a chosen place coincide with specific local conditions. For instance, we construct automorphic representations whose components at a chosen finite place are the p -adic representations we are interested in. To do so, we make use of the following generalization of [F, Prop. III. 3]:

5.21. PROPOSITION. *Let F be a number field. Let $\mathbf{H}$ be a reductive connected F -group. Fix a nonarchimedean place w of F . Let τ_w be a cuspidal H_w -module. Let $\{v_i\}_{i \in I}$ be a finite set of nonarchimedean places. For each $i \in I$, let τ_{v_i} be a square integrable H_{v_i} -module. Let S be a finite set of places which contains the union of $w, \{v_i\}_{i \in I}$, and all the archimedean places. Then, there exists an automorphic representation π of $\mathbf{H}(\mathbb{A}_F)$ such that: (i) $\pi_w = \tau_w$; (ii) $\pi_{v_i} = \tau_{v_i}$ for all $i \in I$; (iii) π_v is unramified for all $v \notin S$.*

Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial representation of $\mathrm{GL}(2, k)$ with trivial central character ω_τ . Let $\{\pi^+, \pi^-\}$ be the local packet of $\mathrm{PGSp}(2, k)$ which lifts to the induced representation $I_{(2,2)}(\tau, \varepsilon\tau)$ of $\mathrm{PGL}(4, k)$, according to [F4, Prop. V. 5].

5.22. PROPOSITION. *There exist nontrivial operators A^+ in $\mathrm{Hom}_G(\pi^+, \varepsilon\pi^+)$, A^- in $\mathrm{Hom}_G(\pi^-, \varepsilon\pi^-)$, such that the following twisted character identities hold for matching functions $f \in C(G)$, $f_i \in C(H_i)$ ($i = 1, 2$):*

$$\begin{aligned}\langle \pi^+, f \rangle_{A^+} &= \frac{1}{2} \langle \tau \otimes_1 1, f_1 \rangle + \frac{1}{2} \langle B_{\mathcal{K}/k}\tau \otimes_2 \omega_\tau, f_2 \rangle, \\ \langle \pi^-, f \rangle_{A^-} &= \frac{1}{2} \langle \tau \otimes_1 1, f_1 \rangle - \frac{1}{2} \langle B_{\mathcal{K}/k}\tau \otimes_2 \omega_\tau, f_2 \rangle.\end{aligned}$$

REMARK. Note that by assumption ω_τ is equal to 1. We nonetheless write ω_τ in the above equations, for we expect the same character identities to hold in the cases where $\omega_\tau \neq 1$.

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F with the following properties:

- There exists a finite place w of F , prime in E , such that $F_w = k$.
- $E_w = \mathcal{K}$. Here, w denotes both the place of F and the unique place of E which lies above it.

Let V be the set of places of F . Let $V^{\mathrm{un}}(E/F) \subset V$ be the set of finite places which are unramified in E . Let $w_1, w_2 \in V$ be two (finite) places, different from w , which are unramified, prime in E .

Using Proposition 5.21, we construct a cuspidal automorphic representation $\mathcal{T}$ of $\mathrm{GL}(2, \mathbb{A}_F)$ with trivial central character such that:

- $\mathcal{T}_w = \tau$;
- $\mathcal{T}_{w_i}$ ($i = 1, 2$) is the Steinberg representation St_{2, w_i} of $\mathrm{GL}(2, F_{w_i})$;
- $\mathcal{T}_v$ is unramified for each finite place $v \notin \{w, w_1, w_2\}$.

In particular, since τ is non- $\mathcal{K}$ -monomial, $\mathcal{T}$ is non- E -monomial.

Let $\mathcal{E}$ be the quadratic character of C_F which corresponds to the extension E/F . In particular, $\mathcal{E}_w = \varepsilon$. View $\mathbf{G} = \mathrm{GSp}(2)$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let $\{\Pi\} = \otimes_v \{\Pi_v\}$ be the unstable packet $[\mathcal{T}, \mathcal{E}\mathcal{T}]$ of $\mathbf{G}(\mathbb{A}_F)$ (see Section 4.3.1). It lifts to the induced representation $I_{(2,2)}(\mathcal{T}, \mathcal{E}\mathcal{T})$ of $\mathrm{GL}(4, \mathbb{A}_F)$. For each place v where $\mathcal{T}_v$ is square integrable, the local packet $\{\Pi_v\}$ consists of two (square integrable) representations Π_v^+, Π_v^- . In particular, $\Pi_w^+ = \pi^+$ and $\Pi_w^- = \pi^-$.

At a place v where $\mathcal{T}_v$ is fully induced (for instance, where $\mathcal{T}_v$ is unramified), the packet $\{\Pi_v\}$ consists of a single representation, denoted by Π_v^+ . For convenience, we put $\Pi_v^- := 0$, and write $\{\Pi_v\} = \{\Pi_v^+, \Pi_v^-\}$.

At the place w_i ($i = 1, 2$), where $\mathcal{T}_{w_i} = \mathrm{St}_{2, w_i}$, the local packet $\{\Pi_{w_i}\}$ consists of the square integrable representation $\Pi_{w_i}^+ = \delta(\varepsilon_{w_i} \nu_{w_i}^{1/2} \mathrm{St}_{2, w_i}, \nu_{w_i}^{-1/2})$ and the cuspidal $\Pi_{w_i}^- = \delta^-(\varepsilon_{w_i} \nu_{w_i}^{1/2} \mathrm{St}_{2, w_i}, \nu_{w_i}^{-1/2})$.

By Corollary 5.15, each representation Π' in $\{\Pi\}$ is $\mathcal{E}$ -invariant. Hence, as in the case of one dimensional representations (Section 5.4.1), the twisted character $\langle \Pi', f \rangle_{\mathcal{E}}$ is equal to a product $\prod_v \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters.

Let π_1 be the automorphic representation $\mathcal{T} \otimes_1 1$ of $\mathbf{H}_1(\mathbb{A}_F)$. Let π_2 be the automorphic representation $B_{E/F}\mathcal{T} \otimes_2 1$ of $\mathbf{H}_2(\mathbb{A}_F)$.

Let $S_0 = V - V^{\text{un}}(E/F)$. Note that w may or may not lie in S_0 . Let $S_1 = S_0 \cup \{w\}$. Let $S = S_1 \cup \{w_1, w_2\}$. By Corollary 4.18, the following $\mathcal{E}$ -trace identity holds for matching functions f in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), 1)$ and f_i in $E(w_1, w_2, \mathbf{H}_i(\mathbb{A}_F), 1)$ ($i = 1, 2$) whose local components at all $v \notin S$ are spherical:

$$(5.13) \quad \sum_{\Pi' \in \{\Pi\}} m(\Pi') \langle \Pi', f \rangle_{\mathcal{E}, S} = \frac{1}{2} \langle \pi_1, f_1 \rangle_S + \frac{1}{2} \langle \pi_2, f_2 \rangle_S.$$

Here, $m(\Pi')$ is the multiplicity of Π' in the discrete spectrum of $\mathbf{G}$. From [F4, Sect. V. 10.3], we have:

$$m(\Pi') = \frac{1}{2} \left(1 + (-1)^{n(\Pi')} \right),$$

where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$. In other words, $\Pi' \in \{\Pi\}$ appears (with multiplicity one) in the discrete spectrum if and only if $\Pi'_v = \Pi_v^-$ for an even number of places v .

Let $P^+(S)$ denote the set of representations Π' in $\{\Pi\}$ such that Π'_v is unramified for all $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an *even* number of places v in S .

Let $P^-(S)$ denote the set of Π' in $\{\Pi\}$ such that Π'_v is unramified for all $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an *odd* number of places v in S . By the multiplicity formula, the equation (5.13) is equivalent to

$$(5.14) \quad 2 \sum_{\Pi' \in P^+(S)} \langle \Pi', f \rangle_{\mathcal{E}, S} = \langle \pi_1, f_1 \rangle_S + \langle \pi_2, f_2 \rangle_S.$$

If a place v of F is archimedean, then $\text{GL}(2, F_v)$ is equal to $\text{GL}(2, \mathbb{C})$ because F is totally imaginary. Hence, $\mathcal{T}_v$ is a fully induced representation.

If v is a finite place not in S , then by construction $\mathcal{T}_v$ is unramified. Since $\mathcal{T}$ is cuspidal, $\mathcal{T}_v$ is fully induced.

Hence, for all $v \in S_1 - \{w\}$, $\mathcal{T}_v$ is fully induced. Suppose $\mathcal{T}_v = I(\alpha_v, \beta_v)$ for some characters α_v, β_v of $F_v^\times$. Then,

	v prime in E	v splits in E
$\pi_{1,v} =$	$I(\alpha_v, \beta_v) \otimes_1 1$	$I(\alpha_v, \beta_v) \otimes_1 (1 \otimes 1)$
$\pi_{2,v} =$	$B_{E/F} I(\alpha_v, \beta_v) \otimes_2 (\alpha_v \beta_v) \circ N_{E_v/F_v}$	$(I(\alpha_v, \beta_v) \times I(\alpha_v, \beta_v)) \otimes_2 \alpha_v \beta_v$

Let Π''_v be the induced representation $\varepsilon_v \alpha_v / \beta_v \rtimes I(\beta_v, \beta_v \varepsilon_v)$ of G_v . By Propositions 5.6, 5.8, and Lemmas 5.12, 5.13, $\langle \pi_{1,v}, f_{1,v} \rangle$ and $\langle \pi_{2,v}, f_{2,v} \rangle$ are equal to $\langle \Pi''_v, f_v \rangle_{A_v}$. Here, A_v is the intertwining operator defined in Section 5.1.1 if v is prime in E ; otherwise, $A_v = 1$.

For each $v \in S_1 - \{w\}$, regard the local components of the test functions at all places $u \neq v$ as fixed. By (5.14) and the linear independence of characters, we conclude that $\{\Pi_v\}$ is the singleton consisting of Π''_v . Hence, the $(S_1 - \{w\})$ -components of both sides of (5.14) cancel one another.

Let $S_3 = S - (S_1 - \{w\}) = \{w, w_1, w_2\}$. After cancellation, (5.14) becomes

$$(5.15) \quad 2 \sum_{\Pi' \in P^+(S_3)} \epsilon(\Pi') \langle \Pi', f \rangle_{\mathcal{E}, S_3} = \prod_{v \in S_3} \langle \pi_{1,v}, f_{1,v} \rangle + \prod_{v \in S_3} \langle \pi_{2,v}, f_{2,v} \rangle.$$

Here, $\epsilon(\Pi') = \pm 1$. It appears in the equation because $\langle \Pi''_v, f_v \rangle_{A_v} = \pm \langle \Pi''_v, f_v \rangle_{\mathcal{E}_v}$ for $v \in S_1 - \{w\}$. Absorbing the signs $\epsilon(\Pi')$ into the local intertwining operators $\rho(\mathcal{E})_v$ at $v \in S_3$ if necessary, we assume that $\epsilon(\Pi') = 1$ for all Π' .

By construction, $\pi_{1,w_i} = \text{St}_{2,w_i} \otimes_1 1$ and $\pi_{2,w_i} = B_{E/F} \text{St}_{2,w_i} \otimes_2 1$ for $i = 1, 2$. Hence, by Lemma 5.20, there exist constants $\epsilon_i^+, \epsilon_i^- = \pm 1$ such that

$$\begin{aligned} \epsilon_i^+ \langle \Pi_{w_i}^+, f_{w_i} \rangle_{\mathcal{E}_{w_i}} - \epsilon_i^- \langle \Pi_{w_i}^-, f_{w_i} \rangle_{\mathcal{E}_{w_i}} &= \langle \pi_{1,w_i}, f_{1,w_i} \rangle, \\ \epsilon_i^+ \langle \Pi_{w_i}^+, f_{w_i} \rangle_{\mathcal{E}_{w_i}} + \epsilon_i^- \langle \Pi_{w_i}^-, f_{w_i} \rangle_{\mathcal{E}_{w_i}} &= \langle \pi_{2,w_i}, f_{2,w_i} \rangle. \end{aligned}$$

The equation (5.15) is therefore equivalent to

$$\begin{aligned} & \left(2 \sum_{\Pi' \in P^+(\{w, w_1\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w, w_1\}} \right) \langle \Pi_{w_2}^+, f_{w_2} \rangle_{\mathcal{E}_{w_2}} \\ & + \left(2 \sum_{\Pi' \in P^-(\{w, w_1\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w, w_1\}} \right) \langle \Pi_{w_2}^-, f_{w_2} \rangle_{\mathcal{E}_{w_2}} \\ & = \epsilon_2^+ \left(\prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle + \prod_{v \in \{w, w_1\}} \langle \pi_{2,v}, f_{2,v} \rangle \right) \langle \Pi_{w_2}^+, f_{w_2} \rangle_{\mathcal{E}_{w_2}} \\ & + \epsilon_2^- \left(\prod_{v \in \{w, w_1\}} \langle \pi_{2,v}, f_{2,v} \rangle - \prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle \right) \langle \Pi_{w_2}^-, f_{w_2} \rangle_{\mathcal{E}_{w_2}}. \end{aligned}$$

By the linear independence of characters, we have:

$$\begin{aligned} (5.16) \quad 2 \sum_{\Pi' \in P^+(\{w, w_1\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w, w_1\}} \\ = \epsilon_2^+ \left(\prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle + \prod_{v \in \{w, w_1\}} \langle \pi_{2,v}, f_{2,v} \rangle \right), \end{aligned}$$

$$\begin{aligned} (5.17) \quad 2 \sum_{\Pi' \in P^-(\{w, w_1\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w, w_1\}} \\ = \epsilon_2^- \left(\prod_{v \in \{w, w_1\}} \langle \pi_{2,v}, f_{2,v} \rangle - \prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle \right). \end{aligned}$$

Equation (5.16) has the same form as (5.15), with w_2 removed from the set $S_3 = \{w, w_1, w_2\}$. By applying the same argument inductively on the set $\{w_1, w_2\}$, we conclude that

$$2 \sum_{\Pi' \in P^+(\{w\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w\}} = \epsilon_1^+ \epsilon_2^+ (\langle \pi_{1,w}, f_{1,w} \rangle + \langle \pi_{2,w}, f_{2,w} \rangle).$$

Hence, there exists a constant $\epsilon^+ = \pm 1$ such that

$$(5.18) \quad 2 \langle \Pi_w^+, f_w \rangle_{\mathcal{E}_w} = 2 \sum_{\Pi' \in P^+(\{w\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w\}} = \epsilon^+ \langle \pi_{1,w}, f_{1,w} \rangle + \epsilon^+ \langle \pi_{2,w}, f_{2,w} \rangle.$$

Similarly, (5.17) implies that there exists $\epsilon_- = \pm 1$ such that

$$(5.19) \quad 2 \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = 2 \sum_{\Pi' \in P^-(\{w\})} \langle \Pi', f \rangle_{\mathcal{E}, \{w\}} = \epsilon^- \langle \pi_{1,w}, f_{1,w} \rangle - \epsilon^- \langle \pi_{2,w}, f_{2,w} \rangle.$$

For $* = +, -$, let $\rho(\mathcal{E})_w^*$ denote the operator in $\text{Hom}_{G_w}(\Pi_w^*, \mathcal{E}_w \Pi_w^*)$ restricted from $\rho(\mathcal{E})$. Let $A^* = \epsilon^* \cdot \rho(\mathcal{E})_w^*$. The proposition then follows from (5.18) and (5.19). $\square$

REMARK. As in Proposition 5.16, we may extend Proposition 5.22 to the case of nontrivial central characters if the multiplicity formula for the global packet $\{\Pi\}$ holds in general.

5.4.2.1. *Monomial Representations.* Let χ be a character of $\mathcal{K}^\times$. The representation $\pi = 1 \rtimes \pi(\chi)$ of G is reducible with two inequivalent tempered constituents π^+, π^- .

5.23. LEMMA. *There exist intertwining operators $A^* \in \text{Hom}_G(\pi^*, \varepsilon \pi^*)$ ($* = +, -$) such that the following holds for matching functions f in $C(G)$ and f_1 in $C(H_1)$:*

$$\langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} = \langle \pi(\chi) \otimes 1, f_1 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There is a place w of F such that $F_w = k$; (ii) w is prime in E , $E_w = \mathcal{K}$.

Let w_1, w_2 be two (finite) places, different from w , which are prime in E .

Let $\mathcal{E}$ be the quadratic character of C_E corresponding to the extension E/F . We identify the generator σ of $\text{Gal}(\mathcal{K}/k)$ with that of $\text{Gal}(E/F)$.

Construct a character $\mathcal{X}$ of C_E such that: (i) $\mathcal{X}_{w_i}$ ($i = 1, 2$) is not fixed by the action of σ ; (ii) $\mathcal{X}_v$ is unramified for each finite place $v \notin \{w_1, w_2\}$.

Let $\pi(\mathcal{X})$ be the cuspidal E -monomial representation of $\text{GL}(2, \mathbb{A}_F)$ associated with $\mathcal{X}$. At each finite place v where $\mathcal{X}_v$ is unramified, $\pi(\mathcal{X})_v = \pi(\mathcal{X}_v)$ is fully induced. Since F is totally imaginary, $\pi(\mathcal{X})_v$ is a fully induced representation of $\text{GL}(2, \mathbb{C})$ for each archimedean place v .

Let π_1 be the automorphic representation $\pi(\mathcal{X}) \otimes 1$ of $\mathbf{H}(\mathbb{A}_F)$. In particular, $\pi_{1,v}$ is parabolically induced for each $v \notin \{w_1, w_2\}$.

Let $\mathcal{T}$ be the following representation of the Heisenberg parabolic subgroup P of $\mathbf{G}(\mathbb{A}_F)$:

$$\mathcal{T} = 1 \otimes \pi(\mathcal{X}) : \begin{pmatrix} a & * & * \\ & g_2 & * \\ & & \frac{\det g_2}{a} \end{pmatrix} \mapsto \pi(\mathcal{X})(g_2).$$

Let $\Pi = 1 \rtimes \pi(\mathcal{X})$, the representation of $\mathbf{G}(\mathbb{A}_F)$ parabolically induced from $\mathcal{T}$. Since $\mathcal{E} \otimes (1 \rtimes \pi(\mathcal{X})) = 1 \rtimes \mathcal{E}\pi(\mathcal{X}) \cong 1 \rtimes \pi(\mathcal{X})$, the representation Π is $\mathcal{E}$ -invariant. At each place v where $\pi(\mathcal{X})_v$ is fully induced, Π_v is irreducible. If $\pi(\mathcal{X})_v$ is cuspidal, then Π_v has two inequivalent tempered constituents Π_v^+, Π_v^- .

The operator $I_{P,\mathcal{T}}(\mathcal{E})$, defined in Section 3.1.1, intertwines Π with $\mathcal{E}\Pi$. For each place v , let $I_{P,\mathcal{T}}(\mathcal{E})_v \in \text{Hom}_{G_v}(\Pi_v, \mathcal{E}_v \Pi_v)$ be the local component of $I_{P,\mathcal{T}}(\mathcal{E})$ at v . For an irreducible constituent Π' of Π , and a test function f on $\mathbf{G}(\mathbb{A}_F)$, put $\langle \Pi', f \rangle_{\mathcal{E}} := \text{tr } \Pi'(f) I_{P,\mathcal{T}}(\mathcal{E})$. It is equal to the product $\prod_{v \in V} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$, where $\langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} := \text{tr } \Pi'_v(f_v) I_{P,\mathcal{T}}(\mathcal{E})_v$.

Let S_0 be a finite set of places of F which contains the union of $\{w_1, w_2\}$, the archimedean places, and the finite places which are ramified in E . Let f, f_1 be matching functions on $\mathbf{G}(\mathbb{A}_F), \mathbf{H}_1(\mathbb{A}_F)$, with elliptic components at w_1, w_2 and spherical components at all the places outside of S_0 . By (4.6) in Section 4.5, we have:

$$\prod_{v \in S_0} (\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} - \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v}) = \prod_{v \in S_0} \langle \pi_{1,v}, f_{1,v} \rangle.$$

Since $\pi_{1,v}$ is fully induced for all $v \neq w_1, w_2$, by Proposition 5.6 and Lemma 5.12 the $(S_0 - \{w_1, w_2\})$ -components of the above equation cancel. Hence,

$$(5.20) \quad \prod_{v \in \{w_1, w_2\}} \left(\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} - \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v} \right) = \prod_{v \in \{w_1, w_2\}} \langle \pi_{1,v}, f_{1,v} \rangle.$$

Now, construct a character $\mathcal{X}'$ of C_E such that: (i) $\mathcal{X}'_w = \chi$; (ii) $\mathcal{X}'_{w_i} = \mathcal{X}_{w_i}$ for $i = 1, 2$; (iii) $\pi(\mathcal{X}')_v$ is fully induced for each $v \neq w_1, w_2$.

Let $\Pi = 1 \rtimes \pi(\mathcal{X}')$. Let $\pi_1 = \pi(\mathcal{X}') \otimes 1$. Let $S = S_0 \cup \{w\}$. By (4.6) and the usual cancellation, we have:

$$\prod_{v \in \{w_1, w_2, w\}} \left(\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} - \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v} \right) = \prod_{v \in \{w_1, w_2, w\}} \langle \pi_{1,v}, f_{1,v} \rangle.$$

By (5.20), the $\{w_1, w_2\}$ -components of the above equation cancel each other; hence,

$$\langle \Pi_w^+, f_w \rangle_{\mathcal{E}_w} - \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{1,w}, f_{1,w} \rangle.$$

Since Π_w^+, Π_w^- are inequivalent, $I_{P,T}(\mathcal{E})_w$ either swaps their spaces, or it maps each space to itself. Since the right-hand side of the above character identity is nonzero in general, we conclude that $I_{P,T}(\mathcal{E})_w$ restricts to $I_{P,T}(\mathcal{E})_w^+$ in $\text{Hom}_{G_w}(\Pi_w^+, \mathcal{E}_w \Pi_w^+)$ and $I_{P,T}(\mathcal{E})_w^-$ in $\text{Hom}_{G_w}(\Pi_w^-, \mathcal{E}_w \Pi_w^-)$.

Let $A^+ = I_{P,T}(\mathcal{E})_w^+$. Let $A^- = -I_{P,T}(\mathcal{E})_w^-$. The lemma follows. $\square$

Let θ, χ be two distinct characters of $\mathcal{K}^\times$ such that none of $\theta, \chi, \theta\chi, \theta^\sigma\chi$ is fixed by the generator σ of $\text{Gal}(\mathcal{K}/k)$.

Suppose $\theta\chi|_{k^\times} = \theta^\sigma\chi|_{k^\times} = \varepsilon$. Then, the central characters of $\pi(\theta\chi)$ and $\pi(\theta^\sigma\chi)$ are trivial. Let $\{\pi^+, \pi^-\}$ be the local packet of G which lifts to the induced representation $I_{(2,2)}(\pi(\theta\chi), \pi(\theta^\sigma\chi))$ of $\text{PGL}(4, k)$ according to [F4, Prop. V. 5].

5.24. PROPOSITION. *There exist operators $A^* \in \text{Hom}_G(\pi^*, \varepsilon\pi^*)$ ($*$ = +, -), and a constant $\epsilon = \pm 1$, such that the following character identities hold for matching functions:*

$$\begin{aligned} \langle \pi^+, f \rangle_{A^+} + \langle \pi^-, f \rangle_{A^-} &= \langle \pi(\chi) \otimes_1 \theta, f_1 \rangle, \\ \langle \pi^+, f \rangle_{A^+} - \langle \pi^-, f \rangle_{A^-} &= \epsilon \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle. \end{aligned}$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There is a place w of F such that $F_w = k$; (ii) w is prime in E , $E_w = \mathcal{K}$.

Let w_1, w_2 be two (finite) places, different from w , which are prime in E .

Let $\mathcal{E}$ be the quadratic character of C_E corresponding to the extension E/F . We identify the generator σ of $\text{Gal}(\mathcal{K}/k)$ with that of $\text{Gal}(E/F)$.

Construct characters $\mathcal{X}, \mathcal{O}$ of C_E such that: (i) $\mathcal{O}_w = \theta, \mathcal{X}_w = \chi$; (ii) $\mathcal{X}_{w_i}$ ($i = 1, 2$) is not fixed by the action of σ ; (iii) $\mathcal{O}_{w_i} = \mathcal{U}_{w_i} \circ N_{E/F}$ ($i = 1, 2$) for some character $\mathcal{U}_{w_i}$ of $F_{w_i}^\times$; (iv) $\mathcal{X}_v, \mathcal{O}_v$ are unramified for each finite place $v \neq w_1, w_2, w$.

Let $\{\Pi\}$ be the global unstable packet $[\pi(\mathcal{O}\mathcal{X}), \pi(\mathcal{O}^\sigma\mathcal{X})]$ of $\mathbf{G}(\mathbb{A}_F)$, described as follows:

- At a place v where $\pi(\mathcal{O}\mathcal{X})_v$ and $\pi(\mathcal{O}^\sigma\mathcal{X})_v$ are fully induced, $\{\Pi_v\}$ is the singleton consisting of an irreducible fully induced representation, denoted by Π_v^+ . For convenience, we put $\Pi_v^- := 0$ and write $\{\Pi_v\} = \{\Pi_v^+, \Pi_v^-\}$.
In particular, $\{\Pi_v\}$ is a singleton for each v which splits in E . Since F is totally imaginary, the archimedean places all split in E .
- At a place v , prime in E , such that $\mathcal{O}_v = \sigma\mathcal{O}_v$ but $\mathcal{X}_v \neq \sigma\mathcal{X}_v$, the local packet $\{\Pi_v\}$ consists of the two inequivalent tempered constituents Π_v^+, Π_v^- of $1 \rtimes \pi(\mathcal{O}_v\mathcal{X}_v)$.

- At a place v , prime in E , where none of $\mathcal{O}_v, \mathcal{X}_v, \mathcal{O}_v \mathcal{X}_v, \mathcal{O}_v {}^\sigma \mathcal{X}_v$ is fixed by σ , the local packet $\{\Pi_v\}$ consists of two inequivalent cuspidal representations Π_v^+, Π_v^- . In particular, $\{\Pi_w\} = \{\pi^+, \pi^-\}$.

From [F4, Sect. 10.3], a representation $\Pi' \in \{\Pi\}$ appears in the discrete spectrum of $\mathbf{G}$ with multiplicity

$$m(\Pi') = \frac{1}{2} \left(1 + (-1)^{n(\Pi')} \right),$$

where $n(\Pi')$ is the number of places v for which $\Pi'_v = \Pi_v^-$. In other words, Π' appears with multiplicity one in the discrete spectrum if and only if $\Pi'_v = \Pi_v^-$ for an even number of places v .

Let π_1 be the representation $\pi(\mathcal{X}) \otimes_1 \mathcal{O}$ of $\mathbf{H}(\mathbb{A}_F)$. Let $\tilde{\pi}_1 = \pi(\mathcal{O}) \otimes_1 \mathcal{X}$. By the construction of $\mathcal{X}$ and $\mathcal{O}$, we have $\pi_{1,w} = \pi(\chi) \otimes_1 \theta$ and $\tilde{\pi}_{1,w} = \pi(\theta) \otimes_1 \chi$. Moreover, the representations $\pi_{1,v}, \tilde{\pi}_{1,v}$ are parabolically induced for each place $v \neq w, w_1, w_2$.

For a finite set of places S , let $P^+(S)$ denote the set of representations Π' in $\{\Pi\}$ such that Π'_v is unramified for each $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an even number of places v in S .

Let $P^-(S)$ denote the set of Π' in $\{\Pi\}$ such that Π'_v is unramified for each $v \notin S$, and $\Pi'_v = \Pi_v^-$ for an odd number of places v in S .

Let S be the finite set of places $\{w, w_1, w_2\}$. Let f, f_1 be matching functions in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), 1)$, $E(w_1, w_2, \mathbf{H}_1(\mathbb{A}_F), 1)$, respectively, whose local components at all the places outside of S are spherical. By (4.3) and the usual cancellation, we obtain:

$$(5.21) \quad 2 \sum_{\Pi' \in P^+(S)} \langle \Pi', f \rangle_{\mathcal{E}, S} = \prod_{v \in S} \langle \pi_{1,v}, f_{1,v} \rangle + \prod_{v \in S} \langle \tilde{\pi}_{1,v}, f_{1,v} \rangle.$$

For $v = w_1, w_2$, we have by construction $\mathcal{O}_v = \mathcal{U}_v \circ N_{E/F}$ for some character $\mathcal{U}_v$ of $F_v^\times$. Hence,

$$\begin{aligned} \pi_{1,v} &= \pi(\mathcal{X}_v) \otimes_1 \mathcal{U}_v \circ N_{E/F} = \mathcal{U}_v \pi(\mathcal{X}_v) \otimes_1 1, \\ \tilde{\pi}_{1,v} &= \mathcal{U}_v I(1, \mathcal{E}_v) \otimes_1 \mathcal{X}_v. \end{aligned}$$

The local packet $\{\Pi_v\}$ consists of the inequivalent tempered constituents Π_v^+, Π_v^- of $1 \rtimes \mathcal{U}_v \pi(\mathcal{X}_v)$. By Lemma 5.23, we have:

$$(5.22) \quad \langle \pi_{1,v}, f_{1,v} \rangle = \epsilon_v^+ \langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} + \epsilon_v^- \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v}.$$

Here, $\epsilon_v^+, \epsilon_v^- = \pm 1$. They appear because each twisted character in Lemma 5.23 may differ from the corresponding character here by a sign. Since $f_{1,v}$ is elliptic and $\tilde{\pi}_{1,v}$ is parabolically induced, the term $\prod_{v \in S} \langle \tilde{\pi}_{1,v}, f_{1,v} \rangle$ in (5.21) vanishes.

Let v be still w_1 or w_2 . Let $\Pi_v = 1 \rtimes \pi(\mathcal{O}_v \mathcal{X}_v) = 1 \rtimes \mathcal{U}_v \pi(\mathcal{X}_v)$. Let A be the intertwining operator in $\text{Hom}_{G_v}(\Pi_v, \mathcal{E}_v \Pi_v)$ defined in Section 5.1.1. By Lemma 5.3, the twisted character $\langle \Pi_v, f_v \rangle_A$ is zero for all elliptic test function f_v on G_v . Since $\Pi_v = \Pi_v^+ + \Pi_v^-$, the twisted character $\langle \Pi_v^+, f_v \rangle_A$ must be equal to $-\langle \Pi_v^-, f_v \rangle_A$ for all elliptic f_v . Hence, there exists a constant $\xi_v = \pm 1$ such that $\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} = \xi_v \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v}$, and

$$\langle \pi_{1,v}, f_{1,v} \rangle = \epsilon_v^+ \langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} + \epsilon_v^- \xi_v \langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v}.$$

Since $\langle \pi_{1,v}, f_{1,v} \rangle \neq 0$ for a general elliptic test function $f_{1,v}$, the twisted character $\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v}$ is nonzero. Moreover, the sign ϵ_v^+ must be equal to $\epsilon_v^- \xi_v$. There is therefore a constant $\epsilon = \pm 1$ such that

$$\langle \pi_{1,v}, f_{1,v} \rangle = 2 \epsilon_v \langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v}.$$

We conclude that the equation (5.21) is equivalent to

$$(5.23) \quad 2 \langle \Pi_{w_2}^+, f_{w_2} \rangle_{\mathcal{E}_{w_2}} \cdot \left(\sum_{\Pi' \in P^+(\{w, w_1\})} \prod_{v \in \{w, w_1\}} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} + \xi_{w_2} \sum_{\Pi' \in P^-(\{w, w_1\})} \prod_{v \in \{w, w_1\}} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} \right) \\ = 2 \epsilon_{w_2} \langle \Pi_{w_2}^+, f_{w_2} \rangle_{\mathcal{E}_{w_2}} \cdot \prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle.$$

Hence,

$$\sum_{\Pi' \in P^+(\{w, w_1\})} \prod_{v \in \{w, w_1\}} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} + \xi_{w_2} \sum_{\Pi' \in P^-(\{w, w_1\})} \prod_{v \in \{w, w_1\}} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} \\ = \epsilon_{w_2} \prod_{v \in \{w, w_1\}} \langle \pi_{1,v}, f_{1,v} \rangle.$$

By repeating the same argument for the place w_1 , we conclude that there are constants $\xi^+, \xi^- = \pm 1$ such that

$$\xi^+ \langle \Pi_w^+, f_w \rangle_{\mathcal{E}_w} + \xi^- \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{1,w}, f_{1,w} \rangle.$$

By the symmetry between $\pi_{1,w} = \pi(\chi) \otimes_1 \theta$ and $\tilde{\pi}_{1,w} = \pi(\theta) \otimes_1 \chi$, we conclude that there are constants $\tilde{\xi}^+, \tilde{\xi}^- = \pm 1$ such that

$$\tilde{\xi}^+ \langle \Pi_w^+, f_w \rangle_{\mathcal{E}_w} + \tilde{\xi}^- \langle \Pi_w^-, f_w \rangle_{\mathcal{E}_w} = \langle \tilde{\pi}_{1,w}, f_{1,w} \rangle.$$

Since $\pi_{1,w} \not\cong \tilde{\pi}_{1,w}$, the ordered couple (ξ^+, ξ^-) is equal to neither $(\tilde{\xi}^+, \tilde{\xi}^-)$ nor $(-\tilde{\xi}^+, -\tilde{\xi}^-)$, or else we would have $\langle \pi_{1,w}, f_{1,w} \rangle = \pm \langle \tilde{\pi}_{1,w}, f_{1,w} \rangle$.

For $*$ = $+/-$, let $\rho(\mathcal{E})_w^*$ be the operator in $\text{Hom}(\Pi_w^*, \mathcal{E}_w \Pi_w^*)$ restricted from $\rho(\mathcal{E})$. The proposition now follows on letting $A^* = \xi^* \rho(\mathcal{E})_w^*$ and $\epsilon = \xi^+ / \tilde{\xi}^+$. $\square$

5.5. Character Identities for Stable Packets

5.5.1. Lifting from H_1 . Let $\mathcal{K}/k$ be the quadratic field extension corresponding to ε via local class field theory.

Let χ be a character of $\mathcal{K}^\times$ such that $\chi \neq \sigma\chi$. From [ST, Prop. 4.8], the induced (local) representation $\nu\varepsilon \rtimes \nu^{-1/2}\pi(\chi)$ of G is reducible of length 2. Its constituents consist of a nontempered quotient, denoted by $L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$, and a square integrable subrepresentation, which we denote here by $\delta(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$.

5.25. LEMMA. *Let L denote the local representation $L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$ of G . There exists an intertwining operator $A \in \text{Hom}_G(L, \varepsilon L)$, and a nonzero constant c , such that the following identity holds for matching functions $f \in C(G)$, $f_1 \in C(H_1)$:*

$$\langle L, f \rangle_A = c \langle 1_{2,k} \otimes_1 \chi, f_1 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There exists a finite place w of F such that $F_w = k$, (ii) w is prime in E , (iii) $E_w = \mathcal{K}$. We identify the generator σ of $\text{Gal}(\mathcal{K}/k)$ with that of $\text{Gal}(E/F)$.

Let V be the set of places of F . Let $V^{\text{un}}(E/F) \subset V$ be the set of finite places which are unramified in E .

Let $\mathcal{E}$ be the quadratic character of C_F corresponding to the extension E/F . In particular, $\mathcal{E}_w = \varepsilon$. We consider $\mathbf{G}$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Construct a character $\mathcal{X}$ of C_E with $\mathcal{X}_w = \chi$.

Let $\{\Pi\} = \bigotimes_v \{\Pi_v\}$ be the stable quasi-packet

$$\{L(\nu\mathcal{E}, \nu^{-1/2}\pi(\mathcal{X}))\}$$

of $\mathbf{G}(\mathbb{A}_F)$ (see Section 4.3.1, [F4, Prop. V. 10.10]). It lifts to the Langlands quotient representation $J(\nu^{1/2}\pi(\mathcal{X}), \nu^{-1/2}\pi(\mathcal{X}))$ of $\mathrm{GL}(4, \mathbb{A}_F)$. Here, by abuse of notation we let ν denote also the normalized absolute value function on $\mathbb{A}_F$. The local components of $\{\Pi\}$ are as follows:

- If $\mathcal{E}_v \neq 1$ and $\mathcal{X}_v \neq {}^\sigma\mathcal{X}_v$, then the local quasi-packet $\{\Pi_v\}$ consists of

$$L(\nu_v\mathcal{E}_v, \nu_v^{-1/2}\pi(\mathcal{X}_v)),$$

the unique nontempered quotient of $\nu_v\mathcal{E}_v \rtimes \nu_v^{-1/2}\pi(\mathcal{X}_v)$. In particular, $\{\Pi_v\}$ is the singleton consisting of L . Since $L(\nu_v\mathcal{E}_v, \nu_v^{-1/2}\pi(\mathcal{X}_v))$ is the unique nontempered quotient of the $\mathcal{E}_v$ -invariant $\nu_v\mathcal{E}_v \rtimes \nu_v^{-1/2}\pi(\mathcal{X}_v)$, it is $\mathcal{E}_v$ -invariant.

- If $\mathcal{E}_v \neq 1$, and $\mathcal{X}_v = \mathcal{U}_v \circ \mathrm{N}_{E/F}$ for some character $\mathcal{U}_v$ of C_F , then $\{\Pi_v\}$ is the quasi-packet

$$\{L(\nu_v\mathcal{E}_v, \mathcal{E}_v \rtimes \mathcal{U}_v\nu_v^{-1/2}), \delta^-(\mathcal{E}_v\nu_v^{1/2}\mathrm{St}_{2,v}, \nu_v^{-1/2}\mathcal{U}_v)\}.$$

By Lemma 5.14, each member of $\{\Pi_v\}$ is $\mathcal{E}_v$ -invariant.

- If v splits into two places v_1, v_2 of E , then $\{\Pi_v\}$ is the singleton consisting of the irreducible representation $\mathcal{X}_1/\mathcal{X}_2 \rtimes \mathcal{X}_2 1_{2, F_v}$. Here, $\mathcal{X}_i := \mathcal{X}_{v_i}$ ($i = 1, 2$), and $1_{2, F_v}$ is the trivial representation of $\mathrm{GL}(2, F_v)$.

It follows from the above remarks that each member of $\{\Pi\}$ is $\mathcal{E}$ -invariant. Hence, for each $\Pi' \in \{\Pi\}$, $\langle \Pi', f \rangle_{\mathcal{E}}$ is the product $\prod_{v \in V} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters. Here, $\langle \Pi'_v, f_v \rangle_{\mathcal{E}_v} := \mathrm{tr} \Pi'_v(f_v) \rho(\mathcal{E})_v$, where $\rho(\mathcal{E})_v$ is the local intertwining operator in $\mathrm{Hom}_{G_v}(\Pi'_v, \mathcal{E}_v \Pi'_v)$ restricted from $\rho(\mathcal{E})$.

Let w_1, w_2 be two (finite) places, different from w , which are prime in E .

Let S_0 be the (finite) set of bad places for $\mathcal{X}$. That is, it is the union of $V - V^{\mathrm{un}}(E/F)$ and the places $v \in V^{\mathrm{un}}(E/F)$ for which $\mathcal{X}_v$ is ramified. In particular, since $\mathcal{X}_w \neq {}^\sigma\mathcal{X}_w$, the place w lies in S_0 . Let $S = S_0 \cup \{w_1, w_2\}$.

The split central character of π_1 is $\mathcal{X}|_{C_F}$. Let $f \in E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \mathcal{X}|_{C_F})$, $f_1 \in E(w_1, w_2, \mathbf{H}_1(\mathbb{A}_F), \mathcal{X}|_{C_F})$ be matching functions whose local components at the places outside of S are spherical.

Since $\{\Pi\}$ is stable, each representation $\Pi' \in \{\Pi\}$ occurs with multiplicity one in the discrete spectrum of $\mathbf{G}$. Here, we are assuming that the multiplicity one property holds for $\mathrm{GSp}(2)$.

By Claim 4.34 and the stability of $\{\Pi\}$, the following twisted trace identity holds:

$$(5.1) \quad \sum_{\Pi' \in \{\Pi\}} \langle \Pi', f \rangle_{\mathcal{E}, S} = \frac{1}{2} \langle 1_2 \otimes 1 \mathcal{X}, f_1 \rangle_S + \frac{1}{2} \langle 1_2 \otimes 1 {}^\sigma\mathcal{X}, f_1 \rangle_S.$$

By the matching condition on the functions f and f_1 , the terms $\langle 1_2 \otimes 1 \mathcal{X}, f_1 \rangle_S$ and $\langle 1_2 \otimes 1 {}^\sigma\mathcal{X}, f_1 \rangle_S$ have equal contribution. For each $\Pi' \in \{\Pi\}$, Π'_w is equal to $L =$

$L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$. Hence, (5.1) is equivalent to

$$\left(\sum_{\Pi' \in \{\Pi\}} \langle \Pi', f \rangle_{\mathcal{E}, S - \{w\}} \right) \cdot \epsilon(\Pi') \cdot \langle L, f_w \rangle_{\mathcal{E}_w} = \prod_{v \in S} \langle 1_{2, F_v} \otimes_1 \mathcal{X}_v, f_{1, v} \rangle.$$

Here, $\epsilon(\Pi') = \pm 1$. It appears in the equation because, for two distinct members Π' , Π'' of $\{\Pi\}$, the local intertwining operators $\rho(\mathcal{E})_w$ for L as a component of Π' and for L as a component of Π'' may differ by a sign.

Fixing the local components of the test functions at each place in $S - \{w\}$, the lemma follows. $\square$

Let χ be a character of $\mathcal{K}^\times$ such that $\chi \neq \sigma\chi$. Let $L = L(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$. Let $\delta = \delta(\nu\varepsilon, \nu^{-1/2}\pi(\chi))$.

5.26. PROPOSITION. *There exist intertwining operators A^L in $\text{Hom}_G(L, \varepsilon L)$, and A^δ in $\text{Hom}_G(\delta, \varepsilon\delta)$, such that the following identities hold for matching functions:*

$$\begin{aligned} \langle L, f \rangle_{A^L} &= \langle 1_{2, k} \otimes_1 \chi, f_1 \rangle, \\ \langle \delta, f \rangle_{A^\delta} &= \langle \text{St}_{2, k} \otimes_1 \chi, f_1 \rangle. \end{aligned}$$

PROOF. Let $I = \nu\varepsilon \rtimes \nu^{-1/2}\pi(\chi)$. Thus, $I = \delta + L$.

Let A be the operator in $\text{Hom}_G(I, \varepsilon I)$ defined in Section 5.1.1. By Proposition 5.6, we have:

$$\langle I, f \rangle_A = \langle I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 \chi, f_1 \rangle.$$

Since δ is the unique subrepresentation of I , and L is the unique quotient, the operator A induces operators $A|_\delta$ in $\text{Hom}_G(\delta, \varepsilon\delta)$ and $A|_L$ in $\text{Hom}_G(L, \varepsilon L)$. We have therefore:

$$(5.2) \quad \langle I, f \rangle_A = \langle \delta, f \rangle_{A|_\delta} + \langle L, f \rangle_{A|_L} = \langle I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 \chi, f_1 \rangle.$$

By Lemma 5.25, there exist a constant c and an operator $A^L \in \text{Hom}_G(L, \varepsilon L)$ such that

$$(5.3) \quad \langle L, f \rangle_{A^L} = c \langle 1_{2, k} \otimes_1 \chi, f_1 \rangle.$$

The operator A^L differs from $A|_L$ by at most a sign. Absorbing any negative sign into the constant c if necessary, we may assume $A^L = A|_L$.

Since

$$I(\nu^{1/2}, \nu^{-1/2}) \otimes_1 \chi = 1_{2, k} \otimes_1 \chi + \text{St}_{2, k} \otimes_1 \chi,$$

by (5.2) and (5.3) we have:

$$\langle \delta, f \rangle_{A|_\delta} - \langle \text{St}_{2, k} \otimes_1 \chi, f_1 \rangle = (1 - c) \langle 1_{2, k} \otimes_1 \chi, f_1 \rangle.$$

The right-hand side of the above equation consists of a nontempered representation, while the representations which appear on the left are square integrable. The central exponents of a square integrable representation all decay. By the linear independence of central exponents (see [FK, Sect. 21]), we conclude that $c = 1$.

Hence,

$$\langle L, f \rangle_{A^L} = \langle 1_{2, k} \otimes_1 \chi, f_1 \rangle, \quad \langle \delta, f \rangle_{A|_\delta} = \langle \text{St}_{2, k} \otimes_1 \chi, f_1 \rangle.$$

Let $A^\delta = A|_\delta$. The proposition follows. $\square$

5.27. PROPOSITION. *Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial, admissible representation of $\mathrm{GL}(2, k)$, χ be a character of $\mathcal{K}^\times$, such that: There does not exist a character ε' of $k^\times$ such that ${}^\sigma\chi/\chi = \varepsilon' \circ \mathrm{N}_{\mathcal{K}/k}$ and $\tau \cong \varepsilon'\tau$. There exist a collection $[\pi]$ of distinct ε -invariant, cuspidal, admissible representations of G , and an operator $A(\pi')$ in $\mathrm{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:*

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes_1 \chi, f_1 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There exists a place w of F such that $F_w = k$; (ii) w is prime in E , with $E_w = \mathcal{K}$. We identify the generator σ of $\mathrm{Gal}(\mathcal{K}/k)$ with that of $\mathrm{Gal}(E/F)$.

Let $\mathcal{E}$ be the quadratic character of C_F corresponding to E/F . Thus, $\mathcal{E}_w = \varepsilon$. We view $\mathbf{G}$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let w_1, w_2 be two (finite) places, different from w , which are prime in E .

Construct a character $\mathcal{X}$ of C_E such that:

- $\mathcal{X}_w = \chi$;
- $\mathcal{X}_{w_i} \neq {}^\sigma\mathcal{X}_{w_i}$ ($i = 1, 2$);
- $\mathcal{X}_v$ is unramified for each finite place $v \neq w, w_1, w_2$.

Construct a cuspidal automorphic representation $\mathcal{T}$ of $\mathrm{GL}(2, \mathbb{A}_F)$ such that:

- $\mathcal{T}_w = \tau$;
- $\mathcal{T}_{w_i} = \mathrm{St}_{2, w_i}$ ($i = 1, 2$);
- $\mathcal{T}_v$ is unramified for each finite place $v \neq w, w_1, w_2$.

In particular, since τ is not $\mathcal{K}$ -monomial, $\mathcal{T}$ is not E -monomial.

Let π_1 be the automorphic representation $\mathcal{T} \otimes_1 \mathcal{X}$ of $\mathbf{H}_1(\mathbb{A}_F)$. Let ${}^\sigma\pi_1$ be $\mathcal{T} \otimes_1 {}^\sigma\mathcal{X}$.

Let V be the set of places of F .

Let $\{\Pi\} = \bigotimes_{v \in V} \{\Pi_v\}$ be the global packet of $\mathbf{G}(\mathbb{A}_F)$ whose global datum is the lift of the global datum parametrizing π_1 (see Section 4.1.1.1). By Lemma 4.20, $\{\Pi\}$ is a stable packet. Hence, each representation $\Pi' \in \{\Pi\}$ occurs with multiplicity one in the discrete spectrum of $\mathbf{G}$. Here, we are assuming that $\mathrm{GSp}(2)$ has the multiplicity one property.

Let $\{\Pi\}^\mathcal{E}$ denote the set of $\mathcal{E}$ -invariant representations in $\{\Pi\}$. For each $\Pi' \in \{\Pi\}^\mathcal{E}$, the character $\langle \Pi', f \rangle_\mathcal{E}$ is a product $\prod_{v \in V} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters.

For a finite set of places S , let $\{\Pi\}_S^\mathcal{E}$ denote the set of representations in $\{\Pi\}^\mathcal{E}$ which have the same unramified local component Π_v^0 at each finite place $v \notin S$.

Let $V^{\mathrm{un}}(E/F) \subset V$ be the set of finite places which are unramified in E . Let $S_0 = V - V^{\mathrm{un}}(E/F)$. Let $S = S_0 \cup \{w, w_1, w_2\}$. Then, $\pi_{1,v}$ is unramified for each place $v \notin S$.

Let ω be the split central character of π_1 . Let f and f_1 be matching functions in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and $E(w_1, w_2, \mathbf{H}_1(\mathbb{A}_F), \omega)$, respectively, such that their local components at the finite places outside of S are spherical.

By Proposition 4.19 and the stability of $\{\Pi\}$, the following twisted trace identity holds:

$$(5.4) \quad \sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \langle \Pi', f \rangle_{\mathcal{E}, S} = \frac{1}{2} \langle \pi_1, f_1 \rangle_S + \frac{1}{2} \langle {}^\sigma\pi_1, f_1 \rangle_S.$$

The matching condition on f and f_1 implies that $\langle \pi_1, f_1 \rangle_S$ and $\langle {}^\sigma \pi_1, f_1 \rangle_S$ have equal contribution to the trace identity. Consequently, (5.4) is equivalent to

$$(5.5) \quad \sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \langle \Pi', f \rangle_{\mathcal{E}, S} = \langle \pi_1, f_1 \rangle_S = \prod_{v \in S} \langle \pi_{1,v}, f_{1,v} \rangle.$$

Since F is totally imaginary, and $\mathcal{T}_v$ is unramified for each finite place $v \neq w, w_1, w_2$, the representation $\pi_{1,v}$ is parabolically induced for each place v in $S - \{w, w_1, w_2\}$. For $i = 1, 2$, the representation π_{1,w_i} is equal to $\text{St}_{2,w_i} \otimes_1 \mathcal{X}_{w_i}$. By Propositions 5.6, 5.26, Lemma 5.12, and the linear independence of characters, the local components at each place $v \in S - \{w\}$ of the representations in $\{\Pi\}_S^\mathcal{E}$ are all equal to the same representation $\Pi_{0,v}$.

Hence, the $(S - \{w\})$ -components of (5.5) cancel up to a sign. That is, there exists a set $\{\epsilon(\Pi') = \pm 1 : \Pi' \in \{\Pi\}_S^\mathcal{E}\}$ such that

$$\sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \epsilon(\Pi') \langle \Pi'_w, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{1,w}, f_{1,w} \rangle.$$

The signs $\epsilon(\Pi')$ appear because, for a place $v \in S - \{w, w_1, w_2\}$ (resp. $v = w_1, w_2$), the twisted character in Proposition 5.6 (resp. Proposition 5.26) may differ from $\langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ by a sign.

For $\Pi' \in \{\Pi\}_S^\mathcal{E}$, let $\rho(\mathcal{E})|_{\Pi'_w}$ denote the operator in $\text{Hom}_{G_w}(\Pi'_w, \mathcal{E}_w \Pi'_w)$ which is defined by the restriction of $\rho(\mathcal{E})$ to Π' . Let

$$[\pi] = \{\Pi'_w : \Pi' \in \{\Pi\}_S^\mathcal{E}\}.$$

Since the local components at each place $v \neq w$ of the representations in $\{\Pi\}_S^\mathcal{E}$ are all equal to the same representation, the elements in $[\pi]$ are distinct.

Since $\pi_{1,w}$ is cuspidal, its central exponents are equal to zero. Hence, by the linear independence of central exponents ([FK, Sect. 21]), every representation in $[\pi]$ is cuspidal. For each $\pi' = \Pi'_w$ in $[\pi]$, let $A(\pi') = \epsilon(\Pi') \cdot \rho(\mathcal{E})|_{\Pi'_w}$. The proposition follows. $\square$

5.5.1.1. Some Additional Twisted Character Identities for H_1 . Using the global character identities listed in Section 4.5, we list the following local character identities. We skip their proofs, which are no different in nature from what we have encountered so far. For an ε -invariant representation π of G , we let A be the (unique up to a sign) nontrivial intertwining operator in $\text{Hom}_G(\pi, \varepsilon\pi)$.

- (1) Let θ, χ be characters of $\mathcal{K}^\times$ such that $\theta\chi = \mu \circ N_{\mathcal{K}/k}$ for some character μ of $k^\times$.

Suppose ${}^\sigma \chi / \chi$ does not factor through $N_{\mathcal{K}/k}$. Then, $\pi({}^\sigma \chi / \chi)$ is cuspidal, and the following holds for matching functions:

$$\left\langle \pi \left(\frac{{}^\sigma \chi}{\chi} \right) \rtimes \mu, f \right\rangle_A = \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle.$$

If ${}^\sigma \chi / \chi = \varepsilon' \circ N_{\mathcal{K}/k}$ for some nontrivial quadratic character ε' of $k^\times$, then the following holds for matching functions:

$$\langle \varepsilon' \times \varepsilon' \varepsilon \rtimes \mu, f \rangle_A = \langle \pi(\theta) \otimes_1 \chi, f_1 \rangle.$$

- (2) Let $\mathcal{K}' \neq \mathcal{K}$ be a quadratic extension of k . Let ε' be the quadratic character of $k^\times$ corresponding to $\mathcal{K}'/k$ via local class field theory.

Let χ be a character of $\mathcal{K}^\times$ such that ${}^\sigma \chi / \chi = \varepsilon' \circ N_{\mathcal{K}/k}$. In particular, $\chi \neq {}^\sigma \chi$.

Let θ be a character of $\mathcal{K}'^\times$ which is not fixed by the action of $\text{Gal}(\mathcal{K}'/k)$. Let μ be a character of $\mathcal{K}'^\times$ which satisfies $\mu \circ N_{\mathcal{K}\mathcal{K}'/\mathcal{K}'} = \chi \circ N_{\mathcal{K}\mathcal{K}'/\mathcal{K}}$. Here, $\mathcal{K}\mathcal{K}'$ is the compositum of $\mathcal{K}$ and $\mathcal{K}'$.

Let $\pi_{\mathcal{K}'}(\theta)$, $\pi_{\mathcal{K}'}(\mu\theta)$ be the $\mathcal{K}'$ -monomial representations associated with θ , $\mu\theta$, respectively.

The following holds for matching functions:

$$\langle \varepsilon \varepsilon' \rtimes \pi_{\mathcal{K}'}(\mu\theta), f \rangle_A = \langle \pi_{\mathcal{K}'}(\theta) \otimes_1 \chi, f_1 \rangle.$$

5.5.2. Lifting from H_2 . Let $\nu, \nu_{\mathcal{K}}$ be the normalized absolute value functions on the p -adic fields $k, \mathcal{K}$, respectively. Let $1_{2,l}$ ($l = k, \mathcal{K}$) denote the trivial representation of $\text{GL}(2, l)$. Let $\text{St}_{2,l}$ denote the Steinberg representation of $\text{GL}(2, l)$.

For a nontrivial quadratic character ζ of $k^\times$, and a cuspidal ζ -invariant representation τ of $\text{GL}(2, k)$, let $L = L(\zeta\nu, \nu^{-1/2}\tau)$ denote the unique nontempered quotient of the induced representation $\zeta\nu \rtimes \nu^{-1/2}\tau$ of G . Let $\delta = \delta(\zeta\nu, \nu^{-1/2}\tau)$ denote the unique square integrable subrepresentation of $\zeta\nu \rtimes \nu^{-1/2}\tau$.

Let χ be a character of $\mathcal{K}^\times$ such that $\chi \neq {}^\sigma\chi$ and ${}^\sigma\chi/\chi = \varepsilon' \circ N_{\mathcal{K}/k}$ for some nontrivial quadratic character ε' of $k^\times$. Thus, the ε -invariant representation $\pi(\chi)$ is also ε' -invariant ([LL]).

5.28. LEMMA. *Let $\zeta = \varepsilon'$ or $\varepsilon'\varepsilon$. There exists an operator A in $\text{Hom}_G(L, \varepsilon L)$, and a nonzero constant c , such that the following holds for matching functions $f \in C(G)$, $f_2 \in C(H_2)$:*

$$\langle L, f \rangle_A = c \langle \chi 1_{2,\mathcal{K}} \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \rangle.$$

PROOF. This is due to Claim 4.35. The proof is otherwise identical with that of Lemma 5.25, and we skip it. $\square$

5.29. PROPOSITION. *Let $\zeta = \varepsilon'$ or $\varepsilon'\varepsilon$. If the extension $\mathcal{K}/k$ is unramified, then there exist operators A^L in $\text{Hom}_G(L, \varepsilon L)$, and A^δ in $\text{Hom}_G(\delta, \varepsilon\delta)$, such that the following identities hold for matching functions:*

$$\begin{aligned} \langle L, f \rangle_{A^L} &= \langle \chi 1_{2,\mathcal{K}} \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \rangle, \\ \langle \delta, f \rangle_{A^\delta} &= \langle \chi \text{St}_{2,\mathcal{K}} \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \rangle. \end{aligned}$$

PROOF. Since $\mathcal{K}/k$ is unramified, $\nu_{\mathcal{K}}$ is equal to $\nu \circ N_{\mathcal{K}/k}$. By Proposition 5.8, we have:

$$\begin{aligned} \left\langle \varepsilon \zeta \nu \rtimes \nu^{-1/2} \pi(\chi), f \right\rangle_A \\ = \langle L, f \rangle_A + \langle \delta, f \rangle_A = \left\langle \chi I(\nu_{\mathcal{K}}^{1/2}, \nu_{\mathcal{K}}^{-1/2}) \otimes_2 \chi|_{k^\times} \cdot \zeta, f_2 \right\rangle. \end{aligned}$$

Here, A is the intertwining operator defined in Section 5.1.1. Using Lemma 5.28, the proposition follows from the same argument used in the proof of Proposition 5.26. $\square$

For a representation τ of $\text{GL}(2, \mathcal{K})$, let ${}^\sigma\tau$ denote the following $\text{GL}(2, \mathcal{K})$ -module on the space of τ :

$${}^\sigma\tau : (g_{ij}) \mapsto \tau((\sigma g_{ij})), \quad \forall (g_{ij}) \in \text{GL}(2, \mathcal{K}).$$

5.30. PROPOSITION. *Let τ be an irreducible, cuspidal, admissible representation of $\text{GL}(2, \mathcal{K})$ such that: (i) $\tau \not\cong {}^\sigma\tau$, and (ii) $\omega_\tau = \mu \circ N_{\mathcal{K}/k}$ for some character μ of $k^\times$. There exist a collection $[\pi]$ of distinct ε -invariant, cuspidal, admissible G -modules, and*

an operator $A(\pi')$ in $\text{Hom}_G(\pi', \varepsilon\pi')$ for each $\pi' \in [\pi]$, such that the following holds for matching functions:

$$\sum_{\pi' \in [\pi]} \langle \pi', f \rangle_{A(\pi')} = \langle \tau \otimes_2 \mu, f_2 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There is a finite place w of F such that $F_w = k$; (ii) w is prime in E , with $E_w = \mathcal{K}$. We identify the generator σ of $\text{Gal}(\mathcal{K}/k)$ with that of $\text{Gal}(E/F)$.

Let $\mathcal{E}$ be the quadratic character of C_F corresponding to E/F . View $\mathbf{G}$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let V be the set of places of F . Let $V^{\text{un}}(E/F) \subset V$ be the set of finite places which are unramified in E .

Let w_1, w_2 be two finite places, different from w , which are unramified, prime in E .

Construct a character $\mathcal{U}$ of C_F such that: (i) $\mathcal{U}_w = \mu$; (ii) $\mathcal{U}_{w_1} = \mathcal{U}_{w_2} = 1$; (iii) $\mathcal{U}_v$ is unramified for each finite place $v \neq w, w_1, w_2$.

Construct a cuspidal automorphic representation $\mathcal{T}$ of $\text{GL}(2, \mathbb{A}_E)$ with the following properties:

- $\omega_{\mathcal{T}} = \mathcal{U} \circ N_{E/F}$;
- $\mathcal{T}_w = \tau$;
- For $i = 1, 2$, the representation $\mathcal{T}_{w_i}$ is equal to $\chi_{w_i} \text{St}_{2, w_i}$, where χ_{w_i} is a character of $E_{w_i}^\times$ such that: (i) $\chi_{w_i} \neq \sigma\chi_{w_i}$, and (ii) $\chi_{w_i}^2 = 1$;
- $\mathcal{T}_v$ is unramified for each finite place $v \neq w, w_1, w_2$.

Let π_2 be the cuspidal automorphic representation $\mathcal{T} \otimes_2 \mathcal{U}$ of $\mathbf{H}_2(\mathbb{A}_F)$. Its local component at w is $\pi_{2, w} = \tau \otimes_2 \mu$. Let $\sigma\pi_2 = \sigma\mathcal{T} \otimes_2 \mathcal{U}$.

Since the number field F is totally imaginary, $\pi_{2, v}$ and $\sigma\pi_{2, v}$ are fully induced for each archimedean place v .

Let $\{\Pi\} = \bigotimes_{v \in V} \{\Pi_v\}$ be the global packet of $\mathbf{G}(\mathbb{A}_F)$ whose global datum is the lift of the global datum parametrizing π_2 . Since all the cases of lifting to the unstable (quasi-)packets of $\mathbf{G}(\mathbb{A}_F)$ have been accounted for, the packet $\{\Pi\}$ is stable. Hence, each representation $\Pi' \in \{\Pi\}$ occurs with multiplicity one in the discrete spectrum of $\mathbf{G}$ (that is, assuming that the multiplicity one property is valid for $\text{GSp}(2)$).

Let $\{\Pi\}^\mathcal{E}$ denote the set of representations in $\{\Pi\}$ which are $\mathcal{E}$ -invariant. For each Π' in $\{\Pi\}^\mathcal{E}$, the twisted character $\langle \Pi', f \rangle_\mathcal{E}$ is a product $\prod_{v \in V} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters.

For a finite set of places S , let $\{\Pi\}_S^\mathcal{E}$ denote the set of representations in $\{\Pi\}^\mathcal{E}$ whose local components at each place $v \notin S$ are the same unramified representation Π_v^0 .

Let $S_0 = V - V^{\text{un}}(E/F)$. Let $S = S_0 \cup \{w, w_1, w_2\}$. Let ω be the split central character of π_2 . Let f and f_2 be matching functions in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega)$ and $E(w_1, w_2, \mathbf{H}_2(\mathbb{A}_F), \omega)$, respectively, such that $f_v, f_{2, v}$ are spherical for each place $v \notin S$.

By Proposition 4.30 and the stability of $\{\Pi\}$, the following trace identity holds:

$$(5.6) \quad \sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \langle \Pi', f \rangle_{\mathcal{E}, S} = \frac{1}{2} \langle \pi_2, f_2 \rangle_S + \frac{1}{2} \langle \sigma\pi_2, f_2 \rangle_S.$$

The matching condition on f, f_2 implies that $\langle \pi_2, f_2 \rangle_S = \langle \sigma\pi_2, f_2 \rangle_S$. Hence, (5.6) is equivalent to

$$(5.7) \quad \sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \langle \Pi', f \rangle_{\mathcal{E}, S} = \langle \pi_2, f_2 \rangle_S = \prod_{v \in S} \langle \pi_{2, v}, f_{2, v} \rangle.$$

For each place $v \in S - \{w, w_1, w_2\}$, the representation $\pi_{2,v}$ is fully induced. For $v = w_1, w_2$, it is equal to $\chi_v \text{St}_{2,v} \otimes 1$. By Propositions 5.8, 5.29, and Lemma 5.13, and by the linear independence of characters, we conclude that the local components at each place $v \in S - \{w\}$ of the representations in $\{\Pi\}_S^\mathcal{E}$ are all equal to the same representation $\Pi_{0,v}$.

Hence, as in the proof of Proposition 5.27, the $(S - \{w\})$ -components of (5.7) cancel up to a sign. That is, there exists a set $\{\epsilon(\Pi') = \pm 1 : \Pi' \in \{\Pi\}_S^\mathcal{E}\}$ such that

$$\sum_{\Pi' \in \{\Pi\}_S^\mathcal{E}} \epsilon(\Pi') \langle \Pi'_w, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{2,w}, f_{2,w} \rangle.$$

For each Π' in $\{\Pi\}_S^\mathcal{E}$, let $\rho(\mathcal{E})|_{\Pi', w}$ denote the operator in $\text{Hom}_{G_w}(\Pi'_w, \mathcal{E}_w \Pi'_w)$ which is defined by the restriction of $\rho(\mathcal{E})$ to Π' . Let

$$[\pi] = \{\Pi'_w : \Pi' \in \{\Pi\}_S^\mathcal{E}\}.$$

Since the local components at each place $v \neq w$ of the representations in $\{\Pi\}_S^\mathcal{E}$ are all equal to the same representation, the elements in $[\pi]$ are distinct.

Since $\pi_{2,w}$ is cuspidal, its central exponents are equal to zero. Hence, by the linear independence of central exponents ([FK, Sect. 21]), every representation in $[\pi]$ is cuspidal. For each $\pi' = \Pi'_w$ in $[\pi]$, let $A(\pi') = \epsilon(\Pi') \cdot \rho(\mathcal{E})|_{\Pi', w}$. The proposition follows. $\square$

5.5.2.1. Some Additional Twisted Character Identities for H_2 . Let μ be a character of $k^\times$. The induced representation $\varepsilon \rtimes \mu 1_{2,k}$ of G is irreducible ([ST, Prop. 4.8]). It is ε -invariant because $\varepsilon \otimes (\varepsilon \rtimes \mu 1_{2,k}) = \varepsilon \rtimes \varepsilon \mu 1_{2,k}$ is equivalent to $\varepsilon^{-1} \rtimes \varepsilon \varepsilon \mu 1_{2,k} = \varepsilon \rtimes \mu 1_{2,k}$.

5.31. CLAIM. *Suppose the characters μ and ε are unramified. There exists an intertwining operator A in $\text{Hom}_G(\varepsilon \rtimes \mu 1_{2,k}, \varepsilon \rtimes \varepsilon \mu 1_{2,k})$ such that the following holds for matching functions $f \in C(G)$, $f_2 \in C(H_2)$:*

$$\langle \varepsilon \rtimes \mu 1_{2,k}, f \rangle_A = \langle (\mu \circ N_{K/k}) 1_{2,K} \otimes_2 \mu^2 \varepsilon, f_2 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There exists a finite place w of F such that $F_w = k$; (ii) w is prime in E , with $E_w = K$. We identify the generator σ of $\text{Gal}(K/k)$ with that of $\text{Gal}(E/F)$.

Let V be the set of places of F . Let $V^{\text{un}}(E/F) \subset V$ be the set of finite places which are unramified in E .

Let $\mathcal{E}$ be the quadratic character of C_F corresponding to E/F . We view $\mathbf{G}$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let w_1, w_2 be two (finite) places, different from w , which are prime in E .

Construct a character $\mathcal{U}$ of C_F such that: (i) $\mathcal{U}_w = \mu$, and (ii) $\mathcal{U}_v$ is unramified for each finite place $v \neq w, w_1, w_2$.

Let 1_2 be the trivial representation of $\text{GL}(2, \mathbb{A}_F)$. Let $\mathcal{T}$ be the following representation of the Heisenberg parabolic subgroup P of $\mathbf{G}(\mathbb{A}_F)$:

$$\mathcal{T} = \mathcal{E} \otimes \mathcal{U} 1_2 : \begin{pmatrix} a & * & * & * \\ & g_2 & * & * \\ & & \det g_2 & * \\ & & & a \end{pmatrix} \mapsto \mathcal{E}(a) \mathcal{U}(\det g_2).$$

Let $\Pi = \mathcal{E} \rtimes \mathcal{U} 1_2$, the representation of $\mathbf{G}(\mathbb{A}_F)$ parabolically induced from $\mathcal{T}$.

- At a place v where $\mathcal{E}_v$ is trivial, Π_v is reducible, with two nontempered constituents $\Pi_v^+ = L(\nu_v, 1 \rtimes \nu_v^{-1/2} \mathcal{U}_v)$ and $\Pi_v^- = L(\nu_v^{1/2} \text{St}_{2,F_v}, \nu_v^{-1/2} \mathcal{U}_v)$ ([ST, Lemma 3.8]).
- If $\mathcal{E}_v \neq 1$, then Π_v is irreducible. We put $\Pi_v^+ := \Pi_v$ and $\Pi_v^- := 0$.

In particular, Π_w is by construction irreducible and unramified.

Let $\{\Pi\} = \bigotimes_{v \in V} \{\Pi_v^+, \Pi_v^-\}$ be the set of irreducible *automorphic* constituents of Π . That is, an irreducible constituent Π' of Π is a member of $\{\Pi\}$ if and only if Π'_v is unramified for almost all v .

The representation Π is $\mathcal{E}$ -invariant. The operator $I_{P,\mathcal{T}}(\mathcal{E})$, defined in Section 3.1.1, intertwines Π with $\mathcal{E}\Pi$. For an irreducible constituent Π' of Π , and a test function f on $\mathbf{G}(\mathbb{A}_F)$, put $\langle \Pi', f \rangle_{\mathcal{E}} := \text{tr } \Pi'(f) I_{P,\mathcal{T}}(\mathcal{E})$. It is a product $\prod_{v \in V} \langle \Pi'_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters.

Let $1_{2,E}$ be the trivial representation of $\text{GL}(2, \mathbb{A}_E)$. Let π_2 be the representation $(\mathcal{U} \circ \text{N}_{E/F}) 1_{2,E} \otimes_2 \mathcal{U}^2 \mathcal{E}$ of $\mathbf{H}_2(\mathbb{A}_F)$. In particular, $\pi_{2,w}$ is equivalent to $(\mu \circ \text{N}_{\mathcal{K}/k}) 1_{2,\mathcal{K}} \otimes_2 \mu^2 \varepsilon$.

Let $S_0 = V - V^{\text{un}}(E/F)$. Let $S_1 = S_0 \cup \{w_1, w_2\}$. Then, Π_v and $\pi_{2,v}$ are unramified for each place $v \notin S_1$. The split central character of π_2 is equal to $\mathcal{U}^2 \mathcal{E}$. Let $f \in E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \mathcal{U}^2 \mathcal{E})$ and $f_2 \in E(w_1, w_2, \mathbf{H}_2(\mathbb{A}_F), \mathcal{U}^2 \mathcal{E})$ be matching functions such that $f_v, f_{2,v}$ are spherical for each $v \notin S_1$. By Claim 4.38, the following holds:

$$(5.8) \quad \prod_{v \in S_1} \left(\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} - \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v} \right) = \prod_{v \in S_1} \langle \pi_{2,v}, f_{2,v} \rangle.$$

For each Π' in $\{\Pi\}$, Π'_w is equal to $\Pi_w = \varepsilon \rtimes \mu 1_{2,k}$. Let $S = S_1 \cup \{w\}$. Let f, f_2 be matching functions with elliptic components at w_1, w_2 , and spherical components at all the places outside of S . Applying (5.8) to S , we obtain:

$$(5.9) \quad \langle \Pi_w, f \rangle_{\mathcal{E}_w} \prod_{v \in S_1} \left(\langle \Pi_v^+, f_v \rangle_{\mathcal{E}_v} - \langle \Pi_v^-, f_v \rangle_{\mathcal{E}_v} \right) = \langle \pi_{2,w}, f_{2,w} \rangle \prod_{v \in S_1} \langle \pi_{2,v}, f_{2,v} \rangle_{\mathcal{E}_v}.$$

By (5.8), the S_1 -components of the above equation cancel; hence,

$$\langle \pi_w, f_w \rangle_{\mathcal{E}_w} = \langle \pi_{2,w}, f_{2,w} \rangle.$$

Let A be the operator in $\text{Hom}_{G_w}(\Pi_w, \mathcal{E}_w \Pi_w)$ which is induced from $I_{P,\mathcal{T}}(\mathcal{E})$. The claim follows. $\square$

5.32. COROLLARY. *Suppose the extension $\mathcal{K}/k$ is unramified. Let μ be an unramified character of $k^\times$. Let π be the induced representation $\varepsilon \rtimes \mu \text{St}_{2,k}$ of G . There exists an operator A in $\text{Hom}_G(\pi, \varepsilon \pi)$ such that the following holds for matching functions:*

$$\langle \pi, f \rangle_A = \langle (\mu \circ \text{N}_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}} \otimes_2 \mu^2 \varepsilon, f_2 \rangle.$$

PROOF. We have:

$$(\mu \circ \text{N}_{\mathcal{K}/k}) I(\nu_{\mathcal{K}}^{1/2}, \nu_{\mathcal{K}}^{-1/2}) = (\mu \circ \text{N}_{\mathcal{K}/k}) 1_{2,\mathcal{K}} + (\mu \circ \text{N}_{\mathcal{K}/k}) \text{St}_{2,\mathcal{K}}.$$

Since the extension $\mathcal{K}/k$ is unramified, ε is an unramified character of $k^\times$, and $\nu_{\mathcal{K}}$ is equal to $\nu \circ \text{N}_{\mathcal{K}/k}$. The corollary then follows from Proposition 5.8 and the previous claim. $\square$

5.33. PROPOSITION. *Let τ be an irreducible, cuspidal, non- $\mathcal{K}$ -monomial representation of $\text{GL}(2, k)$. Let π be the induced representation $\varepsilon \rtimes \tau$ of G . There exists an operator A in $\text{Hom}_G(\pi, \varepsilon \pi)$ such that the following holds for matching functions:*

$$\langle \pi, f \rangle_A = \langle B_{\mathcal{K}/k} \tau \otimes_2 \omega_{\tau} \varepsilon, f_2 \rangle.$$

PROOF. Construct a totally imaginary number field F and a quadratic extension E of F such that: (i) There exists a finite place w of F such that $F_w = k$; (ii) w is prime in E , with $E_w = \mathcal{K}$. We identify the generator σ of $\text{Gal}(\mathcal{K}/k)$ with that of $\text{Gal}(E/F)$.

Let $\mathcal{E}$ be a quadratic character of C_F corresponding to E/F . We consider $\mathbf{G}$ as an F -group, and $\mathbf{H}_1, \mathbf{H}_2$ as the elliptic $\mathcal{E}$ -endoscopic groups of $\mathbf{G}$ over F .

Let V be the set of places of F . Let $V^{\text{un}}(E/F) \subset V$ be the set of finite places which are unramified in E .

Let w_1, w_2 be two (finite) places, different from w , which are unramified, prime in E . Construct a cuspidal automorphic representation $\mathcal{T}$ of $\text{GL}(2, \mathbb{A}_F)$ such that:

- $\mathcal{T}_w = \tau$;
- $\mathcal{T}_{w_i} = \text{St}_{2, F_{w_i}}$ ($i = 1, 2$);
- $\mathcal{T}_v$ is unramified for each finite place v .

Let $\pi_2 = B_{E/F} \mathcal{T} \otimes_2 \omega_{\mathcal{T}} \mathcal{E}$. It is a cuspidal automorphic representation of $\mathbf{H}_2(\mathbb{A}_F)$, with $\pi_{2,w} = B_{K/k} \tau \otimes_2 \omega_{\tau} \mathcal{E}$.

Let Π be the $\mathbf{G}(\mathbb{A}_F)$ -module $\mathcal{E} \rtimes \mathcal{T}$. It is parabolically induced from the representation $\Omega = \mathcal{E} \otimes \mathcal{T}$ of the Heisenberg parabolic subgroup P of $\mathbf{G}(\mathbb{A}_F)$. In particular, $\Pi_w = \mathcal{E} \rtimes \tau$.

Since F is totally imaginary, $\mathcal{T}_v$ is a fully induced representation of $\text{GL}(2, \mathbb{C})$ for each archimedean place v . For each finite place $v \neq w, w_1, w_2$, the representation $\mathcal{T}_v$ is by construction unramified, hence fully induced. Therefore, by [ST, Thm 4.2], [ST, Lemma 3.4], and [ST, Prop. 4.8], the representation Π_v is irreducible for each place v . Consequently, Π is irreducible.

The operator $I_{P,\Omega}(\mathcal{E})$, defined in Section 3.1.1, intertwines Π with $\mathcal{E}\Pi$. For a test function f on $\mathbf{G}(\mathbb{A}_F)$, put $\langle \Pi, f \rangle_{\mathcal{E}_v} := \text{tr } \Pi(f) I_{P,\Omega}(\mathcal{E})$. It is a product $\prod_{v \in V} \langle \Pi_v, f_v \rangle_{\mathcal{E}_v}$ of twisted local characters.

Let $S_0 = V - V^{\text{un}}(E/F)$. Let $S = S_0 \cup \{w, w_1, w_2\}$. The central character of Π and the split central character of π_2 are both equal to $\omega_{\mathcal{T}} \mathcal{E}$. Let f in $E(w_1, w_2, \mathbf{G}(\mathbb{A}_F), \omega_{\mathcal{T}} \mathcal{E})$ and f_2 in $E(w_1, w_2, \mathbf{H}_2(\mathbb{A}_F), \omega_{\mathcal{T}} \mathcal{E})$ be matching functions such that $f_v, f_{2,v}$ are spherical for each place $v \notin S$.

By Claim 4.38, we have:

$$(5.10) \quad \prod_{v \in S} \langle \Pi_v, f_v \rangle_{\mathcal{E}_v} = \prod_{v \in S} \langle \pi_{2,v}, f_{2,v} \rangle.$$

Note that since Π is irreducible, the term Π_v^- in Claim 4.38 is zero for each place v .

By Proposition 5.8, Lemma 5.13, and Corollary 5.32, the $(S_0 \cup \{w_1, w_2\})$ -components of (5.10) cancel up to a sign $\epsilon = \pm 1$. Hence,

$$(5.11) \quad \langle \Pi_w, f_w \rangle_{\mathcal{E}_w} = \epsilon \langle \pi_{2,w}, f_{2,w} \rangle.$$

Let $I_{P,\Omega}(\mathcal{E})_w \in \text{Hom}_{G_w}(\Pi_w, \mathcal{E}_w \Pi_w)$ be the operator induced from $I_{P,\Omega}(\mathcal{E})$. Let $A = \epsilon \cdot I_{P,\Omega}(\mathcal{E})_w$. The proposition follows. $\square$

APPENDIX A

Summary of Global Lifting

A.0.2.2. Notation:

- E/F a quadratic extension of number fields, σ the generator of $\text{Gal}(E/F)$.
- For a number field Q , C_Q denotes the idèle class group of Q , $\widehat{C_Q}$ the set of (unitary) characters of C_Q , and $1_{2,Q}$ the trivial representation of $\text{GL}(2, \mathbb{A}_Q)$.
- ε a character of C_F , with $\varepsilon \neq 1 = \varepsilon \circ N_{E/F}$.
- For an automorphic representation τ of $\text{GL}(2, \mathbb{A}_E)$, $\pi(\tau)$ denotes the automorphic $\text{GL}(4, \mathbb{A}_F)$ -module associated with τ via the twisted endoscopic lifting from $\text{R}_{E/F}\text{GL}(2)$ to $\text{GL}(4)$. $\text{R}_{E/F}$ denotes restriction of scalars from E to F .
- $\mathcal{A}(\text{GL}(2))$ denotes the set of equivalence classes of automorphic representations of $\text{GL}(2, \mathbb{A}_F)$. For $\tau \in \mathcal{A}(\text{GL}(2))$, $B_{E/F}\tau$ denotes the automorphic $\text{GL}(2, \mathbb{A}_E)$ -module obtained from τ via base change.

In the following tables, each entry in the second row is a (pseudo-)(quasi-)packet of the group above it in the first row. Two (quasi-)packets belong to the same row if the global datum parametrizing one is the lift of the global datum parametrizing the other via the L -group embedding associated with their corresponding groups (see Section 4.1.1.1).

A.1. Unstable (quasi-)packets of G

- (1) $\tau \in \mathcal{A}(\text{GL}(2))$ cuspidal non- E -monomial, or one dimensional:

$\text{GL}(4)$	G	H_1	H_2
$I(\tau, \varepsilon\tau)$	$[\tau, \varepsilon\tau]$	$\tau \otimes_1 1$	$B_{E/F}\tau \otimes_2 \omega_\tau$

$[\tau, \varepsilon\tau]$ is the unstable (quasi-)packet which lifts to $I(\tau, \varepsilon\tau)$ of $\text{GL}(4, \mathbb{A}_F)$ (see Section 4.3.1).

REMARK: The above table also covers representations of the form $\tau \otimes_1 \chi$, with $\chi = {}^\sigma\chi$, because by Hilbert 90 $\chi = \mu \circ N_{E/F}$ for some $\mu \in \widehat{C_F}$, and $\tau \otimes_1 \mu \circ N_{E/F} = \mu\tau \otimes_1 1$.

- (2) $\theta, \chi \in \widehat{C_E}$; none of $\theta, \chi, \theta\chi, \theta {}^\sigma\chi$ is fixed by σ .
 $\{\pi(\theta) \otimes_1 \chi\}$ denotes the set of the packets $\pi(\theta) \otimes_1 \chi, \pi(\theta) \otimes_1 {}^\sigma\chi, \pi(\chi) \otimes_1 \theta, \pi(\chi) \otimes_1 {}^\sigma\theta$.

$\text{GL}(4)$	G	H_1
$I(\pi(\theta\chi), \pi(\theta {}^\sigma\chi))$	$[\pi(\theta\chi), \pi(\theta {}^\sigma\chi)]$	$\{\pi(\theta) \otimes_1 \chi\}$

A.2. Stable (quasi-)packets

Every ε -invariant stable (quasi-)packet of $\mathbf{G}(\mathbb{A}_F)$ is lifted **exclusively** from either $\mathbf{H}_1$ or $\mathbf{H}_2$.

A.2.1. Stable (quasi-)packets lifted from H_1 via $\xi_1 : {}^L H_1 \hookrightarrow {}^L G$.

- (1)
- $\tau \in \mathcal{A}(\mathrm{GL}(2))$
- cuspidal;
- $\chi \in \widehat{C_E}$
- with
- $\chi \neq {}^\sigma \chi$
- .

$\mathrm{GL}(4)$	G	H_1
$\pi(\chi B_{E/F} \tau)$	$\xi_1^*(\tau \otimes_1 \chi)$	$\tau \otimes_1 \chi, \tau \otimes_1 {}^\sigma \chi$

$\xi_1^*(\tau \otimes_1 \chi)$ denotes the packet of $\mathbf{G}(\mathbb{A}_F)$ which is the lift of $\tau \otimes_1 \chi$. All of its members are cuspidal.

- (2)
- $\chi \in \widehat{C_E}$
- with
- $\chi \neq {}^\sigma \chi$
- .

$\mathrm{GL}(4)$	G	H_1
$J(\nu^{1/2} \pi(\chi), \nu^{-1/2} \pi(\chi))$	$L(\varepsilon \nu, \nu^{-1/2} \pi(\chi))$	$1_{2,F} \otimes_1 \chi,$ $1_{2,F} \otimes_1 {}^\sigma \chi$

REMARK: Since $\mu 1_{2,F} \otimes_1 \chi = 1_{2,F} \otimes_1 (\mu \circ N_{E/F}) \chi$, $\forall \mu \in \widehat{C_F}$, the table above covers all cases of $\tau \otimes_1 \chi$ where τ is a one dimensional representation of $\mathrm{GL}(2, \mathbb{A}_F)$.

A.2.2. Stable (quasi-)packets lifted from H_2 via $\xi_2 : {}^L H_2 \hookrightarrow {}^L G$.

- (1)
- $\tau \in \mathcal{A}(\mathrm{R}_{E/F} \mathrm{GL}(2))$
- cuspidal;
- $\mu \in \widehat{C_F}$
- with
- $\omega_\tau = \mu \circ N_{E/F}$
- .

$\mathrm{GL}(4)$	G	H_2
$\pi(\tau)$	$\xi_2^*(\tau \otimes_2 \mu)$	$\tau \otimes_2 \mu,$ ${}^\sigma \tau \otimes_2 \mu$

$\xi_2^*(\tau \otimes_2 \mu)$ denotes the packet of $\mathbf{G}(\mathbb{A}_F)$ which is the lift of $\tau \otimes_2 \mu$. All of its members are cuspidal.

- (2)
- $\chi \in \widehat{C_E}$
- ,
- $\varepsilon' \neq \varepsilon \in \widehat{C_F}$
- , with (i)
- $\varepsilon' \neq (\varepsilon')^2 = 1$
- , (ii)
- ${}^\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$
- ;
- $\zeta \in \{\varepsilon', \varepsilon \varepsilon'\}$
- .

$\mathrm{GL}(4)$	G	H_2
$J(\nu^{1/2} \pi(\chi), \nu^{-1/2} \pi(\chi))$	$L(\varepsilon \zeta \nu, \nu^{-1/2} \pi(\chi))$	$\chi 1_{2,E} \otimes_2 \chi _{C_F} \cdot \zeta,$ ${}^\sigma \chi 1_{2,E} \otimes_2 \chi _{C_F} \cdot \zeta$

Each ε -invariant, discrete spectrum, automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ with two elliptic local components appears in exactly one of the tables in Sections A.1 and A.2.

A.3. Induced representations

- (1)
- $\mu \in \widehat{C_F}$
- .

$\mathrm{GL}(4)$	G	H_1	H_2
$I(\mu, \mu, \varepsilon \mu, \varepsilon \mu)$	$\varepsilon \times \varepsilon \rtimes \mu$	$\mu I(1, 1) \otimes_1 1$	$(\mu \circ N_{E/F}) I(1, 1) \otimes_2 \mu^2$

- (2)
- $\mu \in \widehat{C_F}$
- ;
- $\theta, \chi \in \widehat{C_E}$
- ; with (i)
- $\theta \neq {}^\sigma \theta$
- , (ii)
- $\chi \neq {}^\sigma \chi$
- , (iii)
- $\theta \chi = \mu \circ N_{E/F}$
- .
-
- (a)
- $\chi^2 \neq {}^\sigma \chi^2$
- .

$\mathrm{GL}(4)$	G	H_1
$I(\mu \pi({}^\sigma \chi / \chi), \mu, \mu \varepsilon)$	$\pi({}^\sigma \chi / \chi) \rtimes \mu$	$\pi(\theta) \otimes_1 \chi, \pi(\theta) \otimes_1 {}^\sigma \chi$

- (b)
- $\exists \varepsilon' \in \widehat{C_F}$
- with (i)
- $\varepsilon' \neq (\varepsilon')^2 = 1$
- , (ii)
- ${}^\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$
- .

$\mathrm{GL}(4)$	G	H_1
$I(\mu, \mu \varepsilon' \varepsilon, \mu \varepsilon, \mu \varepsilon')$	$\varepsilon' \times \varepsilon' \varepsilon \rtimes \mu$	$\pi(\theta) \otimes_1 \chi$

- (3)
- $\theta \in \widehat{C_E}$
- with
- $\theta \neq {}^\sigma \theta$
- .

$\mathrm{GL}(4)$	G	H_1
$I(\pi(\theta), \pi(\theta))$	$1 \rtimes \pi(\theta)$	$\pi(\theta) \otimes_1 1$
$I(\pi(\theta), \pi(\theta))$	$\varepsilon \rtimes \pi(\theta)$	$I(1, 1) \otimes_1 \theta, I(1, 1) \otimes_1 {}^\sigma \theta$

- (4) $[E', F] = 2$, $E' \neq E$, with $\text{Gal}(E'/F) = \langle \sigma' \rangle$.

$\varepsilon' \in \widehat{C_F}$ with $\varepsilon' \neq \varepsilon' \circ N_{E'/F} = 1$.

$\chi \in \widehat{C_E}$; $\mu, \theta \in \widehat{C_{E'}}$; (i) ${}^\sigma \chi / \chi = \varepsilon' \circ N_{E/F}$, (ii) $\mu \circ N_{EE'/E'} = \chi \circ N_{EE'/E}$,

- (iii) $\mu \theta \neq {}^{\sigma'}(\mu \theta)$.

GL(4)	G	H_1
$I(\pi_{E'}(\mu \theta), \varepsilon \pi_{E'}(\mu \theta))$	$\varepsilon \varepsilon' \rtimes \pi_{E'}(\mu \theta)$	$\pi_{E'}(\theta) \otimes_1 \chi$

- (5) $\tau \in \mathcal{A}(\text{GL}(2))$ cuspidal non- E -monomial.

GL(4)	G	H_2
$I(\tau, \varepsilon \tau)$	$\varepsilon \rtimes \tau$	$B_{E/F} \tau \otimes_2 \omega_\tau \varepsilon$

- (6) $\mu \in \widehat{C_F}$.

GL(4)	G	H_2
$I(\mu, \mu, \varepsilon \mu, \varepsilon \mu)$	$1 \times \varepsilon \rtimes \mu$	$(\mu \circ N_{E/F}) I(1, 1) \otimes_2 \mu^2 \varepsilon$

Each parabolically induced $\mathbf{G}(\mathbb{A}_F)$ -module which contributes discretely to the fine χ -expansion of the ε -twisted trace formula of $\mathbf{G}(\mathbb{A}_F)$ appears in exactly one of the tables in Section A.3.

APPENDIX B

Fundamental Lemma

Let F be a p -adic field. Let ε be a character of $F^\times$ with $\varepsilon^2 = 1$. To prove the Fundamental Lemma in the context of $\mathbf{G}$ and $\mathbf{H}_1, \mathbf{H}_2$, by [H2] it suffices to prove it for the case where F has odd residual characteristic, and then only for the unit elements in the Hecke algebras of the groups, when $\varepsilon \neq 1$ and also when $\varepsilon = 1$. Moreover, we may assume that ε is an unramified character. The case of $\varepsilon = 1$ is straightforward, given the results of [F3]. Hence, in this chapter we assume that F has odd residual characteristic, and that ε is nontrivial.

We fix once and for all an element $A \in F^\times - F^{\times 2}$ such that $F_A := F(\sqrt{A})$ is the unramified quadratic extension of F which corresponds to ε via local class field theory. Consider $\mathbf{G} = \mathrm{GSp}(2)$ as an F -group. Let $G = \mathbf{G}(F)$. Recall the definitions in Chapter 2 of the elliptic ε -endoscopic groups of $\mathbf{G}$:

- (1) $H_1 = \mathbf{H}_1(F) = \{(g, \xi) \in \mathrm{GL}(2, F) \times F_A^\times : \det g = \mathrm{N}_{F_A/F} x\}$.
- (2) $H_2 = \mathbf{H}_2(F)$
 $= \{(g, c) \in \mathrm{GL}(2, F_A) \times F^\times\} / \{(\mathrm{diag}(z, z), \mathrm{N}_{F_A/F} z^{-1}) : z \in F_A^\times\}$.

Let R be the ring of integers in F . Fix a uniformizer ϖ of F . Let K be the maximal compact, hyperspecial subgroup $\mathbf{G}(R)$ of G . Let 1_K in $C_c^\infty(G)$ be the characteristic function of K . Let dg be the Haar measure on G for which the volume of K is one. For a function $f \in C_c^\infty(G)$, define the ε -twisted orbital integral of f at an element $\delta \in G$ as follows:

$$O_\delta(f) = \int_{Z_G(\delta) \backslash G} f(g^{-1} \delta g) \varepsilon(g) dg.$$

Here, $Z_G(\delta)$ is the centralizer of δ in G .

For $i = 1, 2$, the group $\mathbf{H}_i$ is defined over R . Let $K_i = \mathbf{H}_i(R)$. Let dh_i be the Haar measure on H_i for which K_i has volume one. Let $1_{K_i} \in C_c^\infty(H_i)$ be the characteristic function of K_i . For $H = H_1$ or H_2 , $f_H \in C_c^\infty(H)$, and $\delta_H \in H$, let

$$\begin{aligned} O_{\delta_H}(f_H) &= \int_{Z_H(\delta_H) \backslash H} f_H(h^{-1} \delta_H h) dh, \\ SO_{\delta_H}(f_H) &= \sum_{\delta'_H} O_{\delta_H}(f_H). \end{aligned}$$

Here, the sum in the second expression is over representatives δ'_H of the F -conjugacy classes within the stable conjugacy class of δ_H .

B.1. THEOREM. *Let $i = 1, 2$. For a regular element $\delta \in G$ and a $\mathbf{G}$ -regular δ_i in H_i which is a norm of δ , there exists a choice of transfer factor $\Delta(\delta_i, \delta) \in \mathbb{C}$ such that the following holds:*

$$SO_{\delta_i}(1_{K_i}) = \sum_{\delta'} \Delta(\delta_i, \delta') O_{\delta'}(1_K).$$

The sum on the right is over representatives δ' of the F -conjugacy classes within the stable conjugacy class of δ .

REMARK. Since $O_{\delta'}(1_K)$ is equal to zero if δ is not stably conjugate to an element in K , it suffices to prove the theorem for the elements δ which lie in K .

We shall prove the theorem only for the *elliptic* regular elements. For elements which lie in parabolic subgroups, the theorem reduces via Harish-Chandra descent (see, for example, [H]) to the Fundamental Lemma for lower rank groups.

An element $g \in G$ is called **absolutely semisimple** if $g^m = 1$ for some integer m prime to the residual characteristic of F . An element g is called **topologically unipotent** if $g^{(q^N)}$ converges to 1 as the integer N approaches infinity. Here, q is the cardinality of the residue field of F . By an extension of a theorem of Kazhdan's ([F3, Prop. I. H. 1]), an element $\delta \in K$ decomposes uniquely as $\delta = su = us$, where s is absolutely semisimple, u is topologically unipotent, and $s, u \in K$.

With regard to proving Theorem B.1, we restrict further our attention to the elliptic regular elements which are topologically unipotent. For elements whose absolutely semisimple parts are nontrivial, the method of semisimple reduction ([F3, Part. III]) reduces the theorem to the Fundamental Lemma for lower rank groups.

B.0.0.1. *Notation.*

- For a field k and an element $B \in k^\times$, let k_B denote the possibly trivial field extension $k(\sqrt{B})$ of k .
- Let $\mathrm{GL}(2, k)^{k_B}$ denote the subgroup $\{g \in \mathrm{GL}(2, k) : \det g \in N_{k_B/k} k_B^\times\}$ of $\mathrm{GL}(2, k)$.
- For $\xi = a + b\sqrt{B} \in k_B^\times$, $a, b \in k$, put

$$\phi^{k:B}(\xi) := \phi^B(\xi) := \begin{pmatrix} a & bB \\ b & a \end{pmatrix}.$$

Observe that if $[k_B : k] = 2$, then $\det \phi^B(\xi) = N_{k_B/k} \xi$ for all $\xi \in k_B^\times$.

- For $\rho \in k^\times$, $\xi \in k_B$, put

$$\phi_\rho^{k:B}(\xi) := \phi_\rho^B(\xi) := \begin{pmatrix} a & bB\rho \\ b/\rho & a \end{pmatrix}.$$

- Let T_B^ρ denote the maximal torus $\{\phi_\rho^B(\xi) : \xi \in k_B^\times\}$ in $\mathrm{GL}(2, k)$. It is elliptic if $[k_B : k] = 2$.
- Let $L = k_B(\sqrt{\xi})$, where $\xi \in k_B^\times$. For $l = \alpha + \beta\sqrt{\xi} \in L^\times$, $\alpha, \beta \in k_B$, put

$$\phi^{k:B;\xi}(l) := \begin{pmatrix} \phi^B(\alpha) & \phi^B(\beta)\phi^B(\xi) \\ \phi^B(\beta) & \phi^B(\alpha) \end{pmatrix}.$$

- For $\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} x & y \\ z & t \end{pmatrix} \in \mathrm{GL}(2)$, put

$$\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} x & y \\ z & t \end{pmatrix}\right] := \begin{pmatrix} a & x & b & y \\ c & z & d & t \end{pmatrix} \in \mathrm{GL}(4).$$

B.1. Norm Correspondence—Elliptic Elements

We now describe the norm correspondence between the elliptic regular elements of G and the elliptic $\mathbf{G}$ -regular elements of H_1, H_2 . But first, we need to classify the stable conjugacy classes of the elliptic regular elements of the groups. This is equivalent to classifying the stable conjugacy classes of the elliptic maximal tori. We list below the

representatives of these stable conjugacy classes. Moreover, we list the conjugacy classes within each stable conjugacy class.

B.1.1. Elliptic Tori of G . The ε -twisted orbital integral of a function on G at an element $\delta \in G$ is zero if $Z_\delta(G)$ is not contained in the kernel of ε . In fact, as we shall see, elliptic $\mathbf{G}$ -regular elements of the ε -endoscopic groups can be the norms of only those elliptic regular elements in G whose centralizers are contained in $\ker \varepsilon$. Thus, for our purpose, it suffices to list only the stable conjugacy classes of the elliptic tori which lie in $\ker \varepsilon$.

Recall that F_A is the quadratic extension of F which corresponds to ε via local class field theory. The stable conjugacy classes of the elliptic tori of G are computed in [F3, Sect. I. C]. The representatives of those classes which lie in the kernel of ε are as follows:

- $T_{I,A} = \left\{ [\phi^A(\gamma), \phi^A(\zeta)] : \gamma, \zeta \in F_A^\times; N_{F_A/F}\gamma = N_{F_A/F}\zeta \right\};$
- $T_{III,A,AD} = \left\{ [\phi^A(\gamma), \phi^{AD}(\zeta)] : \gamma \in F_A^\times, \zeta \in F_{AD}^\times; N_{F_A/F}\gamma = N_{F_{AD}/F}\zeta \right\},$
 $D, AD \in F^\times - AF^{\times 2};$
- $T_{III,D,AD} = \left\{ [\phi^D(\gamma), \phi^{AD}(\zeta)] : \gamma \in F_D^\times, \zeta \in F_{AD}^\times; N_{F_D/F}\gamma = N_{F_{AD}/F}\zeta \right\},$
 $D, AD \in F^\times - AF^{\times 2};$
- $T_{IV,A,D} = \left\{ \phi^{F:A:D}(l) : l \in L^\times; N_{L/F_A}l \in F^\times \right\},$
 $L = F_A(\sqrt{D}), D \in F_A^\times - F_A^{\times 2}.$

Note that $T_{III,A,AD}$ and $T_{III,D,AD}$ are distinct cases, for A depends on ε and is not arbitrary.

With the exception of $T_{I,A}$, the conjugacy class of each elliptic maximal torus listed above are stable. In the case of $T_{I,A}$, there are two conjugacy classes within the stable conjugacy class, described as follows:

For $\xi = a + b\sqrt{A} \in F_A^\times$ ($a, b \in F$), put $\phi_\varpi^A(\xi) := \begin{pmatrix} a & b\varpi A \\ b/\varpi & a \end{pmatrix}$. Put

$$T_{I,A}^\varpi := \begin{pmatrix} 1 & & \\ & 1 & \\ & & \varpi \end{pmatrix}^{-1} T_{I,A} \begin{pmatrix} 1 & & \\ & 1 & \\ & & \varpi \end{pmatrix} = \{ [\phi^A(\gamma), \phi_\varpi^A(\zeta)] : N_{F_A/F}\gamma = N_{F_A/F}\zeta \}.$$

From [F3, Sect. I. C], the tori $T_{I,A}$ and $T_{I,A}^\varpi$ are representatives of the conjugacy classes within the stable conjugacy class of $T_{I,A}$.

B.1.2. Elliptic Tori of H_1 . There are two stable conjugacy classes of elliptic tori in H_1 . They are represented by the following tori:

- $T_{1,A} = \{ (\phi^A(\gamma), \zeta) : \gamma, \zeta \in F_A^\times; N_{F_A/F}\gamma = N_{F_A/F}\zeta \};$
- $T_{1,AD} = \{ (\phi^{AD}(\gamma), \zeta) : \gamma \in F_{AD}^\times, \zeta \in F_A^\times; N_{F_{AD}/F}\gamma = N_{F_A/F}\zeta \},$
 $D, AD \in F^\times - AF^{\times 2}.$

An element of H_1 has the form $(g, \zeta) \in \mathrm{GL}(2, F) \times F_A^\times$, with $\det g = N_{F_A/F}\zeta$. Suppose two elements $(g, \zeta), (g', \zeta')$ in H_1 are stably conjugate to each other. Then, $\zeta = \zeta'$. Since $\mathbf{H}_1(\bar{F}) = \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times \times \bar{F}^\times$, and there is no distinction between stable conjugacy and F -conjugacy for $\mathrm{GL}(2)$, the elements g, g' are conjugate via some element of $\mathrm{GL}(2, F)$.

Let $\mathrm{GL}(2, F)^{F_A}$ denote the subgroup of elements in $\mathrm{GL}(2, F)$ whose determinants lie in $N_{F_A/F} F_A^\times$. We have an exact sequence:

$$1 \rightarrow \mathrm{GL}(2, F)^{F_A} \rightarrow \mathrm{GL}(2, F) \xrightarrow{\det} F^\times / N_{F_A/F} F_A^\times \rightarrow 1.$$

Since $[F^\times : N_{F_A/F} F_A^\times] = 2$, $\mathrm{GL}(2, F)^{F_A}$ is a subgroup of index 2 in $\mathrm{GL}(2, F)$. Thus, there are at most two conjugacy classes within the stable conjugacy class of each element $g \in \mathrm{GL}(2, F)^{F_A}$.

B.2. CLAIM. *A regular element $\delta_1 = (\phi^A(\gamma), \zeta) \in T_{1,A}$ is stably conjugate but not conjugate to $\delta_1^\varpi = (\phi_\varpi^A(\gamma), \zeta)$.*

PROOF. Since $\phi_\varpi^A(\gamma) = \begin{pmatrix} 1 & \varpi \\ & 1 \end{pmatrix}^{-1} \phi^A(\gamma) \begin{pmatrix} 1 & \varpi \\ & 1 \end{pmatrix}$, it is clear that δ_1 is stably conjugate to δ_1^ϖ . Suppose g is an element in $\mathrm{GL}(2, F)$ such that $g^{-1} \phi^A(\gamma) g = \phi_\varpi^A(\gamma)$. Then, $g \begin{pmatrix} 1 & \varpi \\ & 1 \end{pmatrix}$ commutes with $\phi^A(\gamma)$; hence, $g \begin{pmatrix} 1 & \varpi \\ & 1 \end{pmatrix} = \begin{pmatrix} x & Az \\ z & x \end{pmatrix}$ for some $x, z \in F$. The determinant of g is therefore equal to $\varpi^{-1}(x^2 - z^2 A)$, which does not lie in $N_{F_A/F} F_A^\times$, for $x^2 - z^2 A$ lies in $N_{F_A/F} F_A^\times$, while ϖ does not because F_A/F is unramified. Hence, $g \notin \mathrm{GL}(2, F)^{F_A}$, which implies that δ_1^ϖ is not conjugate to δ_1 in H_1 . $\square$

B.3. COROLLARY. *The conjugacy class of the tori in H_1 which are stably conjugate but not conjugate to $T_{1,A}$ is represented by*

$$T_{1,A}^\varpi := \{(\phi_\varpi^A(\gamma), \zeta) : \gamma, \zeta \in F_A^\times; N_{F_A/F} \gamma = N_{F_A/F} \zeta\}.$$

B.4. CLAIM. *The conjugacy class of $T_{1,AD}$ is stable.*

PROOF. Suppose an element $\delta' = (t', \zeta)$ in H_1 is stably conjugate to the element $\delta = (\phi^{AD}(\xi), \zeta) \in T_{1,AD}$, where $N_{F_{AD}/F} \xi = N_{F_A/F} \zeta$. There exists g in $\mathrm{GL}(2, F)$ such that $t' = g^{-1} \phi^{AD}(\xi) g$. Since $\mathrm{GL}(2, F)$ is the disjoint union of $\mathrm{GL}(2, F)^{F_A}$ and $\mathrm{GL}(2, F)^{F_A} \mathrm{diag}(1, \varpi)$, we may assume that $g = \mathrm{diag}(1, \varpi)$. Since by assumption the residual characteristic of F is odd, and F_A/F is unramified, the extension F_{AD}/F is ramified. Hence, there exists an element $\gamma \in F_{AD}^\times$ such that $N_{F_{AD}/F} \gamma = \varpi^{-1}$. Since $\phi^{AD}(\xi)$ commutes with $\phi^{AD}(\gamma)$, t' is equal to

$$(\phi^{AD}(\gamma) g)^{-1} \cdot \phi^{AD}(\xi) \cdot (\phi^{AD}(\gamma) g).$$

The element $\phi^{AD}(\gamma) g$, having determinant 1, lies in $\mathrm{GL}(2, F)^{F_A}$; hence, δ' is conjugate to δ in H_1 . $\square$

B.1.3. Elliptic Tori of H_2 . For an element $D \in F_A^\times$, and $l = \alpha + \beta\sqrt{D} \in F_A(\sqrt{D})^\times$ ($\alpha, \beta \in F_A$), put

$$\phi^{F_A:D}(l) := \begin{pmatrix} \alpha & \beta D \\ \beta & \alpha \end{pmatrix} \in \mathrm{GL}(2, F_A).$$

A maximal elliptic torus in H_2 has the following form:

$$T_{2,D} = \{(\phi^{F_A:D}(l), c) : l \in E^\times, c \in F^\times\},$$

where $D \in F_A^\times - F_A^{\times 2}$ and $E = F_A(\sqrt{D})$. There are 3 possibilities for D :

- $D \in F^\times - AF^{\times 2}$; hence, E/F is Galois, biquadratic.
- $D \notin F^\times$, E/F cyclic of degree 4.
- $D \notin F^\times$, E/F non-Galois, with Galois closure $\tilde{E}$ of E over F . Then, the Galois group $\mathrm{Gal}(\tilde{E}/F)$ is equivalent to D_4 , the dihedral group of order 8.

Since H_2 is a quotient group of $\mathrm{GL}(2, F_A) \times F^\times$, there is no distinction here between conjugacy and stable conjugacy.

B.1.4. Norm Correspondence—Explicit Description. We now use the results in Section 2.4 to describe explicitly the norm correspondence between elliptic regular elements.

The groups of $\bar{F}$ -points of $\mathbf{H}_1, \mathbf{H}_2$ are as follows:

$$\mathbf{H}_1(\bar{F}) = \{(g, \alpha, \beta) \in \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times \times \bar{F}^\times : \det g = \alpha\beta\},$$

$$\mathbf{H}_2(\bar{F}) = (\mathrm{GL}(2, \bar{F}) \times \mathrm{GL}(2, \bar{F}) \times \bar{F}^\times) / \{(\alpha I_2, \beta I_2, (\alpha\beta)^{-1}) : \alpha, \beta \in \bar{F}^\times\}.$$

Recall the definitions of N_1, N_2 :

$$N_1 : \mathbf{H}_1(\bar{F}) \ni (\mathrm{diag}(a, \lambda/a); b, \lambda/b) \mapsto \mathrm{diag}(b, a, \lambda/a, \lambda/b) \in \mathbf{G}(\bar{F}),$$

$$N_2 : \mathbf{H}_2(\bar{F}) \ni (\mathrm{diag}(a, b), \mathrm{diag}(c, d), 1) \mapsto \mathrm{diag}(ac, ad, bc, bd) \in \mathbf{G}(\bar{F}).$$

By definition, an element $\delta_i \in \mathbf{H}_i(F)$ ($i = 1, 2$) is a norm of an element $\delta \in \mathbf{G}(F)$ if δ_i is conjugate to a diagonal element t_i in $\mathbf{H}_i(\bar{F})$, and δ to a diagonal element t in $\mathbf{G}(\bar{F})$, such that $t = N_i(t_i)$.

For an algebraic group $\mathbf{H}$, and elements $g, h \in \mathbf{H}(\bar{F})$, we write $g \sim h$ if g, h are conjugate under $\mathbf{H}(\bar{F})$.

B.1.4.1. Matching $T_{1,A}$. Let σ be the generator of $\mathrm{Gal}(F_A/F)$. Put $\bar{\gamma} := \sigma\gamma$ for $\gamma \in F_A$.

B.5. CLAIM. *Let γ, ζ be elements in $F_A^\times - F^\times$ such that $N_{F_A/F}\gamma = N_{F_A/F}\zeta$. The element $\delta_1 = (\phi^A(\gamma), \zeta)$ in $T_{1,A}$ is a norm of $[\phi^A(\zeta), \phi^A(\gamma)] \in T_{1,A}$.*

PROOF. The element $(\phi^A(\gamma), \zeta)$ is conjugate to $(\mathrm{diag}(\gamma, \bar{\gamma}); \zeta, \bar{\zeta})$ in $\mathbf{H}_1(\bar{F})$. We have:

$$N_1 : (\mathrm{diag}(\gamma, \bar{\gamma}); \zeta, \bar{\zeta}) \mapsto \mathrm{diag}(\zeta, \gamma, \bar{\gamma}, \bar{\zeta}).$$

The element $\mathrm{diag}(\zeta, \gamma, \bar{\gamma}, \bar{\zeta})$ is conjugate to δ in $\mathbf{G}(\bar{F})$. □

It is clear from the proof of the claim that

$$(\phi^A(\gamma), \zeta), (\phi^A(\gamma), \bar{\zeta}), (\phi^A(\zeta), \gamma), (\phi^A(\zeta), \bar{\gamma}) \in T_{1,A}$$

are all norms of δ . If $\gamma \neq \zeta, \bar{\zeta}$, the elements listed above are not stably conjugate to one another. This is an example of a norm correspondence which is not one to one.

B.1.5. Matching $T_{1,AD}$. Let D be an element of $F^\times - AF^{\times 2}$. Recall our notation: $F_{AD} = F(\sqrt{AD})$, $F_D = F(\sqrt{D})$. Let E be the biquadratic extension $F(\sqrt{D}, \sqrt{A})$ of F .

Suppose $\mathrm{Gal}(E/F) = \langle \sigma, \tau \rangle$, such that $F_A = E^\tau$, $F_D = E^\sigma$, and $F_{AD} = E^{\sigma\tau}$. Here E^μ denotes the subfield of E fixed by $\mu \in \mathrm{Gal}(E/F)$. Thus, $\mathrm{Gal}(F_D/F) = \langle \tau \rangle = \langle \sigma\tau \rangle$, $\mathrm{Gal}(F_A/F) = \langle \sigma \rangle = \langle \tau\sigma \rangle$, and $\mathrm{Gal}(F_{AD}/F) = \langle \sigma \rangle = \langle \tau \rangle$.

B.6. CLAIM. *Let ζ be an element of $F_{AD}^\times - F^\times$, γ an element of $F_A^\times - F^\times$, such that $N_{F_{AD}/F}\zeta = N_{F_A/F}\gamma$. The element $\delta_1 = (\phi^{AD}(\zeta), \gamma)$ in $T_{1,AD}$ is a norm of $\delta = [\phi^A(\gamma), \phi^{AD}(\zeta)] \in T_{III,A,AD}$.*

PROOF. Consider δ_1, δ as elements in $\mathbf{H}_1(\bar{F}), \mathbf{G}(\bar{F})$, respectively. We have:

$$\delta_1 \sim (\mathrm{diag}(\zeta, \tau\zeta); \gamma, \sigma\gamma) \xrightarrow{N_1} \mathrm{diag}(\gamma, \zeta, \tau\zeta, \sigma\gamma) \sim \delta.$$

The element δ lies in $T_{III,A,AD}$ because

$$\det \phi^{AD}(\zeta) = N_{F_{AD}/F}\zeta = N_{F_A/F}\gamma = \det \phi^A(\gamma).$$

□

Observe that the elements $(\phi^{AD}(\zeta), \gamma)$ and $(\phi^{AD}(\zeta), \sigma\gamma)$ in $T_{1,AD}$, though not stably conjugate to each other, are both norms of $[\phi^A(\gamma), \phi^{AD}(\zeta)] \in T_{III,A,AD}$.

B.1.6. Matching $T_{2,D}$, Biquadratic Case. Let D be an element in $F^\times - AF^{\times 2}$. Then, $E = F_A(\sqrt{D})$ is a Galois, biquadratic extension of F . Suppose $\text{Gal}(E/F) = \langle \sigma, \tau \rangle$, such that $F_D = E^\sigma$, $E_A = E^\tau$, and $F_{AD} = E^{\sigma\tau}$. Then, $\text{Gal}(F_D/F) = \langle \tau \rangle = \langle \sigma\tau \rangle$, $\text{Gal}(F_A/F) = \langle \sigma \rangle = \langle \tau\sigma \rangle$, and $\text{Gal}(F_{AD}/F) = \langle \sigma \rangle = \langle \tau \rangle$.

B.7. CLAIM. For $c \in F^\times$ and $l \in E^\times$, the element $\delta_2 = (\phi^{F_A:D}(l), c)$ in $T_{2,D}$ is a norm of $\delta = [\phi^{F:D}(c\zeta'), \phi^{F:AD}(c\zeta)] \in T_{III,D,AD}$, where $\zeta' = N_{E/F_D}l = l\sigma l$ and $\zeta = N_{E/F_{AD}}l = l\sigma\tau l$.

PROOF. Recall that $\mathbf{H}_2(\bar{F})$ a quotient of $\text{GL}(2, \bar{F}) \times \text{GL}(2, \bar{F}) \times \bar{F}^\times$. The image of δ_2 in $\mathbf{H}_2(\bar{F})$ is represented by $(\phi^{F_A:D}(l), c\sigma\phi^{F_A:D}(l), 1)$.

Suppose $l = \alpha + \beta\sqrt{D}$ for some $\alpha, \beta \in F_A$. Since $D \in F^\times$ and $\sigma\sqrt{D} = \sqrt{D}$, the element $(\phi^{F_A:D}(l), c\sigma\phi^{F_A:D}(l), 1)$ is equal to

$$\left(\begin{pmatrix} \alpha & \beta D \\ \beta & \alpha \end{pmatrix}, c \begin{pmatrix} \sigma\alpha & \sigma\beta D \\ \sigma\beta & \sigma\alpha \end{pmatrix}, 1 \right) \sim (\text{diag}(l, \tau l), \text{diag}(c\sigma l, c\tau\sigma l), 1) \in \mathbf{H}_2(\bar{F}),$$

which maps via N_2 to

$$\begin{aligned} \text{diag}(cl\sigma l, cl\tau\sigma l, c\tau l \cdot \sigma l, \tau l \cdot \tau\sigma l) &= \text{diag}(c\zeta', c\zeta, c\tau\zeta, c\tau\zeta') \\ &\sim \delta = [\phi^{F:D}(c\zeta'), \phi^{F:AD}(c\zeta)]. \end{aligned}$$

Since $\det \phi^{F:D}(c\zeta') = c^2 N_{E/F}l = \det \phi^{F:AD}(c\zeta)$, the element δ lies in $T_{III,D,AD}$. $\square$

If $l \notin F_D^\times$, the regular elements $(\phi^{F_A:D}(l), c), (\sigma\phi^{F_A:D}(l), c) \in T_{2,D}$, which are not stably conjugate to each other, are both norms of the (possibly singular) element $[\phi^{F:D}(c\zeta'), \phi^{F:AD}(c\zeta)] \in T_{III,D,AD}$.

B.1.7. Matching $T_{2,D}$, Non-Biquadratic Case. Let D be an element in $F_A^\times - F_A^{\times 2}$. Let $\tilde{E}$ be the Galois closure of $E = F(\sqrt{D})$ over F . Let σ be the element of order 4 in $\text{Gal}(\tilde{E}/F)$ defined by $\sigma\sqrt{A} = -\sqrt{A}$, $\sigma\sqrt{D} = \sqrt{\sigma D}$, $\sigma^2\sqrt{D} = -\sqrt{D}$, $\sigma^3\sqrt{D} = -\sqrt{\sigma D}$. In particular, $\text{Gal}(F_A/F)$ is generated by the restriction of σ to F_A .

B.8. CLAIM. For $c \in F^\times$ and $l \in E^\times$, the element $\delta_2 = (\phi^{F_A:D}(l), c)$ in $T_{2,D}$ is a norm of $\delta = \phi^{F:A:D}(cl\sigma l) \in T_{IV,A,D}$.

PROOF. The image of δ_2 in $\mathbf{H}_2(\bar{F})$ is represented by $(\phi^{F_A:D}(l), c\sigma\phi^{F_A:D}(l), 1)$. Suppose $l = \alpha + \beta\sqrt{D}$ for some $\alpha, \beta \in F_A$. Then, δ_2 is equal to

$$\left(\begin{pmatrix} \alpha & \beta D \\ \beta & \alpha \end{pmatrix}, c \begin{pmatrix} \sigma\alpha & \sigma\beta\sigma D \\ \sigma\beta & \sigma\alpha \end{pmatrix}, 1 \right) \sim (\text{diag}(l, \sigma^2 l), \text{diag}(c\sigma l, c\sigma^3 l), 1) \in \mathbf{H}_2(\bar{F}).$$

We have:

$$\begin{aligned} N_2 : (\text{diag}(l, \sigma^2 l), \text{diag}(c\sigma l, c\sigma^3 l), 1) &\mapsto \text{diag}(cl\sigma l, cl\sigma^3 l, c\sigma^2 l \cdot \sigma l, c\sigma^2 l \cdot \sigma^3 l) \\ &\sim \delta = \phi^{F:A:D}(cl\sigma l). \end{aligned}$$

Since $N_{E/F_A}l\sigma l = l\sigma l \cdot \sigma^2 l \cdot \sigma^3 l \in F_A^\sigma = F$, δ lies in $T_{IV,A,D}$. $\square$

If $l \notin F_D^\times$, the elements $(\phi^{F_A:D}(l), c)$ and $(\sigma\phi^{F_A:D}(l), c) \in T_{2,D}$ are not stably conjugate to each other, but they are both norms of the element $\phi^{F:A:D}(cl\sigma l)$ in $T_{IV,A,D}$.

B.1.7.1. Summary. In summary, the norm correspondence between the stable conjugacy classes of the elliptic regular elements is as follows (bar denotes action of σ):

(1) Case of H_1

$$T_{1,A} \ni \left\{ \begin{array}{l} (\phi^A(\gamma), \zeta), \\ (\phi^A(\gamma), \bar{\zeta}), \\ (\phi^A(\zeta), \gamma), \\ (\phi^A(\zeta), \bar{\gamma}) \end{array} \right\} \longleftrightarrow [\phi^A(\gamma), \phi^A(\zeta)] \in T_{1,A}.$$

$$T_{1,AD} \ni \left\{ \begin{array}{l} (\phi^{AD}(\zeta), \gamma), \\ (\phi^{AD}(\zeta), \bar{\gamma}) \end{array} \right\} \longleftrightarrow [\phi^A(\gamma), \phi^{AD}(\zeta)] \in T_{III,A,AD}.$$

(2) Case of H_2

$$D \in F^\times - AF^{\times 2};$$

$$T_{2,D} \ni \left\{ \begin{array}{l} (\phi^{F_A:D}(l), c), \\ (\sigma \phi^{F_A:D}(l), c) \end{array} \right\} \longleftrightarrow [\phi^D(cl\sigma l), \phi^{AD}(cl\sigma \tau l)] \in T_{III,D,AD}.$$

$$D \in F_A^\times - F_A^{\times 2};$$

$$T_{2,D} \ni \left\{ \begin{array}{l} (\phi^{F_A:D}(l), c), \\ (\sigma \phi^{F_A:D}(l), c) \end{array} \right\} \longleftrightarrow \phi^{F:A:D}(cl\sigma l) \in T_{IV,A,D}.$$

B.2. Comparison of Orbital Integrals

Recall that ε is the nontrivial, unramified, quadratic character of $F^\times$, and F_A is the quadratic, unramified extension of F . It corresponds to ε via local class field theory. We fix a uniformizer ϖ in the ring of integers R of F . Since F_A/F is unramified, ϖ is also a uniformizer for F_A .

Let $K = \mathbf{G}(R)$, $K_i = \mathbf{H}_i(R)$ ($i = 1, 2$).

For $\mathbf{H} = \mathbf{G}$, $\mathbf{H}_1$, or $\mathbf{H}_2$, fix a pair $(\mathbf{B}, \mathbf{T})$, where $\mathbf{T}$ is the maximal diagonal torus of $\mathbf{H}$, and $\mathbf{B}$ is the upper triangular Borel subgroup of $\mathbf{H}$ containing $\mathbf{T}$. Let Δ be the set of roots of $\mathbf{H}$ with respect to $(\mathbf{B}, \mathbf{T})$. We define the **discriminant** $D_H(\delta)$ of an element $\delta \in \mathbf{H}(F)$ as follows:

$$D_H(\delta) := \prod_{\alpha \in \Delta} |\alpha(t_\delta) - 1|_{\bar{F}}.$$

Here, $|\cdot|_{\bar{F}}$ is the unique absolute value on $\bar{F}$ extending the normalized p -adic absolute value on F , and t_δ is an element in $\mathbf{T}(\bar{F})$ which is conjugate to δ in $\mathbf{H}(\bar{F})$. The discriminant $D_H(\delta)$ is independent of the choice of t_δ . Note that $D_H(\delta) = 0$ if δ is not regular.

For a p -adic field k with ring of integers R_k , each element $x \in k^\times$ may be written as $x = u \varpi_k^{\text{ord}_k x}$, where u is a unit in R_k , ϖ_k is a uniformizer of k , and $\text{ord}_k x$ is an integer. The function $\text{ord}_k : k^\times \rightarrow \mathbb{Z}$ is independent of the choice of ϖ_k . Let q be the cardinality of the residue field $R/(\varpi)$ of F . Put $\text{ord} := \text{ord}_F$. Let R_{F_A} be the ring of integers of F_A .

For elements g, h in a group, put

$$\text{Int}(g)h := ghg^{-1}.$$

B.2.1. Comparison for $T_{1,A}, T_{1,A}$.

B.2.1.1. Stable Orbital Integral— $T_{1,A}$.

B.9. CLAIM. Let ζ, ξ be elements in $R_{F_A}^\times - R^\times$ such that $N_{F_A/F}\zeta = N_{F_A/F}\xi$. Suppose $\zeta = a + b\sqrt{A}$, where $a, b \in F$. Let $\delta_1 = (\phi^A(\zeta), \xi) \in T_{1,A}$. The following equality holds:

$$SO_{\delta_1}(1_{K_1}) = \frac{q+1}{q-1} q^{\text{ord } b} - \frac{2}{q-1}.$$

PROOF. Let $\mathrm{GL}(2, F)^{F_A}$ be the subgroup of elements in $\mathrm{GL}(2, F)$ whose determinants lie in $N_{F_A/F} F_A^\times$. Let $K' = \mathrm{GL}(2, R)$. Since F_A/F is unramified, K' is a maximal compact subgroup in $\mathrm{GL}(2, F)^{F_A}$. The group $K_1 = \mathbf{H}_1(R)$ is then equal to $(K' \times R_{F_A}^\times)' := \{(g, \gamma) \in K' \times R_{F_A}^\times : \det g = N_{F_A/F} \gamma\}$.

For $\rho \in F^\times$, put $\phi_\rho^A(\alpha) := \mathrm{Int}(\mathrm{diag}(1, \rho^{-1}))\phi^A(\alpha)$. There are two conjugacy classes within the stable conjugacy class of δ_1 . One is represented by $\delta_1^1 := \delta_1$, and the other by $\delta_1^\varpi := (\phi_\varpi^A(\zeta), \xi)$.

For $\rho \in \{1, \varpi\}$, put

$$T_1^\rho := Z_{H_1}(\delta_1^\rho) = \{(\phi_\rho^A(\alpha), \beta) : N_{F_A/F} \alpha = N_{F_A/F} \beta\}.$$

Put $T_1 := T_1^1 = T_{1,A}$. Let dh denote the fixed Haar measure dh_1 on H_1 .

By definition,

$$SO_{\delta_1}(1_{K_1}) = \sum_{\rho \in \{1, \varpi\}} \int_{T_1^\rho \backslash H_1} 1_{K_1}(h^{-1} \delta_1^\rho h) dh.$$

The quotient $T_1^\rho \backslash H_1$ is equal to $T_A^\rho \backslash \mathrm{GL}(2, F)^{F_A}$, where $T_A^\rho := \{\phi_\rho^A(\gamma) : \gamma \in F_A^\times\}$.

From [F3, p. 30], we have the decomposition

$$\mathrm{GL}(2, F) = \dot{\bigcup}_{j \geq 0} T_A^\rho \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j-\mathrm{ord} \rho} \end{pmatrix} K',$$

where the disjoint union is over all nonnegative integers j . Since K' is a subgroup of $\mathrm{GL}(2, F)^{F_A}$, and $\varpi \notin N_{F_A/F} F_A^\times$, the above decomposition implies that

$$\mathrm{GL}(2, F)^{F_A} = \dot{\bigcup}_{j-\mathrm{ord} \rho \text{ even} \geq 0} T_A^\rho \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j-\mathrm{ord} \rho} \end{pmatrix} K'.$$

Let δ_1' be the $\mathrm{GL}(2, F)$ -component $\phi^A(\zeta)$ of δ_1 . Put $\delta_1'^\varpi := \phi_\varpi^A(\zeta)$. For any $h = (h', \gamma) \in H_1$, the element $\mathrm{Int}(h^{-1}) \delta_1$ lies in K_1 if and only if $\mathrm{Int}(h'^{-1}) \delta_1'$ lies in K' . The stable orbital integral $SO_{\delta_1}(1_{K_1})$ is therefore equal to

$$\begin{aligned} & \int_{T_A \backslash \mathrm{GL}(2, F)^{F_A}} 1_{K'}(\mathrm{Int}(h'^{-1}) \delta_1') dh' + \int_{T_A^\varpi \backslash \mathrm{GL}(2, F)^{F_A}} 1_{K'}(\mathrm{Int}(h'^{-1}) \delta_1'^\varpi) dh' \\ &= \sum_{j \text{ even} \geq 0} \int_{K' \cap \mathrm{Int}(\mathrm{diag}(1, \varpi^{-j})) T_A \backslash K'} 1_{K'}(\mathrm{Int}(k^{-1} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-j} \end{pmatrix}) \delta_1') dk \\ & \quad + \sum_{j \text{ odd} \geq 0} \int_{K' \cap \mathrm{Int}(\mathrm{diag}(1, \varpi^{1-j})) T_A^\varpi \backslash K'} 1_{K'}(\mathrm{Int}(k^{-1} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-j} \end{pmatrix}) \delta_1') dk. \end{aligned}$$

Here, dk is the Haar measure on K' such that K' has volume one.

For a nonnegative integer j , let

$$R_{F_A}(j) = \{a + b\sqrt{A} : a, b \in F; |a| \leq 1, |b| \leq |\varpi^j| = q^{-j}\}.$$

Then, the group $K' \cap \mathrm{Int}(\begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-j} \end{pmatrix}) T_A$ is isomorphic to $R_{F_A}(j)^\times$.

Hence, for $\rho \in \{1, \varpi\}$ we have:

$$[K' \cap T_A : K' \cap \mathrm{Int}(\begin{pmatrix} 1 & 0 \\ 0 & \varpi^{\mathrm{ord} \rho - j} \end{pmatrix}) T_A^\rho] = [R_{F_A}^\times : R_{F_A}(j)^\times].$$

Consequently,

$$\begin{aligned}
 (B.1) \quad SO_{\delta_1}(1_{K_1}) &= \sum_{j \text{ even } \geq 0} [R_{F_A}^\times : R_{F_A}(j)^\times] 1_{K'} \left(\begin{pmatrix} a & bA\varpi^j \\ b\varpi^{-j} & a \end{pmatrix} \right) \\
 &+ \sum_{j \text{ odd } \geq 0} [R_{F_A}^\times : R_{F_A}(j)^\times] 1_{K'} \left(\begin{pmatrix} a & bA\varpi^j \\ b\varpi^{-j} & a \end{pmatrix} \right) \\
 &= \sum_{j \geq 0} [R_{F_A}^\times : R_{F_A}(j)^\times] 1_{K'} \left(\begin{pmatrix} a & bA\varpi^j \\ b\varpi^{-j} & a \end{pmatrix} \right).
 \end{aligned}$$

Since F_A/F is unramified, the cardinality q_{F_A} of the residue field $R_{F_A}/(\varpi)$ is q^2 . The index $[R_{F_A}^\times : R_{F_A}(j)^\times]$ is therefore equal to

$$\begin{aligned}
 [R_{F_A}^\times : 1 + \varpi^j R_{F_A}] / [R_{F_A}(j)^\times : 1 + \varpi^j R_{F_A}] \\
 = \frac{(q_{F_A} - 1)q_{F_A}^{j-1}}{(q - 1)q^{j-1}} = \begin{cases} 1 & \text{if } j = 0, \\ (q + 1)q^{j-1} & \text{if } j > 0. \end{cases}
 \end{aligned}$$

Since $1_{K'} \left(\begin{pmatrix} a & bA\varpi^j \\ b\varpi^{-j} & a \end{pmatrix} \right)$ is zero unless $j \leq \text{ord } b$, we conclude that

$$SO_{\delta_1}(1_K) = 1 + \frac{q+1}{q} \sum_{j=1}^{\text{ord } b} q^j = \frac{q+1}{q-1} q^{\text{ord } b} - \frac{2}{q-1}.$$

□

B.2.1.2. Twisted Orbital Integral— $T_{I,A}$.

Notation:

- $\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} x & y \\ z & t \end{pmatrix} \right] := \begin{pmatrix} a & x & y & b \\ c & z & t & d \end{pmatrix}.$
- $C_0 := \{[g_1, g_2] : g_1, g_2 \in \text{GL}(2, F); \det g_1 = \det g_2\} \subset G.$
- $C'_0 = (\text{GL}(2, F) \times \text{GL}(2, F))'$
 $:= \{(g_1, g_2) \in \text{GL}(2, F) \times \text{GL}(2, F) : \det g_1 = \det g_2\}.$
- $T_A := \{\phi^A(\xi) : \xi \in F_A^\times\}.$
- For $\rho \in F^\times$, put $T_A^\rho := \text{Int}(\text{diag}(1, \rho^{-1}))T_A$, and
 $T'_\rho := (T_A \times T_A^\rho)' := \{(t_1, t_2) \in T_A \times T_A^\rho : \det t_1 = \det t_2\} \subset C'_0.$
- For $\rho \in F^\times$, put $\phi_\rho^A(\gamma) := \text{Int}(\text{diag}(1, \rho^{-1}))\phi^A(\gamma)$. Put
 $T_\rho := T_{I,A}^\rho := \{[\phi^A(\alpha), \phi_\rho^A(\beta)] : N_{F_A/F}\alpha = N_{F_A/F}\beta\} \subset G.$
- Put $K_0 := (\text{GL}(2, R) \times \text{GL}(2, R))'$, where the prime denotes equal determinants.
- For an integer m , put $K_m := \{(g, h) \in C'_0 : g - ehe \in \varpi^m M_2(R)\}$. Here, $M_2(R)$ denotes the set of all 2×2 matrices with coefficients in R , and $e := \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \in \text{GL}(2, F).$

Let ξ, ζ be elements in $R_{F_A}^\times - R^\times$ such that $\xi \neq \zeta$ or $\sigma\zeta$, and $N_{F_A/F}\xi = N_{F_A/F}\zeta$. Let δ be the regular element $[\phi^A(\xi), \phi^A(\zeta)]$ in $T_{I,A}$. Let δ_ϖ be the element $[\phi^A(\xi), \phi_\varpi^A(\zeta)]$. It is stably conjugate but not conjugate to δ . We assume that δ is topologically unipotent.

Suppose $\xi = a_1 + b_1\sqrt{A}$, $\zeta = a_2 + b_2\sqrt{A}$, where a_i, b_i ($i = 1, 2$) are elements in F , with $b_i \neq 0$. Let $N_i = \text{ord } b_i$. Let $X = \text{ord}(a_1 - a_2)$.

B.10. CLAIM. *The sum $O_\delta(1_K) + O_{\delta\varpi}(1_K)$ of ε -twisted orbital integrals is equal to*

$$(-q)^{N_1}(-q)^X \left[\frac{(q+1)q^{N_2} - 2}{q-1} \right].$$

PROOF. We make use of the following decomposition of G ([F3, Lemma I. J. 1], see also [We]):

$$\mathrm{GSp}(2, F) = \dot{\bigcup}_{m \geq 0} C_0 z(m) K.$$

Here, the disjoint union is over nonnegative integers m , and

$$z(m) := \begin{pmatrix} 1 & 0 & \varpi^{-m} & 0 \\ 0 & 1 & 0 & \varpi^{-m} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Put $K_m^{C_0} := C_0 \cap z(m) K z(m)^{-1}$. For $\rho \in \{1, \varpi\}$, put $\delta_\rho := [\phi^A(\xi), \phi_\rho^A(\zeta)]$. we have:

$$\begin{aligned} O_{\delta_\rho}(1_K) &= \int_{T_\rho \backslash G} 1_K(g^{-1} \delta_\rho g) \varepsilon(g) dg \\ &= \sum_{m \geq 0} |K| \int_{T_\rho \backslash C_0 / C_0 \cap z(m) K z(m)^{-1}} 1_K(z(m)^{-1} g^{-1} \delta_\rho g z(m)) \varepsilon(g z(m)) dg \\ &= \sum_{m \geq 0} \int_{T_\rho \backslash C_0 / C_0 \cap z(m) K z(m)^{-1}} 1_K(z(m)^{-1} g^{-1} \delta_\rho g z(m)) \varepsilon(g) dg \\ &= \sum_{m \geq 0} \int_{T_\rho \backslash C_0 / K_m^{C_0}} 1_{K_m^{C_0}}(g^{-1} \delta_\rho g) \varepsilon(g) dg. \end{aligned}$$

Note that ε is trivial on K because it is unramified. Also, $|K| = 1$ by our choice of measure.

Let

$$C'_0 = \{(g_1, g_2) \in \mathrm{GL}(2, F) \times \mathrm{GL}(2, F) : \det g_1 = \det g_2\}.$$

Let $e = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$, $w = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \in \mathrm{GL}(2, F)$. For an integer m , define an isomorphism $\phi_m : C'_0 \rightarrow C_0$ as follows:

$$\phi_m((g_1, g_2)) = \begin{pmatrix} 1 & & \\ & \varpi^m & \\ & & \varpi^m \end{pmatrix} [g_1, ew g_2 we] \begin{pmatrix} 1 & & \\ & \varpi^m & \\ & & \varpi^m \end{pmatrix}^{-1}.$$

For $\rho \in \{1, \varpi\}$, let $\delta'_\rho = (\phi^A(\xi), \phi_\rho^A(\zeta)) \in C'_0$. Let $T'_\rho = Z_{C'_0}(\delta'_\rho)$. For an integer $m \geq 0$, let $\delta_{\rho, m} = \phi_m^{-1}(\delta_\rho) \in C'_0$. Let $T_{\rho, m} = Z_{C'_0}(\delta_{\rho, m})$.

Let

$$K_m = \{(g_1, g_2) \in C'_0 : g_1 - e g_2 e \in \varpi^m M_2(R)\},$$

where $M_2(R)$ is the set of 2×2 matrices with coefficients in R . Then, ϕ_m maps K_m isomorphically onto $K_m^{C_0}$ ([F3, p. 38]).

Let $K_0 = (\mathrm{GL}(2, R) \times \mathrm{GL}(2, R))'$

$$:= \{(g_1, g_2) \in \mathrm{GL}(2, R) \times \mathrm{GL}(2, R) : \det g_1 = \det g_2\}.$$

Then, $[K : K_m^{C_0}] = [K_0 : K_m]$. We have:

$$(B.2) \quad O_{\delta_\rho}(1_K)$$

$$= \sum_{m \geq 0} [K_0 : K_m] \int_{T_{\rho, m} \setminus C'_0} 1_{\phi_m(K_m)} (\phi_m(g')^{-1} \delta_\rho \phi_m(g')) \varepsilon(\phi_m(g')) dg'.$$

By abuse of notation, for $g' = (g_1, g_2) \in C'_0$, let $\varepsilon(g')$ denote the value of $\varepsilon(\phi_m(g'))$. It is equal to $\varepsilon(\det g_1) = \varepsilon(\det g_2)$ and is independent of m .

We change variables $g' \mapsto ((\begin{smallmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{smallmatrix}) we) g'$. Since

$$\varepsilon\left(\left(\begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{pmatrix} we\right)\right) = \varepsilon(\det \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{pmatrix}) = \varepsilon(\varpi)^m = (-1)^m,$$

the expression (B.2) is equal to

$$(B.3) \quad \sum_{m \geq 0} [K_0 : K_m] \int_{T_{\rho'} \setminus C'_0} 1_{K_m}(g'^{-1} \delta'_\rho g') \varepsilon\left(\left(\begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{-m} \end{pmatrix} we\right) g'\right) dg' \\ = \sum_{m \geq 0} (-1)^m [K_0 : K_m] \int_{T_{\rho'} \setminus C'_0} 1_{K_m}(g'^{-1} \delta'_\rho g') \varepsilon(g') dg'.$$

For $\rho \in \{1, \varpi\}$, we have the following decomposition of C'_0 ([F3, p. 45]):

$$C'_0 = \bigcup_{r \in \mathcal{R}_\rho} T'_\rho \cdot r \cdot K_0,$$

where

$$\mathcal{R}_\rho := \left\{ r_{j_1, j_2, \epsilon} = \left(\varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_1} \end{pmatrix}, \phi_\rho^A(\epsilon') \varpi^{-\lfloor (j_2 - \text{ord } \rho)/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \epsilon \varpi^{j_2 - \text{ord } \rho} \end{pmatrix} \right) \right\}.$$

Here, (i) j_1, j_2 are nonnegative integers, with $j_1 - j_2 + \text{ord } \rho$ even; (ii) ϵ is a representative in $R^\times$ of a class in $R^\times / R^{\times 2}$; (iii) ϵ' is an element of $R_{F_A}^\times$ such that $N_{F_A/F} \epsilon' = \epsilon$.

Let $R_{T'_\rho} = K_0 \cap T'_\rho$. The sum (B.3) is then equal to

$$\sum_{m \geq 0} (-1)^m [K_0 : K_m] \sum_{r \in \mathcal{R}_\rho} [R_{T'_\rho} : T'_\rho \cap r K_0 r^{-1}] \varepsilon(r) \int_{K_0} 1_{K_m}(k^{-1} r^{-1} \delta'_\rho r k) dk.$$

For $r = r_{j_1, j_2, \epsilon} \in \mathcal{R}_\rho$, we have

$$\varepsilon(r) = \varepsilon\left(\left(\varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_1} \end{pmatrix}, \phi_\rho^A(\epsilon') \varpi^{-\lfloor (j_2 - \text{ord } \rho)/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \epsilon \varpi^{j_2 - \text{ord } \rho} \end{pmatrix}\right)\right) \\ = \varepsilon\left(\det\left(\varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_1} \end{pmatrix}\right)\right) = (-1)^{j_1}.$$

Thus, $O_{\delta_\rho}(1_K)$ is equal to

$$(B.4) \quad \sum_{m \geq 0} (-1)^m [K_0 : K_m] \sum_{\substack{j_1, j_2 \geq 0; \\ j_1 - j_2 + \text{ord } \rho \text{ even}}} (-1)^{j_1} [R_{T'_\rho} : T'_\rho \cap r K_0 r^{-1}] \\ \cdot \int_{K_0} 1_{K_m}(k^{-1} r^{-1} \delta'_\rho r k) dk.$$

Hence, the sum describing the ε -twisted orbital integral $O_{\delta_\rho}(1_K)$ is quite similar to that describing its nontwisted counterpart in [F3]. They are identical but for the factors $(-1)^m, (-1)^{j_1}$ which appear above in the ε -twisted case.

The quantities $[R_{T'_\rho} : T'_\rho \cap r K_0 r^{-1}]$ and $[K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1} r^{-1} \delta'_\rho r k) dk$ are computed in [F3, Sect. II. C]. Recall that $\xi = a_1 + b_1 \sqrt{A}$, $\zeta = a_2 + b_2 \sqrt{A}$, and $N_i = \text{ord } b_i$ ($i = 1, 2$), $X = \text{ord}(a_1 - a_2)$. We follow the notation in [F3] and let $N = \min(N_1, N_2)$,

$\nu_i = N_i - j_i$. Let $\delta(m = 0)$ be 0 if $m \neq 0$, and 1 if $m = 0$. Let $\delta(m \geq 1)$ be 0 if $m < 1$, and 1 if $m \geq 1$. It follows from [F3, Sect. II. C] that $O_\delta(1_K) + O_{\delta_\varpi}(1_K)$ is the sum of the following three quantities:

I.

$$(B.5) \quad \sum_{m=0}^N (-1)^m (\delta(m=0) + \delta(m \geq 0)(1 - q^{-2})q^{3m})$$

$$\left[\sum_{\nu_1=m}^{N_1-1} \sum_{\nu_2=m}^{N_2-1} \left(\frac{q+1}{q} \right)^2 (-q)^{N_1-\nu_1} q^{N_2-\nu_2} \right. \\ \left. + \sum_{\nu_1=m}^{N_1-1} \frac{q+1}{q} (-q)^{N_1-\nu_1} + \sum_{\nu_2=m}^{N_2-1} \frac{q+1}{q} q^{N_2-\nu_2} + 1 \right],$$

II.

$$(B.6) \quad \sum_{0=\nu}^{N-1} \sum_{m=\nu+1}^{2N-\nu} (-1)^m \frac{1}{2} \left(\frac{q+1}{q} \right)^2 (-q)^{N_1-\nu} q^{N_2-\nu} 2q^{m+2\nu},$$

III.

$$(B.7) \quad \sum_{m=N+1}^{X-N} (-1)^m \frac{q+1}{q} q^{2N+m} +$$

$$\sum_{\nu=0}^{N-1} \sum_{m=2N-\nu+1}^{X-\nu} (-1)^m \frac{1}{2} \left(\frac{q+1}{q} \right)^2 (-q)^{N-\nu} q^{N-\nu} 2q^{m+2\nu}.$$

The sum I + II + III has the same form as the sum on [F3, p. 60] (even though they describe orbital integrals for different groups). There, it is computed to be

$$(-q)^{N_1} (-q)^X \left[\frac{((q+1)q^{N_2} - 2)}{q-1} \right].$$

The proof of Claim B.10 is now complete. $\square$

B.2.1.3. Transfer Factor— $T_{1,A}, T_{1,A}$. Let ξ, ζ be distinct elements in $F_A^\times - F^\times$ such that $\xi \neq \zeta, \sigma\zeta$. Let δ be the element $[\phi^A(\xi), \phi^A(\zeta)]$ in $T_{1,A} \subset G$. Let $\delta_1 = (\phi^A(\xi), \zeta) \in T_{1,A} \subset H_1$. Then, δ is regular and δ_1 is a norm of δ .

There are two conjugacy classes in the stable conjugacy class of δ . One is represented by δ , and the other by $\delta_\varpi := [\phi^A(\xi), \phi_\varpi^A(\zeta)]$. Since we are proving the matching of orbital integrals for the unit elements in the Hecke algebras, we may assume without loss of generality that $\delta_1 \in K_1$ and $\delta \in K$, or in other words $\xi, \zeta \in R_{F_A}^\times - R^\times$.

Extend ε to $F_A^\times$ by letting $\varepsilon(x) = (-1)^{\text{ord}_{F_A} x}$ for all $x \in F_A^\times$. We define the transfer factors for the pairs (δ_1, δ) and $(\delta_1, \delta_\varpi)$ as follows:

$$\Delta(\delta_1, \delta) = \Delta(\delta_1, \delta_\varpi) = \varepsilon \left(\left((\xi + \sigma\xi) - (\zeta + \sigma\zeta) \right) \left(\frac{\xi - \sigma\xi}{2\sqrt{A}} \right) \right) \left(\frac{D_G(\delta)}{D_{H_1}(\delta_1)} \right)^{1/2}.$$

We do not explain here why the above expression constitutes a transfer factor. For those interested in further details, [LS] and [H2] are both useful sources on transfer factors.

Suppose $\xi = a_1 + b_1\sqrt{A}$, $\zeta = a_2 + b_2\sqrt{A}$, where $a_i, b_i \in F$ ($i = 1, 2$). Let $N_i = \text{ord } b_i$. Let $X = \text{ord}(a_1 - a_2)$.

B.11. CLAIM. *The following holds:*

$$\Delta(\delta_1, \delta) = \Delta(\delta_1, \delta_\varpi) = (-q)^{-N_1 - X}.$$

PROOF. The proof is a straightforward computation, and we skip it. $\square$

B.12. PROPOSITION. *The following identity holds:*

$$SO_{\delta_1}(1_{K_1}) = \Delta(\delta_1, \delta)O_\delta(1_K) + \Delta(\delta_1, \delta_\varpi)O_{\delta_\varpi}(1_K).$$

PROOF. This follows from Claims B.9, B.10, and B.11. $\square$

Let $\tilde{\delta}_1 = (\phi^A(\xi), \zeta) \in T_{1,A}$, i.e. we swap ξ and ζ in δ_1 . Then, $\tilde{\delta}_1$ is not stably conjugate to δ_1 , but it is also a norm of $\delta = [\phi^A(\xi), \phi^A(\zeta)]$. We define the transfer factors for the pairs $(\tilde{\delta}_1, \delta)$ and $(\tilde{\delta}_1, \delta_\varpi)$ as follows:

$$\Delta(\tilde{\delta}_1, \delta) = \Delta(\delta_1, \delta), \quad \Delta(\tilde{\delta}_1, \delta_\varpi) = -\Delta(\delta_1, \delta).$$

B.13. PROPOSITION. *The following holds:*

$$SO_{\tilde{\delta}_1}(1_{K_1}) = \sum_{\rho \in \{1, \varpi\}} \Delta(\tilde{\delta}_1, \delta_\rho)O_{\delta_\rho}(1_K).$$

PROOF. Put $O_\delta^{\text{us}}(1_K) := O_\delta(1_K) - O_{\delta_\varpi}(1_K)$. Thus,

$$\sum_{\rho \in \{1, \varpi\}} \Delta(\tilde{\delta}_1, \delta_\rho)O_{\delta_\rho}(1_K) = \Delta(\tilde{\delta}_1, \delta)O_\delta^{\text{us}}(1_K).$$

It follows from (B.4) that $O_\delta^{\text{us}}(1_K)$ is equal to

$$\begin{aligned} &= \sum_{\substack{r_{j_1, j_2, \epsilon}; \\ j_1 + j_2 \text{ even}}} \varepsilon(r)[R_{T'} : T' \cap rK_0r^{-1}] \sum_{m \geq 0} (-1)^m [K_0 : K_m] \\ &\quad \cdot \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk \\ &- \sum_{\substack{r_{j_1, j_2, \epsilon}; \\ j_1 + j_2 \text{ odd}}} \varepsilon(r)[R_{T'_\varpi} : T'_\varpi \cap rK_0r^{-1}] \sum_{m \geq 0} (-1)^m [K_0 : K_m] \\ &\quad \cdot \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'_\varpi rk) dk \\ &= \sum_{\rho \in \{1, \varpi\}} \sum_{\substack{r_{j_1, j_2, \epsilon}; \\ j_1 + j_2 + \text{ord } \rho \text{ even}}} (-1)^{j_1 + j_2} \varepsilon(r)[R_{T'_\rho} : T'_\rho \cap rK_0r^{-1}] \sum_{m \geq 0} (-1)^m [K_0 : K_m] \\ &\quad \cdot \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'_\rho rk) dk. \end{aligned}$$

Hence, the quantity $O_\delta^{\text{us}}(1_K)$ is obtained from $\sum_{\rho \in \{1, \varpi\}} O_{\delta_\rho}(1_K)$ on multiplying each summand in the expressions (B.5) through (B.7) by

$$(-1)^{j_1 + j_2} = (-1)^{N_1 - \nu_1 + N_2 - \nu_2}.$$

In other words, $O_\delta^{\text{us}}(1_K)$ is equal to the sum of the following three quantities:

I.

$$(B.8) \quad \sum_{m=0}^N (-1)^m (\delta(m=0) + \delta(m \geq 0)(1 - q^{-2})q^{3m})$$

$$\left[\sum_{\nu_1=m}^{N_1-1} \sum_{\nu_2=m}^{N_2-1} \left(\frac{q+1}{q} \right)^2 q^{N_1-\nu_1} (-q)^{N_2-\nu_2} \right. \\ \left. + \sum_{\nu_1=m}^{N_1-1} \frac{q+1}{q} q^{N_1-\nu_1} + \sum_{\nu_2=m}^{N_2-1} \frac{q+1}{q} (-q)^{N_2-\nu_2} + 1 \right],$$

II.

$$(B.9) \quad \sum_{0=\nu}^{N-1} \sum_{m=\nu+1}^{2N-\nu} (-1)^m \frac{1}{2} \left(\frac{q+1}{q} \right)^2 q^{N_1-\nu} (-q)^{N_2-\nu} 2q^{m+2\nu},$$

III.

$$(B.10) \quad \sum_{m=N+1}^{X-N} (-1)^m \frac{q+1}{q} q^{2N+m} + \\ \sum_{\nu=0}^{N-1} \sum_{m=2N-\nu+1}^{X-\nu} (-1)^m \frac{1}{2} \left(\frac{q+1}{q} \right)^2 q^{N-\nu} (-q)^{N-\nu} 2q^{m+2\nu}.$$

The sum of the above quantities is the same as $\sum_{\rho \in \{1, \varpi\}} O_{\tilde{\delta}_\rho}(1_K)$, where

$$\tilde{\delta} = [\phi^A(\zeta), \phi^A(\xi)],$$

i.e., δ with the two $\mathrm{GL}(2, F)$ -components swapped.

By Claims B.10 and B.11, we have

$$O_\delta^{\mathrm{us}}(1_K) = \sum_{\rho \in \{1, \varpi\}} O_{\tilde{\delta}_\rho}(1_K) = (-q)^{N_2+X} \left[\frac{((q+1)q^{N_1} - 2)}{q-1} \right],$$

$$\Delta(\tilde{\delta}_1, \delta) = (-q)^{-N_2-X}.$$

By Claim B.9, the stable orbital integral $SO_{\tilde{\delta}_1}(1_{K_1})$ is equal to

$$\frac{q+1}{q-1} q^{N_1} - \frac{2}{q-1}.$$

The proposition follows. $\square$

REMARK: Since $\delta = [\phi^A(\xi), \phi^A(\zeta)]$ is conjugate to $\tilde{\delta} = [\phi^A(\zeta), \phi^A(\xi)]$, it seems contradictory that $\sum_{\rho \in \{1, \varpi\}} O_{\delta_\rho}(1_K)$ should differ from $\sum_{\rho \in \{1, \varpi\}} O_{\tilde{\delta}_\rho}(1_K)$. However, note that we have chosen $[\phi^A(\xi), \phi_\varpi^A(\zeta)]$ to be the representative of the conjugacy class, different from that of δ , in the stable conjugacy class of δ . In other words, the choice of the representative involves a bias towards the second component of $[\phi^A(\xi), \phi^A(\zeta)]$. More concretely, there exists an element $g \in G$ such that $\tilde{\delta} = g^{-1}\delta g$, but $[\phi^A(\zeta), \phi_\varpi^A(\xi)]$ is not conjugate to $[\phi^A(\xi), \phi_\varpi^A(\zeta)]$ via the same g .

B.2.2. Comparison for $T_{1,AD}$, $T_{III,A,AD}$. Let D be an element in $F^\times - AF^{\times 2} \cup F^{\times 2}$. Note that since by assumption F_A/F is unramified, A lies in $R^\times - R^{\times 2}$. Let $E = F_A(\sqrt{D}) = F(\sqrt{A}, \sqrt{D})$. Suppose $\text{Gal}(E/F) = \langle \sigma, \tau \rangle$, such that $F_A = E^\tau$, $F_{AD} = E^\sigma$, and $\text{Gal}(F_A/F) = \langle \sigma \rangle$, $\text{Gal}(F_{AD}/F) = \langle \tau \rangle$. Since by assumption the residual characteristic of F is odd, the quadratic extensions F_D/F , F_{AD}/F are ramified.

For a p -adic field k , let R_k denote the ring of integers in k . Let ϖ be a uniformizer of F .

Since F_{AD}/F and F_D/F are ramified, without loss of generality we assume that $AD = -\varpi$, $D = -A^{-1}\varpi$. We let $\varpi_{AD} = \sqrt{-\varpi}$ be the uniformizer of F_{AD} , thus $N_{F_{AD}/F} \varpi_{AD} = \varpi$. (We may also let $\sqrt{-A^{-1}\varpi}$ be the uniformizer of F_D , but we do not use it in this section).

Let ξ be an element in $R_{F_A}^\times$, ζ an element in $R_{F_{AD}}^\times$, such that $\xi, \zeta \notin R^\times$, and $N_{F_A/F} \xi = N_{F_{AD}/F} \zeta$. Let

$$\begin{aligned} \delta_1 &= (\phi^{AD}(\zeta), \xi) \in T_{1,AD} \cap K_1, \\ \delta &= [\phi^A(\xi), \phi^{AD}(\zeta)] \in T_{III,A,AD} \cap K. \end{aligned}$$

Then, δ is regular, and δ_1 is a norm of δ . Recall from Section B.1 that the respective conjugacy classes of δ_1, δ are stable. We assume that δ is topologically unipotent.

Suppose $\xi = a_1 + b_1\sqrt{A}$, $\zeta = a_2 + b_2\sqrt{AD}$, where $a_i, b_i \in F$ ($i = 1, 2$). Let $N_i = \text{ord } b_i$. Let $X = \text{ord}(a_1 - a_2)$.

B.2.2.1. Stable Orbital Integral— $T_{1,AD}$.

B.14. CLAIM. *The stable orbital integral $O_{\delta_1}(1_{K_1})$ is equal to*

$$\frac{q^{N_2+1} - 1}{q - 1}.$$

PROOF. Recall that $\text{GL}(2, F)^{F_A}$ denotes $\{g \in \text{GL}(2, F) : \det g \in N_{F_A/F} F_A^\times\}$. Since the field extension F_A/F is unramified, $K' = \text{GL}(2, R)$ is a maximal compact subgroup in $\text{GL}(2, F)^{F_A}$.

Let

$$T_{AD}^{F_A} = \{\phi^{AD}(\gamma) : \gamma \in F_{AD}^\times; N_{F_{AD}/F} \gamma \in N_{F_A/F} F_A^\times\} \subset \text{GL}(2, F)^{F_A}.$$

For a nonnegative integer j , put $r_j = \phi^{AD}(\varpi_{AD}^{-j}) \begin{pmatrix} 1 & \\ & \varpi^j \end{pmatrix}$. Lemma I. I. 1 of [F3] implies that $\text{GL}(2, F)^{F_A}$ decomposes as follows:

$$(B.11) \quad \text{GL}(2, F)^{F_A} = \bigcup_{j \geq 0} T_{AD}^{F_A} \cdot r_j \cdot K'.$$

Let

$$R_{F_{AD}}(j) = \{a + b\sqrt{AD} : a, b \in F; |a| \leq 1, |b| \leq |\varpi^j| = q^{-j}\}.$$

We have $K' \cap r_j^{-1} T_{AD}^{FA} r_j = K' \cap r_j^{-1} T_{AD} r_j \cong R_{FAD}(j)^\times$, and $[R_{FAD}^\times : R_{FAD}(j)^\times]$ is equal to q^j for all nonnegative integers j ([F3, Sect. I. I.]). Hence,

$$\begin{aligned}
O_{\delta_1}(1_{K_1}) &= \int_{T_{1,AD} \setminus H_1} 1_{K_1}(h^{-1} \delta_1 h) dh \\
&= \int_{T_{AD}^{FA} \setminus \text{GL}(2, F)^{FA}} 1_{K'}(h^{-1} \phi^{AD}(\xi) h) dh \\
&= \sum_{j \geq 0} \int_{K' \cap \text{Int}(r_j^{-1}) T_{AD}^{FA} \setminus K'} 1_{K'}(\text{Int}(k^{-1} r_j^{-1}) \delta'_1) dk \\
&= \sum_{j \geq 0} [R_{FAD}^\times : R_{FAD}(j)^\times] 1_{K'} \left(\begin{pmatrix} 1 & \\ & \varpi^j \end{pmatrix}^{-1} \phi^{AD}(\xi) \begin{pmatrix} 1 & \\ & \varpi^j \end{pmatrix} \right) \\
&= \sum_{j \geq 0} q^j \cdot 1_{K'} \left(\begin{pmatrix} a & b^{AD} \varpi^j \\ b \varpi^{-j} & a \end{pmatrix} \right) = \sum_{j=0}^{N_2} q^j = \frac{q^{N_2+1} - 1}{q - 1}.
\end{aligned}$$

□

B.2.2.2. Twisted Orbital Integral— $T_{III,A,AD}$. Recall: $\xi = a_1 + b_1 \sqrt{A} \in R_{FA}^\times$, $\zeta = a_2 + b_2 \sqrt{AD} \in R_{FAD}^\times$, $N_i = \text{ord } b_i$ ($i = 1, 2$), $X = \text{ord}(a_1 - a_2)$. Recall that $\delta = [\phi^A(\xi), \phi^{AD}(\zeta)] \in T_{III,A,AD}$.

B.15. CLAIM. *The ε -twisted orbital integral $O_\delta(1_K)$ is equal to*

$$(-q)^{N_1+X} \left(\frac{q^{N_2+1} - 1}{q - 1} \right).$$

PROOF. As before, C'_0 denotes the group $(\text{GL}(2, F) \times \text{GL}(2, F))'$, and K_0 denotes $(\text{GL}(2, R) \times \text{GL}(2, R))'$, where the prime indicates equal determinants. Recall that $K_m := \{(g, h) \in C'_0 : g - ehe \in \varpi^m M_2(R)\}$, $e := \text{diag}(1, -1)$.

Let $\delta' = (\phi^A(\xi), \phi^{AD}(\zeta))$. Let

$$T' = Z_{C'_0}(\delta') = \{(\phi^A(\alpha), \phi^{AD}(\beta)) : \alpha \in F_A^\times, \beta \in F_{AD}^\times; N_{FA/F} \alpha = N_{FAD/F} \beta\}.$$

Repeating the procedure used to compute the orbital integral in the case of $T_{I,A}$, we obtain:

$$(B.12) \quad O_\delta(1_K) = \sum_{m \geq 0} (-1)^m [K_0, K_m] \int_{T' \setminus C'_0} 1_{K_m}(g^{-1} \delta' g) dg.$$

Recall that ϖ_{AD} is a uniformizer of F_{AD} such that $N_{FAD/F} \varpi_{AD} = \varpi$. From [F3, Sect. II. G], we have the decomposition

$$C'_0 = \bigcup_{r \in \mathcal{R}} T' \cdot r \cdot K_0.$$

Here, $\mathcal{R}$ is a set indexed by the couplets of nonnegative integers j_1, j_2 , and the classes in $R^\times / R^{\times 2}$. For a representative $\epsilon \in R^\times$ of a class in $R^\times / R^{\times 2}$, let ϵ' be an element in $F_A^\times$ such that $N_{FA/F} \epsilon' = \epsilon^{-1}$. For an integer j , let s_j be 0 if j is even, 1 if j is odd. The set $\mathcal{R}$ consists of the elements

$$r_{j_1, j_2, \epsilon} = \left(\phi^A(\epsilon') \varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \epsilon \varpi^{j_1} \end{pmatrix}, \phi^{AD}(\varpi_{AD}^{-j_2 + s_{j_1}}) \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_2} \end{pmatrix} \right) \in C'_0,$$

Let $R_{T'} = T' \cap K_0$. Then, (B.12) is equal to

$$(B.13) \quad \sum_{r \in \mathcal{R}} [R_{T'} : T' \cap r K_0 r^{-1}] \varepsilon(r) \sum_{m \geq 0} (-1)^m [K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1} r^{-1} \delta' r k) dk.$$

The quantities $[R_{T'} : T' \cap rK_0r^{-1}]$ and $[K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk$ are computed in [F3, p. 64]. Moreover,

$$\begin{aligned} \varepsilon \left(\phi^A(\epsilon') \varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \epsilon \varpi^{j_1} \end{pmatrix}, \phi^{AD}(\varpi_{AD}^{-j_2+s_{j_1}}) \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_2} \end{pmatrix} \right) \\ = \varepsilon \left(\det \left(\varpi^{-\lfloor j_1/2 \rfloor} \begin{pmatrix} 1 & 0 \\ 0 & \varpi^{j_1} \end{pmatrix} \right) \right) = (-1)^{j_1}. \end{aligned}$$

It follows that (B.13) has the same form as $1/2$ times the sum on page 78 of [F3], which is computed there to be

$$2(-q)^{N_1+X} \left(\frac{q^{N_2+1} - 1}{q - 1} \right).$$

□

B.2.2.3. Transfer Factor— $T_{1,AD}$, $T_{III,A,AD}$. Recall: $\xi = a_1 + b_1\sqrt{A} \in R_{F_A}^\times$, $\zeta = a_2 + b_2\sqrt{AD} \in R_{F_{AD}}^\times$, $N_i = \text{ord } b_i$ ($i = 1, 2$), $X = \text{ord}(a_1 - a_2)$. Recall that $\delta_1 = (\phi^{AD}(\zeta), \xi) \in T_{1,AD}$ and $\delta = [\phi^A(\xi), \phi^{AD}(\zeta)] \in T_{III,A,AD}$.

Recall that $E = F(\sqrt{A}, \sqrt{D})$. We extend ε to $E^\times$ by letting

$$\varepsilon(x) = (\sqrt{-1})^{\text{ord}_E x}, \quad \forall x \in E^\times.$$

We define the transfer factor $\Delta(\delta_1, \delta)$ as follows:

$$\varepsilon \left(\left((\xi + \sigma\xi) - (\zeta + \tau\zeta) \right) \cdot \left(\frac{\xi - \sigma\xi}{2\sqrt{A}} \right) \right) \left(\frac{D_G(\delta)}{D_{H_1}(\delta_1)} \right)^{1/2}.$$

B.16. PROPOSITION. *The following equality holds:*

$$O_{\delta_1}(1_{K_1}) = \Delta(\delta_1, \delta) O_\delta(1_K).$$

PROOF. This follows from Claims B.14, B.2.2.2, and the fact that

$$\Delta(\delta_1, \delta) = (-q)^{-N_1-X}.$$

□

B.2.3. Comparison for $T_{2,D}$, $T_{III,D,AD}$. Let D be an element in $F^\times - AF^{\times 2} \cup F^{\times 2}$. The quadratic extensions F_D/F , F_{AD}/F are ramified, for the residual characteristic of F is odd and F_A/F is unramified.

Let E be the biquadratic extension $F(\sqrt{A}, \sqrt{D})$ of F . The quadratic extensions E/F_D , E/F_{AD} are unramified. Suppose $\text{Gal}(E/F) = \langle \sigma, \tau \rangle$, such that $F_A = E^\tau$, $F_D = E^\sigma$, and $F_{AD} = E^{\sigma\tau}$. Thus, $\text{Gal}(F_A/F) = \langle \sigma \rangle = \langle \sigma\tau \rangle$, $\text{Gal}(F_D/F) = \langle \tau \rangle = \langle \tau\sigma \rangle$, and $\text{Gal}(F_{AD}/F) = \langle \sigma \rangle = \langle \tau \rangle$.

For a p -adic field k , let R_k denote the ring of integers in k . Let $R = R_F$.

Let l be an element in $R_E^\times - R^\times$ such that $N_{E/F_D}l, N_{E/F_{AD}}l \notin F^\times$. Let $\xi = N_{E/F_D}l$, $\zeta = N_{E/F_{AD}}l$. Let c be an element in $R^\times$.

Suppose $l = \alpha + \beta\sqrt{D}$ for some $\alpha, \beta \in F_A$. Put

$$\phi^{FA:D}(l) := \begin{pmatrix} \alpha & \beta D \\ \beta & \alpha \end{pmatrix} \in \text{GL}(2, F_A).$$

Recall that $K_2 = \mathbf{H}_2(R)$. Let

$$\delta_2 = (\phi^{FA:D}(l), c)_* \in T_{2,D} \cap K_2,$$

where $(\phi^{FA:D}(l), c)_*$ denotes the image of $(\phi^{FA:D}(l), c)$ in H_2 . It is a norm of the regular element

$$\delta = c [\phi^D(\xi), \phi^{AD}(\zeta)] \in T_{III,D,AD} \cap K.$$

The conjugacy classes of δ_2 and δ are stable. We assume that δ is topologically unipotent.

We depart from the previous notation and let q be the cardinality of the residue field of F_A . Let q_0 be the cardinality of the residue field of F . Since F_A/F is unramified, $q = q_0^2$.

Let $n = \text{ord}_{F_A}\beta$. Suppose $\xi = N_{E/F_D}l = a_1 + b_1\sqrt{D}$, and $\zeta = N_{E/F_{AD}}l = a_2 + b_2\sqrt{AD}$, where $a_i, b_i \in F$ ($i = 1, 2$). Let $N_i = \text{ord}_F b_i$. Let B_i be the unit in $R^\times$ such that $b_i = B_i\varpi^{N_i}$. Let $N = \min(N_1, N_2)$. Let $X = \text{ord}(a_1 - a_2)$.

B.2.3.1. *Stable Orbital Integral— $T_{2,D}$.*

B.17. CLAIM. *The stable orbital integral $O_{\delta_2}(1_{K_2})$ is equal to*

$$\frac{q^{N+1} - 1}{q - 1}.$$

PROOF. By [F3, Lemma I. I. 2], we have:

$$O_{\delta_2}(1_{K_2}) = \frac{q^{n+1} - 1}{q - 1}.$$

By [F3, Lemma II. L. 1], n is equal to N . □

B.2.3.2. *Twisted Orbital Integral— $T_{III,D,AD}$.* Let κ be the (unique) nontrivial quadratic character of the group $R^\times/R^{\times 2}$.

B.18. CLAIM. *The ε -twisted orbital integral $O_\delta(1_K)$ is equal to*

$$2\kappa(B_1B_2)(-1)^{N_1}q_0^{1+N_1+N_2}\left(\frac{q^{N+1} - 1}{q - 1}\right).$$

PROOF. Fix an element $\epsilon_0 \in R^\times - R^{\times 2}$. Without loss of generality, we may assume that $A = \epsilon_0$, $D = -\epsilon_0\varpi$, and $AD = -\varpi$. We fix uniformizers $\varpi_D = \sqrt{-\epsilon_0\varpi}$, $\varpi_{AD} = \sqrt{-\varpi}$ for F_D, F_{AD} , respectively. Thus, $N_{F_D/F}\varpi_D = \epsilon_0\varpi$, and $N_{F_{AD}/F}\varpi_{AD} = \varpi$.

Let

$$C'_0 = \{(g_1, g_2) \in \text{GL}(2, F)^2 : \det g_1 = \det g_2\},$$

$$K_0 = \{(k_1, k_2) \in \text{GL}(2, R)^2 : \det k_1 = \det k_2\}.$$

Let $K' = \text{GL}(2, R)$. For $D' = D, AD$, let

$$T_{D'} = \{\phi^{D'}(\gamma) : \gamma \in F_{D'}^\times\} \subset \text{GL}(2, F).$$

We have the decompositions

$$\text{GL}(2, F) = \bigcup_{j_1 \geq 0} T_D \begin{pmatrix} 1 & \\ & \varpi^{j_1} \end{pmatrix} K' = \bigcup_{j_2 \geq 0} T_{AD} \begin{pmatrix} 1 & \\ & \varpi^{j_2} \end{pmatrix} K'.$$

Hence, each element $g \in C'_0$ may be written as

$$g = (t_1, t_2) \cdot \left(\phi^D(\varpi_D^{-j_1}) \begin{pmatrix} 1 & \\ & (\epsilon_0\varpi)^{j_1} \end{pmatrix}, \phi^{AD}(\varpi_{AD}^{-j_2}) \begin{pmatrix} 1 & \\ & \varpi^{j_2} \end{pmatrix} \right) \cdot (k_1, k_2),$$

where $(t_1, t_2) \in T_D \times T_{AD}$, and $(k_1, k_2) \in \text{GL}(2, R)^2$, such that $\det(t_1k_2)$ is equal to $\det(t_2k_2)$. If $\det k_1 \neq \det k_2$, we may multiply k_1 by $\text{diag}(x, x)$ from T_D , for some $x \in R^\times$, such that $\det k_1$ and $\det k_2$ differ by $\epsilon_0 \in R^\times - R^{\times 2}$.

Let $s(\epsilon)$ be 0 if $\epsilon = 1$, 1 if $\epsilon = \epsilon_0$. Let $\mathcal{R}$ be the set of elements in C'_0 of the form:

$$r_{j_1, j_2, \epsilon} = \left(\phi^D(\varpi_D^{-j_1-s(\epsilon)}) \begin{pmatrix} 1 & \\ & \epsilon \cdot (\epsilon_0\varpi)^{j_1} \end{pmatrix}, \phi^{AD}(\varpi_{AD}^{-j_2-s(\epsilon)}) \begin{pmatrix} 1 & \\ & \varpi^{j_2} \end{pmatrix} \right),$$

where j_1, j_2 range over nonnegative integers, and ϵ over the set $\{1, \epsilon_0\}$. Then, C'_0 decomposes as follows:

$$(B.14) \quad C'_0 = \bigcup_{\epsilon \in \{1, \epsilon_0\}} \bigcup_{j_1, j_2 \geq 0} T \cdot r_{j_1, j_2, \epsilon} \cdot K_0.$$

Let $T = T_{\text{III}, D, AD}$. Recall that we put

$$C_0 := \{[g_1, g_2] : g_1, g_2 \in \text{GL}(2, F); \det g_1 = \det g_2\} \subset \text{GSp}(2, F).$$

It is isomorphic to C'_0 via $[g_1, g_2] \rightarrow (g_1, g_2)$. By abuse of notation, we let T and δ denote also their images in C'_0 .

For $m \in \mathbb{Z}$, recall that

$$K_m := \{(g, h) \in C'_0 : g - ehe \in \varpi^m M_2(R)\}, \quad e = \text{diag}(1, -1).$$

Let $R_T = T \cap K$. By the same technique used in the case of $T_{\text{I}, A}$, we have:

$$(B.15) \quad O_\delta(1_K) = \sum_{m \geq 0} (-1)^m \sum_{j_1, j_2 \geq 0} \sum_{\epsilon \in \{1, \epsilon_0\}} \varepsilon(r) \cdot [R_T : T \cap rK_0 r^{-1}] \int_{K_m \setminus K_0} 1_{K_m}(k^{-1} r^{-1} \delta r k) dk.$$

Here, r is the abbreviation of $r_{j_1, j_2, \epsilon}$.

Recall that $\xi = a_1 + b_1 \sqrt{D}$, $\zeta = a_2 + b_2 \sqrt{AD}$, where $b_i = B_i \varpi^{N_i}$, $B_i \in R^\times$, $N_i \in \mathbb{Z}$. For nonnegative integers j_1 and j_2 , let $\nu_1 = N_1 - j_1$ and $\nu_2 = N_2 - j_2$. Let $D_1 = D$ and $D_2 = AD$. For $r = r_{j_1, j_2, \epsilon}$, we have:

$$r^{-1} \delta r = \left(\begin{pmatrix} a_1 & b'_1 D'_1 \\ b'_1 & a_1 \end{pmatrix}, \begin{pmatrix} a_2 & b'_2 D'_2 \\ b'_2 & a_2 \end{pmatrix} \right),$$

where

$$\begin{aligned} b'_1 &= (B_1 \varpi^{\nu_1}) / (\epsilon \epsilon_0^{j_1}), & D'_1 &= D_1 \epsilon^2 \cdot (\epsilon_0 \varpi)^{2j_1}. \\ b'_2 &= B_2 \varpi^{\nu_2}, & D'_2 &= D_2 \varpi^{2j_2}. \end{aligned}$$

Put $R_m := R / (\varpi^m R)$. For $a \in R$, let $\bar{a}$ denote its image in R_m . Following the same argument in [F3, p. 64], $\int_{K_m \setminus K_0} 1_{K_m}(k^{-1} r^{-1} \delta r k) dk$ is equal to the cardinality of

$$L_{m, r} := \left\{ x = \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix} \in SL(2, R_m) : \begin{pmatrix} \bar{a}_1 & \bar{b}'_1 D'_1 \\ \bar{b}'_1 & \bar{a}_1 \end{pmatrix} \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix} = \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix} \begin{pmatrix} \bar{a}_2 & \bar{b}'_2 D'_2 \\ \bar{b}'_2 & \bar{a}_2 \end{pmatrix} \right\}.$$

Recall that $X = \text{ord}(a_1 - a_2)$. Transcribing Lemma II. G. 3 of [F3] to the current situation, the following holds:

B.19. CLAIM. *The set $L_{m, r}$ ($r = r_{j_1, j_2, \epsilon}$) is nonempty precisely when the following conditions are satisfied:*

- (1) $0 \leq m \leq X$.
- (2) $\nu_1, \nu_2 \geq 0$.
- (3) $m \leq \nu_1$ if and only if $m \leq \nu_2$.
- (4) If $\nu_1 < m$ or $\nu_2 < m$, then $\nu_1 = \nu_2$; we denote the common value by ν .
- (5) If $\nu < m$, then $(B_1 / \epsilon \epsilon_0^{j_1}) / B_2 \in R^{\times 2}$.
- (6) If $\nu < m$, then $m \leq 2N_i - \nu + \text{ord } D_i$ ($i = 1, 2$).

REMARK. The only differences between the above conditions and those in Lemma II. G. 3 of [F3] are item 5 and the definition of D'_i ($i = 1, 2$).

Recall that q_0 is the cardinality of the residue field of F .

B.20. CLAIM. *If $L_{m,r}$ is nonempty, then*

$$\#L_{m,r} = \begin{cases} 1 & \text{if } m = 0, \\ (q_0^2 - 1)q_0^{3m-1} & \text{if } 1 \leq m \leq \nu_1 \text{ (equivalently: } 1 \leq m \leq \nu_2), \\ 2q_0^{m+2\nu} & \text{if } \nu < m. \end{cases}$$

PROOF. This follows from Lemma II. G. 3 of [F3]. $\square$

B.21. CLAIM. *For $r = r_{j_1, j_2, \epsilon}$, the following holds:*

$$[R_T : T \cap rK_0r^{-1}] = q_0^{j_1+j_2}.$$

PROOF. For $k = F_D$ or F_{AD} , and a nonnegative integer j , put $R_k(j) := R + \varpi^j R_k$. The group $R_T = T \cap K$ is isomorphic to $(R_{F_D}^\times \times R_{F_{AD}}^\times)'$, where the prime indicates equal determinants. The computation of $[R_T : T \cap rK_0r^{-1}]$ is equivalent to computing the cardinality of the kernel in the short exact sequence

$$\begin{aligned} 1 &\rightarrow (R_{F_D}^\times \times R_{F_{AD}}^\times)' / (R_{F_D}(j_1)^\times \times R_{F_{AD}}(j_2)^\times)' \\ &\rightarrow (R_{F_D}^\times \times R_{F_{AD}}^\times) / (R_{F_D}(j_1)^\times \times R_{F_{AD}}(j_2)^\times) \\ &\rightarrow (R_{F_D}^\times \times R_{F_{AD}}^\times) / \left[(R_{F_D}^\times \times R_{F_{AD}}^\times)' (R_{F_D}(j_1)^\times \times R_{F_{AD}}(j_2)^\times) \right] \rightarrow 1. \end{aligned}$$

As shown in [F3, Lemma I. I. 2], for $k = F_D, F_{AD}$,

$$[R_k^\times : R_k(j)^\times] = [R_k^\times : 1 + \varpi^j R_k] / [R_k^\times : 1 + \varpi^j R] = q_0^j.$$

Hence, the cardinality of the middle term of the sequence is $q_0^{j_1+j_2}$. The image in the short exact sequence is isomorphic via $N_{F_D/F} \times N_{F_{AD}/F}$ to

$$\begin{aligned} \text{(B.16)} \quad & (N_{F_D/F} R_{F_D}^\times \times N_{F_{AD}/F} R_{F_{AD}}^\times) / \\ & \{ x \cdot N_{F_D/F} R_{F_D}(j_1)^\times \times x \cdot N_{F_{AD}/F} R_{F_{AD}}(j_2)^\times : \\ & \quad x \in N_{F_D/F} R_{F_D}^\times \cap N_{F_{AD}/F} R_{F_{AD}}^\times \}. \end{aligned}$$

Since the field extensions F_D/F and F_{AD}/F are ramified, we have

$$N_{F_D/F} R_{F_D}^\times = N_{F_{AD}/F} R_{F_{AD}}^\times = R^{\times 2}.$$

Consequently, the set (B.16) has cardinality 1. The claim follows. $\square$

Rearranging the terms in (B.15), and noting that $\varepsilon(r_{j_1, j_2, \epsilon})$ is equal to

$$\varepsilon \left(\det \left(\phi^D(\varpi_D^{-j_1}) \phi^D(\varpi_D^{-s(\epsilon)}) \right) \cdot (\epsilon_0 \varpi)^{j_1} \right) = \varepsilon \left((\epsilon_0 \varpi)^{s(\epsilon)} \right) = (-1)^{s(\epsilon)},$$

the expression (B.15) is the sum of the following two quantities:

$$\text{(B.17)} \quad \sum_{\epsilon=1, \epsilon_0} (-1)^{s(\epsilon)} \sum_{\substack{0 \leq \nu_1 \leq N_1 \\ 0 \leq \nu_2 \leq N_2}} q_0^{N_1 - \nu_1 + N_2 - \nu_2} \left(1 + (1 - q_0^{-2}) \sum_{m=1}^{\min(\nu_1, \nu_2)} q_0^{3m} \right),$$

$$\text{(B.18)} \quad \sum_{\epsilon=1, \epsilon_0} (-1)^{s(\epsilon)} q_0^{N_1 + N_2 - 2\nu} \sum_{0 \leq \nu \leq N} \sum_{\nu < m \leq X - \nu} (-1)^m L_{m, r_{j_1, j_2, \epsilon}}.$$

Here, ν is the common value of ν_1 and ν_2 when $\nu_1 = \nu_2 < m$.

The second sum in (B.17) is independent of ϵ ; hence, (B.17) is equal to zero.

In (B.18), suppose $L_{m,r} \neq 0$. Since $\nu < m$, by Claim B.19 we have $B_1/(B_2 \epsilon \epsilon_0^{j_1}) \in R^{\times 2}$. If B_1/B_2 lies in $R^{\times 2}$, then since $\epsilon_0 \in R^\times - R^{\times 2}$, j_1 is even or odd depending on

(in that order) ϵ being 1 or ϵ_0 . If $B_1/B_2 \notin R^{\times 2}$, then j_1 is odd or even depending on (in that order) ϵ being 1 or ϵ_0 . Recall that κ is the character of $R^\times$ which is 1 at squares and -1 otherwise. Thus, $(-1)^{s(\epsilon)}$ is equal to $\kappa(B_1 B_2)(-1)^{j_1}$ if $L_{m,r_{j_1,j_2},\epsilon} \neq 0$.

By the comments above and Claim B.20, we conclude that

$$\begin{aligned} O_\delta(1_K) &= \kappa(B_1 B_2) \sum_{0 \leq \nu \leq N} (-1)^{N_1 - \nu} q_0^{N_1 + N_2 - 2\nu} \sum_{\nu < m \leq X - \nu} (-1)^m 2q_0^{m+2\nu} \\ &= 2\kappa(B_1 B_2)(-1)^{N_1} q_0^{N_1 + N_2} \sum_{0 \leq \nu \leq N} (-1)^\nu \sum_{\nu < m \leq X - \nu} (-q_0)^m \\ &= 2\kappa(B_1 B_2)(-1)^{N_1} q_0^{1+N_1+N_2} \frac{1}{q_0 + 1} \sum_{0 \leq \nu \leq N} [(-q_0)^X q_0^{-\nu} - q_0^\nu] \\ &= 2\kappa(B_1 B_2)(-1)^{N_1} q_0^{1+N_1+N_2} \frac{1}{q_0^2 - 1} ((-1)^X q_0^{X-N} - 1)(q_0^{N+1} - 1). \end{aligned}$$

By Lemma II. L. 1 of [F3], $X = 2N + 1$. Hence,

$$\begin{aligned} O_\delta(1_K) &= 2\kappa(B_1 B_2)(-1)^{N_1} q_0^{1+N_1+N_2} \left(\frac{(q_0^2)^{N+1} - 1}{q_0^2 - 1} \right) \\ &= 2\kappa(B_1 B_2)(-1)^{N_1} q_0^{1+N_1+N_2} \left(\frac{q^{N+1} - 1}{q - 1} \right). \end{aligned}$$

□

B.2.3.3. Transfer Factor— $T_{2,D}, T_{\text{III},D,AD}$. We define the transfer factor $\Delta(\delta_2, \delta)$ to be

$$\frac{1}{2} \kappa \left(\frac{\xi - \tau\xi}{2\sqrt{D}} \cdot \frac{\zeta - \sigma\tau\zeta}{2\sqrt{AD}} \right) \varepsilon \left(\frac{\xi - \tau\xi}{2\sqrt{A}} \right) \left(\frac{D_G(\delta)}{D_{H_2}(\delta_2)} \right)^{1/2}.$$

B.22. CLAIM. *The following holds:*

$$\Delta(\delta_2, \delta) = \frac{1}{2} \kappa(B_1 B_2)(-1)^{N_1} q_0^{-1-N_1-N_2}.$$

PROOF. Note that by hypothesis $\xi = N_{E/F_D} l = l\sigma l$ and $\zeta = N_{E/F_{AD}} l = l\sigma\tau l$, where $l \in R_E^\times$. The claim then follows from a straightforward computation. □

B.23. PROPOSITION. *The following identity holds:*

$$O_{\delta_2}(1_{K_2}) = \Delta(\delta_2, \delta) O_\delta(1_K).$$

PROOF. This follows from Claims B.17, B.18, and B.22. □

B.2.4. Comparison for $T_{2,D}, T_{\text{IV},A,D}$. Let D be an element in $F_A^\times - F_A^{\times 2} \cup F^\times$. Let $E = F_A(\sqrt{D}) = F(\sqrt{D})$. Let $\tilde{E}$ be the Galois closure of E over F . Then, $\text{Gal}(\tilde{E}/F) = \text{Gal}(E/F)$ is cyclic of order 4 if E/F is Galois, and $\text{Gal}(\tilde{E}/F) = D_4$ if E/F is non-Galois.

Let σ be the element in $\text{Gal}(\tilde{E}/F)$ defined by $\sigma\sqrt{A} = -\sqrt{A}$, $\sigma\sqrt{D} = \sqrt{\sigma D}$, $\sigma^2\sqrt{D} = -\sqrt{D}$, $\sigma^3\sqrt{D} = -\sqrt{\sigma D}$. Note that σ is of order 4, and it generates $\text{Gal}(E/F)$ if E/F is Galois. Moreover, $\text{Gal}(F_A/F)$ is generated by the restriction of σ to F_A .

Let q be the cardinality of the residue field of F_A . Let q_0 be the cardinality of the residue field of F . Since by assumption F_A/F is unramified, $q = q_0^2$. We fix a uniformizer ϖ for both F and F_A .

Let l be an element in $R_E^\times - R_{F_A}^\times$, and c an element in $R^\times$. Suppose $l = \alpha + \beta\sqrt{D}$, where $\alpha, \beta \in F_A$. Let

$$\delta_2 = (\phi^{F_A:D}(l), c)_* := \left(\begin{pmatrix} \alpha & \beta D \\ \beta & \alpha \end{pmatrix}, c \right)_* \in T_{2,D} \cap K_2.$$

Here, lower $*$ indicates image in H_2 .

For an element $\gamma = a + b/\sqrt{A} \in F_A^\times$, with $a, b \in F$, put $\gamma := \begin{pmatrix} a & b/A \\ b & a \end{pmatrix}$. Suppose $l\sigma l = \xi_1 + \xi_2\sqrt{D}$, with $\xi_1, \xi_2 \in F_A$. Let

$$\delta = c \begin{pmatrix} \xi_1 & \xi_2 D \\ \xi_2 & \xi_1 \end{pmatrix} \in T_{IV,A,D} \cap K.$$

The element δ is regular, and δ_2 is a norm of δ . The conjugacy classes of δ_2 and δ are stable. We assume that δ is topologically unipotent.

Let $n = \text{ord}_{F_A} \beta$, and $N = \text{ord}_{F_A} \xi_2$. Let B be the element in $R_{F_A}^\times$ such that $\xi_2 = B\varpi^N$.

B.2.4.1. *Stable Orbital Integral— $T_{2,D}$.*

B.24. CLAIM. *The stable orbital integral $O_{\delta_2}(1_{K_2})$ is equal to*

- $\frac{(q+1)q^n - 2}{q-1}$ if E/F_A is unramified,
- $\frac{q^{n+1} - 1}{q-1}$ if E/F_A is ramified.

PROOF. This follows from Lemma I. 1. 2 of [F3]. □

B.2.4.2. *Twisted Orbital Integral— $T_{IV,A,D}$.* Let κ be the nontrivial quadratic character of the group $R_{F_A}^\times / R_{F_A}^{\times 2}$.

B.25. CLAIM. *The ε -twisted orbital integral $O_\delta(1_K)$ is equal to*

- $(-q)^N \left(\frac{(q+1)q^N - 2}{q-1} \right)$ if E/F_A is unramified,
- $-2\kappa(B)(-q)^N q_0 \left(\frac{q^{N+1} - 1}{q-1} \right)$ if E/F_A is ramified.

PROOF. Let

$$C_A = \text{GL}(2, F_A)' := \{g \in \text{GL}(2, F_A) : \det g \in F^\times\},$$

$$\mathbf{C}_A = \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \eta \end{pmatrix} : \begin{pmatrix} \alpha & \beta \\ \gamma & \eta \end{pmatrix} \in C_A \right\}.$$

For a nonnegative integer m , let $u_m = \begin{pmatrix} 1 & 0 & 0 & \varpi^{-m}/A \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$. Let

$$K_m = \text{GL}(2, R_{F_A}(m))',$$

$$K_m^A = \mathbf{C}_A \cap u_m K u_m^{-1}.$$

Here, $R_{F_A}(m) := R + (\varpi^m \sqrt{A})R$, and the prime indicates determinant in $F^\times$.

For $a = a_1 + a_2/\sqrt{A}$, $b = b_1 + b_2/\sqrt{A}$, $c = \dots \in F_A^\times$, put

$$\phi \begin{pmatrix} a & b \\ c & d \end{pmatrix} := \begin{pmatrix} a_1 & a_2/A & b_2 & b_1 \\ a_2 & a_1 & b_1 A & b_2 \\ c_2/A & c_1/A & d_1 & d_2/A \\ c_1 & c_2/A & d_2 & d_1 \end{pmatrix}.$$

We define an isomorphism $\phi_m : C_A \rightarrow \mathbf{C}_A$ as follows:

$$\phi_m : \begin{pmatrix} \alpha & \beta \\ \gamma & \eta \end{pmatrix} \mapsto \phi \left(\begin{pmatrix} 1 & 0 \\ 0 & A\varpi^m \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{A} \end{pmatrix} \begin{pmatrix} \alpha & \beta \\ \gamma & \eta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/\sqrt{A} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/(A\varpi^m) \end{pmatrix} \right).$$

In particular, ϕ_m maps K_m isomorphically onto K_m^A ([F3, Lemma I. J. 3]).

We have the disjoint decomposition ([F3, Lemma I. J. 1]):

$$(B.19) \quad G = \dot{\bigcup}_{m \geq 0} \mathbf{C}_A u_m K.$$

Let $T = T_{IV,A,D}$. Noting that $\varepsilon(u_m) = 1$, we have:

$$(B.20) \quad \begin{aligned} O_\delta(1_K) &= \int_{T \setminus G} 1_K(g^{-1}\delta g)\varepsilon(g) dg \\ &= \sum_{m \geq 0} |K|_G \int_{T \setminus \mathbf{C}_A / \mathbf{C}_A \cap u_m K u_m^{-1}} 1_K(u_m^{-1}h^{-1}\delta h u_m)\varepsilon(h u_m) dh \\ &= \sum_{m \geq 0} |K|_G \int_{T \setminus \mathbf{C}_A / K_m^A} 1_{K_m^A}(h^{-1}\delta h)\varepsilon(h) dh. \end{aligned}$$

Let δ' be the element in C_A such that $\phi(\delta') = \delta$. Let $T' = Z_{C_A}(\delta')$. For $m \geq 0$, let $\delta_m = \phi_m^{-1}(\delta) \in C_A$. Let $T_m = Z_{C_A}(\delta_m) = \phi_m^{-1}(T)$. We rewrite (B.20) as follows:

$$(B.21) \quad O_\delta(1_K) = \sum_{m \geq 0} |K|_G \int_{T_m \setminus C_A / K_m} 1_{\phi_m(K_m)}(\phi_m(h')^{-1}\delta\phi_m(h'))\varepsilon(\phi_m(h'))dh'.$$

Here, h' denotes the element in C_A such that $\phi_m(h') = h \in \mathbf{C}_A$.

The similitude factor of $\phi(h')$ is $(\det h')^{-1}$; hence,

$$\varepsilon(\phi_m(h')) = \varepsilon((\det h')^{-1}) = \varepsilon(\det h').$$

We change variables $h' \mapsto \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/(A\varpi^m) \end{pmatrix} h'$. The expression (B.21) becomes

$$\sum_{m \geq 0} |K|_G |K_m|_{C_A}^{-1} \int_{T' \setminus C_A} 1_{K_m}(h'^{-1}\delta'h')\varepsilon(A\varpi^m)\varepsilon(\det h') dh'.$$

Since F_A/F is unramified, $\varepsilon(A)$ is equal to one. Consequently, $\varepsilon(A\varpi^m) = (-1)^m$ for each nonnegative integer m .

Let $\tilde{T}'$ be the centralizer of δ' in $\mathrm{GL}(2, F_A)$. It contains T' . Let

$$\begin{aligned} K' &= \mathrm{GL}(2, R_{F_A})' = \{k \in \mathrm{GL}(2, R_{F_A}) : \det k \in F^\times\}, \\ T'_0 &= T' \cap K'. \end{aligned}$$

Suppose E/F_A is unramified. Since

$$\mathrm{GL}(2, F_A) = \dot{\bigcup}_{j \geq 0} \tilde{T}' \cdot \begin{pmatrix} 1 & \\ & \varpi^j \end{pmatrix} \cdot \mathrm{GL}(2, R_{F_A}),$$

the group C_A decomposes as follows ([F3, Lemma II. M. 3]):

$$C_A = \dot{\bigcup}_{j, \epsilon} T' \cdot r_{j, \epsilon} \cdot K'.$$

The disjoint union is over nonnegative integers j and $\epsilon \in R_{F_A}^\times / R_{F_A}^{\times 2}$ if $j \geq 1$. Here, $r_{j, \epsilon} = t_\epsilon \begin{pmatrix} 1 & 0 \\ 0 & \epsilon\varpi^j \end{pmatrix}$, where t_ϵ is an element in $\tilde{T}'$ such that $\det t_\epsilon = \epsilon^{-1}$.

By the above decomposition of C_A , the twisted orbital integral $O_\delta(1_K)$ is equal to

$$(B.22) \quad \sum_{m \geq 0} (-1)^m \sum_{r=r_{j,\epsilon}} \varepsilon(\det r) [T'_0 : T' \cap rK'r^{-1}] \cdot [K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk.$$

Since F_A/F is unramified, $\varepsilon(r_{j,\epsilon})$ is equal to $(-1)^j$. Thus, $O_\delta(1_K)$ is equal to

$$(B.23) \quad \sum_{m \geq 0} (-1)^m \sum_{r=r_{j,\epsilon}} (-1)^j [T'_0 : T' \cap rK'r^{-1}] \cdot [K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk.$$

Recall: $\delta = c \begin{pmatrix} \xi_1 & \xi_2 D \\ \xi_2 & \xi_1 \end{pmatrix} \in T$, $N = \text{ord}_{F_A} \xi_2$. Let $X = \text{ord}_{F_A}(\xi_1 - \sigma\xi_1)$.

By the computation of $[T'_0 : T' \cap rK'r^{-1}]$, $[K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk$ in [F3, Sect. II. M], and (B.23), the twisted orbital integral $O_\delta(1_K)$ is equal to the sum of the following two quantities if E/F_A is unramified:

$$\begin{aligned} \text{I.} & \left(\sum_{m=0} 1 + \frac{q+1}{q} \sum_{1 \leq m \leq N} (-q_0)^{3m} \right) \left(\sum_{\nu=m} 1 + \frac{q+1}{q} \sum_{m \leq \nu < N} (-q)^{N-\nu} \right), \\ \text{II.} & \sum_{0 \leq \nu < N} \frac{q+1}{q} (-q)^{N-\nu} \sum_{\nu < m \leq X-\nu} (-q_0)^m q^\nu. \end{aligned}$$

The sum I is equal to

$$(B.24) \quad \frac{(-q)^N}{1-q_0} \left(1 - \left(\frac{q+1}{q_0} \right) q_0^N + q_0^{-1} \right),$$

and the sum II is equal to

$$(B.25) \quad \frac{(q+1)(-q)^N}{(q_0+1)(1-q_0)q_0} (q_0^{X-N+1} - q_0^{X+1} + q_0^N - 1).$$

From [F3, Sect. II. M], since δ is topologically unipotent, X is equal to $2N$. Hence, $O_\delta(1_K) = \text{I} + \text{II}$ is equal to

$$\begin{aligned} &= \frac{(-q)^N}{1-q_0} \left(1 - \left(\frac{q+1}{q_0} \right) q_0^N + q_0^{-1} + \frac{q+1}{(q_0+1)q_0} q_0^{X-N+1} \right) \\ &= \frac{(-q)^N}{1-q_0} \frac{1}{(q_0+1)q_0} ((q+1)(-q_0^{2N+1} - 1) + q_0^2 + 2q_0 + 1) \\ &= \frac{(-q)^N}{-(q-1)} \frac{1}{q_0} ((q+1)(-q^N q_0) - q - 1 + q + 2q_0 + 1) \\ &= (-q)^N \left[\frac{(q+1)q^N - 2}{q-1} \right]. \end{aligned}$$

This completes the proof of the claim for the case where E/F_A is unramified.

Now, suppose E/F_A is ramified. Recall that we have fixed a uniformizer ϖ for both F and F_A . Since E/F_A is ramified, and $D \in F_A^\times - F^\times$, there exists an element $\epsilon_0 \in R_{F_A}^\times - R_{F_A}^{\times 2} \cup F^\times$ such that $D = -\epsilon_0 \varpi$. Then, $N_{E/F_A} \sqrt{D} = \epsilon_0 \varpi$, and the element $t_0 = \phi^{F_A:D}(\sqrt{D}) \in \tilde{T}'$ has determinant $\epsilon_0 \varpi$.

The group C_A has a disjoint decomposition as follows:

$$C_A = \bigcup_{j,\epsilon} T' \cdot r_{j,\epsilon} \cdot K',$$

where j ranges over the nonnegative integers, and ϵ over the set $\{1, \epsilon_0\}$. Here, $r_{j,\epsilon}$ is an element in $\tilde{T}' \left(\begin{smallmatrix} 1 \\ (\epsilon_0 \varpi)^j / \epsilon \end{smallmatrix} \right)$ such that $\det r_{j,\epsilon} = \begin{cases} 1 & \text{if } \epsilon = 1, \\ \varpi & \text{if } \epsilon = \epsilon_0. \end{cases}$

The twisted orbital integral $O_\delta(1_K)$ may then be expressed as the sum

$$(B.26) \quad \sum_{m \geq 0} (-1)^m \sum_{r=r_{j,\epsilon}} (-1)^{s(\epsilon)} [T'_0 : T' \cap rK'r^{-1}] \cdot [K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk,$$

where $s(\epsilon)$ is equal to 0 if $\epsilon = 1$, 1 if $\epsilon = \epsilon_0$.

Recall that $\delta = c \begin{pmatrix} \xi_1 & \xi_2 D \\ \xi_2 & \xi_1 \end{pmatrix} \in \mathbf{C}_A$. Its preimage under ϕ in C_A is the element $\delta' = c \phi^{F_A:D}(\xi_1 + \xi_2 \sqrt{D})$. Recall that B is the element in $R_{F_A}^\times$ such that $\xi_2 = B\varpi^N$ ($N = \text{ord}_{F_A} \xi_2$).

For a nonnegative integer j , let $\nu = N - j$. Then,

$$r_{j,\epsilon}^{-1} \delta' r_{j,\epsilon} = \begin{pmatrix} \xi_1 & \xi_2' D' \\ \xi_2' & \xi_1 \end{pmatrix},$$

where $\xi_2' = (B\epsilon/\epsilon_0^j)\varpi^\nu$, $D' = (D/\epsilon^2)(\epsilon_0\varpi)^{2j}$.

Let $X = \text{ord}_{F_A}(\xi_1 - \sigma\xi_1)$. By the same argument used in the proof of Lemma II. M. 6 in [F3], the integral $\int_{K_0} 1_{K_m}(k^{-1}r_{j,\epsilon}^{-1}\delta'rk) dk$ is nonzero precisely when $0 \leq \nu \leq N$ and $0 \leq m \leq X$. Moreover, if $m > \nu$, then $\nu + m \leq X$ and ϵ_0^j lies in $B\epsilon R_{F_A}^{\times 2}$.

For $\nu < m \leq X - \nu$, the condition $\epsilon_0^j \in B\epsilon R_{F_A}^{\times 2}$ implies that j is even if $B\epsilon \in R_{F_A}^{\times 2}$, and odd otherwise. Thus, $\epsilon \in \{1, \epsilon_0\}$ depends on j . Moreover, $(-1)^{s(\epsilon)}$ is equal to $\kappa(B)(-1)^j$, where κ is the nontrivial quadratic character of the group $R_{F_A}^\times / R_{F_A}^{\times 2}$.

The quantities $[T'_0 : T' \cap rK'r^{-1}]$ and $[K_0 : K_m] \int_{K_0} 1_{K_m}(k^{-1}r^{-1}\delta'rk) dk$ are computed in [F3, Sect. II. M]. Recall that q_0 is the cardinality of the residue field of F , and $q = q_0^2$ that of F_A . The twisted orbital integral $O_\delta(1_K)$ is equal to the sum of the following:

$$\begin{aligned} (1) & \sum_{\epsilon=1, \epsilon_0} (-1)^{s(\epsilon)} \sum_{0 \leq \nu \leq N} q^{N-\nu}, \\ (2) & \sum_{\epsilon=1, \epsilon_0} (-1)^{s(\epsilon)} \sum_{0 \leq \nu \leq N} \frac{q^{N-\nu}(q+1)}{q} \sum_{1 \leq m \leq \nu} (-q_0)^{3m}, \\ (3) & \sum_{0 \leq \nu \leq N} \kappa(B)(-q)^{N-\nu} \sum_{\nu < m \leq X-\nu} 2(-q_0)^m q^\nu \\ & = 2\kappa(B)(-q)^N \sum_{0 \leq \nu \leq N} (-1)^\nu \sum_{\nu < m \leq X-\nu} (-q_0)^m. \end{aligned}$$

The first two sums are zero, for their inner sums are independent of ϵ . The twisted orbital integral $O_\delta(1_K)$ is therefore equal to the third sum, which is

$$2\kappa(B)(-q)^N (-q_0) \left(\frac{(-1)^X q_0^{X-N} - (-1)^X q_0^{X+1} + q_0^{N+1} - 1}{q - 1} \right).$$

From [F3, Sect. II. M], $X = 2N + 1$; hence,

$$O_\delta(1_K) = -2\kappa(B)(-q)^N q_0 \left(\frac{q_0^{2N+2} - 1}{q - 1} \right) = -2\kappa(B)(-q)^N q_0 \left(\frac{q^{N+1} - 1}{q - 1} \right).$$

□

B.2.4.3. Transfer Factor— $T_{2,D}$, $T_{IV,A,D}$. Recall: $l = \alpha + \beta\sqrt{D} \in R_E^\times$, $c \in R^\times$, $l\sigma l = \xi_1 + \xi_2\sqrt{D}$. We let δ_2 be the element $(\phi^{F_A:D}(l), c)_*$ in $T_{2,D}$. It is a norm of $\delta = c \begin{pmatrix} \xi_1 & \xi_2 D \\ \xi_2 & \xi_1 \end{pmatrix} \in T_{IV,A,D}$. Recall that $n = \text{ord}_{F_A}\beta$, $N = \text{ord}_{F_A}\xi_2$, and $\xi_2 = B\varpi^N$, with $B \in R_{F_A}^\times$.

Suppose E/F_A is unramified. We extend ε to $E^\times$ by letting $\varepsilon(x) = (-1)^{\text{ord}_E x}$ for all $x \in E^\times$. We define the transfer factor for (δ_2, δ) as follows:

$$\Delta(\delta_2, \delta) = \varepsilon \left(\frac{\xi - \sigma^2 \xi}{2\sqrt{D}} \right) \left(\frac{D_G(\delta)}{D_{H_2}(\delta_2)} \right)^{1/2}.$$

B.26. PROPOSITION. *The following identity holds:*

$$O_{\delta_2}(1_{K_2}) = \Delta(\delta_2, \delta) O_\delta(1_K).$$

PROOF. By Lemma M. II. 1 of [F3], we have $n = N$. The proposition then follows from Claims B.24 and B.25. $\square$

Suppose E/F_A is ramified. Let ε_{E/F_A} be the quadratic character associated with the extension E/F_A . We define the transfer factor for (δ_2, δ) as follows:

$$\Delta(\delta_2, \delta) = -\frac{1}{2} \varepsilon_{E/F_A} \left(\frac{l\sigma l - \sigma^2 l \sigma^3 l}{2\sqrt{D}} \right) \left(\frac{D_G(\delta)}{D_{H_2}(\delta_2)} \right)^{1/2}.$$

B.27. PROPOSITION. *The following identity holds:*

$$O_{\delta_2}(1_{K_2}) = \Delta(\delta_2, \delta) O_\delta(1_K).$$

PROOF. Note that $(l\sigma l - \sigma^2 l \sigma^3 l)/(2\sqrt{D})$ is equal to $\xi_2 = B\varpi^N$. Recall that $D = -\epsilon_0\varpi$, where $\epsilon_0 \in R_{F_A}^\times - R_{F_A}^{\times 2}$. The element $\xi_2 \in F_A^\times$ may be written as $\xi_2 = B\epsilon_0^{-N}(\epsilon_0\varpi)^N$. The element $\epsilon_0\varpi = N_{E/F_A}\sqrt{D}$ is a norm from $E^\times$ but ϵ_0 is not. Moreover, since $R_{F_A}^\times \cap N_{E/F_A}E^\times = R_{F_A}^{\times 2}$, the restriction of ε_{E/F_A} to $R_{F_A}^\times$ coincides with κ . Hence, $\varepsilon_{E/F_A}(\xi_2) = (-1)^N \kappa(B)$.

The Jacobian factor $(D_G(\delta)/D_{H_2}(\delta_2))^{1/2}$ is equal to

$$|\xi_2 \sqrt{D}|_{\bar{F}} = |\xi_2|_{F_A} |D|_{F_A}^{1/2} = q^{-N} q^{-1/2} = q^{-N} q_0^{-1}.$$

By Lemma II. M. 1 of [F3], $n = N$. The proposition then follows from Claims B.24 and B.25. $\square$

Bibliography

- [A] Arthur, J.; *A trace formula for reductive groups I*. Duke Math. J. 45 (1978), 911–952.
- [A1] —; *A trace formula for reductive groups II*. Compositio Math. 40 (1980), 87–121.
- [A2] —; *The local behaviour of weighted orbital integrals*. Duke Math. J. 56 (1988), 223–293.
- [A3] —; *Automorphic representations of $\mathrm{GSp}(4)$* . Contributions to automorphic forms, geometry, and number theory, Johns Hopkins Univ. Press, Baltimore, MD (2004), 65–81.
- [AC] Arthur, J., Clozel, L.; *Simple algebras, base change, and the advanced theory of the trace formula*. Annals of Math. Studies No. 120, Princeton University Press (1989).
- [Be] Bernstein, J.; *Representations of p -adic groups*. Lectures given at Harvard University, Fall 1992, Notes by K. E. Rumelhart. <http://www.math.tau.ac.il/~bernstei/texts/harv93.pdf>.
- [BZ] Bernstein, I. N.; Zelevinskii, A. V.; *Representations of the group $\mathrm{GL}(n, F)$, where F is a non-archimedean local field*. Russian Math. Surveys 31 (1976), 1–68.
- [Bo] Borel, A.; *Automorphic L -functions*. Proc. Symp. Pure Math. 33 II, Amer. Math. Soc., Providence, RI (1979), 27–61.
- [BJ] Borel, A.; Jacquet, H.; *Automorphic forms and automorphic representations*. Proc. Symp. Pure Math. 33 I, Amer. Math. Soc., Providence, RI (1979), 189–208.
- [CLL] Clozel, L.; Labesse, J.-P.; Langlands, R.; *Morning seminar on the trace formula*. IAS notes (1984). <http://www.math.ubc.ca/~cass/Langlands/friday/friday.html>.
- [F] Flicker, Y.; *Rigidity for automorphic forms*. J. Analyse Math. 49 (1987), 135–202.
- [F1] —; *Regular trace formula and base change lifting*. Amer. J. of Math. 110 (1988) 739–764.
- [F2] —; *On the symmetric square lifting, Part I*. in *Automorphic forms on low rank groups*. World Scientific (2006).
- [F3] —; *Matching of orbital integrals on $\mathrm{GL}(4)$ and $\mathrm{GSp}(2)$* . Mem. Amer. Math. Soc. No. 655 (1999).
- [F4] —; *Lifting automorphic forms of $\mathrm{PGSp}(2)$, Part I*. in *Automorphic forms and Shimura varieties of $\mathrm{PGSp}(2)$* . World Scientific (2005).
- [FK] Flicker, Y.; Kazhdan, D.; *Metaplectic correspondence*. Inst. Hautes Études Sci. Publ. Math. 64 (1987), 53–110.
- [FK1] —; *A simple trace formula*. J. d’Analyse Math. 50 (1988), 189–200.
- [Fo] Folland, G.; *Real analysis. Modern techniques and their applications*. Second edition. Pure and Applied Mathematics (New York). A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York (1999).
- [H] Hales, T.; *A simple definition of transfer factors for unramified groups*. Contemp. Math. 145, Amer. Math. Soc., Providence, RI (1993), 109–134.
- [H1] —; *Unipotent representations and unipotent classes in $\mathrm{SL}(N)$* . Amer. J. Math. 115 (1993), 1347–1383.
- [H2] —; *On the fundamental lemma for standard endoscopy: reduction to unit elements*. Can. J. Math. 47 (1995), 974 – 994.
- [HC] Harish-Chandra; *Harmonic analysis on reductive p -adic groups*. Notes by G. van Dijk. Lecture Notes in Mathematics 162. Springer-Verlag, Berlin-New York (1970).
- [HC1] —; *Admissible invariant distributions on reductive p -adic groups*. Queen’s Papers in Pure Appl. Math. 48 (1978), 281–347. See also:
Admissible invariant distributions on reductive p -adic groups. Preface and notes by Stephen DeBacker and Paul J. Sally, Jr. University Lecture Series, 16. Amer. Math. Soc., Providence, RI (1999).
- [JL] Jacquet, H.; Langlands, R.P.; *Automorphic forms on $\mathrm{GL}(2)$* . Lecture Notes in Mathematics 114. Springer-Verlag, Berlin-New York (1970).
- [JS] Jacquet, H.; Shalika, J. A.; *On Euler products and the classification of automorphic representations I*. Amer. J. of Math. 103 (1981), 499–558.
- [JS1] —; *On Euler products and the classification of automorphic forms II*. Amer. J. Math. 103 (1981), 777–815.
- [K] Kazhdan, D.; *On Lifting*. Lecture Notes in Math., 1041, Springer, Berlin (1984), 209–249.

- [KS] Kottwitz, R.; Shelstad, D.; *Foundations of twisted endoscopy*. Astérisque No. 255 (1999).
- [LL] Labesse, J.-P.; Langlands, R.; *L-indistinguishability for $SL(2)$* . Canad. J. Math. 31 (1979), 726–785.
- [La] Lang, S; *Algebraic number theory*. Springer-Verlag, New York (1994).
- [L] Langlands, R.; *Base change for $GL(2)$* . Annals of Math. Studies, 96. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo (1980).
- [L1] —; *Les débuts d'une formule des traces stable*. Publications Mathématiques de l'Université Paris VII, 13. Université de Paris VII, U.E.R. de Mathématiques, Paris, 1983.
- [LS] Langlands, R.; Shelstad, D.; *On the definition of transfer factor*. Math. Ann. 278 (1987), 219–271.
- [PR] Platonov, P.; Rapinchuk, A.; *Algebraic groups and number theory*. Academic Press (1994).
- [R] Rodier, F.; *Les représentations de $GSp(4, k)$ où k est un corps local*. C. R. Acad. Sci. Paris Sér. A-B 283 (1976), no. 7, Ai, A429–A431.
- [ST] Sally, P.; Tadic, M.; *Induced representations and classifications for $GSp(2, F)$ and $Sp(2, F)$* . Mém. Soc. Math. France (N.S.) No. 52 (1993), 75–133.
- [Sh] Shahidi, F.; *On nonvanishing of L -functions*. Bull. Amer. Math. Soc. (N.S.) 2 (1980), no. 3, 462–464.
- [Sp] Springer, T. A.; *Linear algebraic groups*. Birkhäuser, Boston (1998).
- [T] Tate, J.; *Number theoretic background*. Proc. Symp. Pure Math. 33 II (1979), 3–26.
- [Wa] Waldspurger, J.-L.; *Sur les intégrales orbitales tordues pour les groupes linéaires: un lemme fondamental*. Canad. J. Math. 43 (1991), no. 4, 852–896.
- [Wa1] —; *Le lemme fondamental implique le transfert*. Compositio Math. 105 (1997), 153–236.
- [We] Weissauer, R; *A special case of the fundamental lemma I*. Preprint. <http://www.mathi.uni-heidelberg.de/~weissaue/papers/ramanuge.dvi>.

List of Symbols

- 1_2 , 12, 100
 $1_{\mathrm{GL}(2, \mathbb{A}_F)}$, 57, 65
 $A(\pi)$, 12, 13
 A^L , 13
 A^δ , 13
 A_2 , 87
 $\mathcal{A}(\mathbf{H}, \omega)$, 3
 $\mathcal{A} \setminus$, 22
 $\mathbb{A}_F$, 3
 $\mathbf{A}_0$, 29, 31
 $\mathbf{A}_M$, 29
 $\mathbf{a}$, a , 16
 $\mathfrak{a}_M$, 29
 $\mathfrak{a}_M^*$, 29
 $a_{\mathbf{H}}$, 42
 $a_{\{\text{packet}\}}$, 48
 $B_{E/F}$, *see* Index: base change
 $[\ , \]$, 57
 $[\text{repn.}]$, 12, 13, 115, 118
 $\langle \ , \ \rangle$, 4, 8, 11, 66
 $\langle \ , \ \rangle_A$, 7, 10, 87
 $\langle \ , \ \rangle_{\varepsilon}$, 5, 66
 $[(\), (\)]$, 19
 $\{\Pi\}$, *see* Index: packet
 $\{\chi\}$, *see* Index: quadruple
 $\{\text{packet}\}_S$, *see* Index: packet, global, S -part
 b_ξ , 27
 $\bar{\ , \ }$, 18
 $C(H_v, \omega_v)$, 4
 $C(H_v, \omega_v)$, 29
 $C(\ , \)$, 85
 $C(\ , \ \omega)$, 7, 85
 $C(\mathbf{H}(\mathbb{A}_F), \omega)$, 4, 29
 C_0, C'_0 , 135
 $C_A, \mathbf{C}_A$, 148
 C_F , 3
 $\mathrm{CL}_{\mathrm{ss}}(\ , \)$, 22
 $\mathcal{C}(\ , \), \mathcal{C}(\ , \ , \)$, *see* Index: global datum
 C_0 , 99
 $[c_v]_{v \in \ , \ [c(\pi_v)]_{v \in \ , \ }$, *see* Index: global datum
 χ_ζ , 29
 $\chi_{\text{repn.}}$, *see* Index: Harish-Chandra character
 $\chi_{\text{repn.}}^A, \chi_{\tau_2}^\varepsilon$, *see* Index: Harish-Chandra character, twisted
 $D(\ , \)$, 36
 $D_H(\ , \)$, *see* discriminant
 $D_{M \setminus G}$, 90
 $D_{T \setminus H}$, 91
 $D_{T_K}(\ , \)$, 88
 $d(\ , \ , \)$, 17
 $\Delta(\ , \)$, *see* Index: transfer factor
 $\delta(\ , \)$, 29
 $\delta(\text{char.}, \text{repn.})$, 117
 $\delta(\text{repn.}, \text{char.})$, 12, 59
 $\delta^-(\text{repn.}, \text{char.})$, 12, 58, 59, 100
 $\delta_M(\ , \)$, 87
 DOR , *see* Index: representation, discretely occuring
 $E(u_1, u_2, \ , \)$, 41
 e , 17
 ε , 6, 27, 85
 $\varepsilon\text{-DOR}$, *see* Index: representation, discretely occuring
 $\varepsilon \otimes \pi$, *see* $\varepsilon\pi$
 $\varepsilon\pi$, 7
 ε_S , 40
 $f^\vee$, *see* Satake Transform
 f_S , 40
 $(\mathbf{G}, 1, \mathbf{a})$, 15
 $\mathrm{GL}(2, k)^K$, 87
 $\mathrm{GL}(2, \)^E$, 35
 $\mathrm{GSp}(2)$, 6, 10, 85
 $\bar{G}$, *see* $\mathrm{PGSp}(2)$
 $\mathbf{G}$, *see* $\mathrm{GSp}(2)$
 $\tilde{G}$, 82
 $\text{group}^{\text{reg}}$, 85
 $(\mathbf{H}, \mathcal{H}, s, \xi)$, 15
 H_1 , *see* $\mathbf{H}_1$
 H_2 , *see* $\mathbf{H}_2$
 H_S , 40
 $\text{Hom}_{\text{group}}(\text{repn.}, \text{repn.})$, 85
 $\hat{H}$, 15

- $\hat{H}_1$, 17–19
 $\hat{H}_2$, 19–21
 $\mathcal{H}$, 15
 $\mathcal{H}_1$, 17–19
 $\mathcal{H}_2$, 19–21
 $\mathcal{H}(H_v, \omega_v)$, 4
 $\mathcal{H}(H_v, \omega_v)$, 29
 $\mathbf{H}_1$, 7, 10, 17–19, 86
 $\mathbf{H}_2$, 7, 10, 19–21, 86
 $\mathcal{H}()$, 27
 $\mathcal{H}^\vee$, 47
 $\mathcal{H}_1$, 17
 $\mathcal{H}_2$, 19
 $H_P()$, 29
 h_w , 15

 I^+ , I^- , 37
 I^c , *see* Index: trace formula, continuous part
 I^d , *see* Index: trace formula, discrete part
 I_2 , 20
 $I_{P,\tau}()$, 29
 $I_{P,\tau}(\cdot, \cdot)$, 29
 $I_{P,\tau}(\cdot, \times \varepsilon)$, 30
 $I_{P,\tau}(\varepsilon)$, 30
 Int , 133
 $\text{Ind}_{W_{E/F}^{E/F}}$, 8
 ι , 23

 J , 6
 $J(\text{repn.}, \text{repn.})$, *see* Index: Langlands quotient
 $J_{\mathcal{O}}()$, 41
 $J_{\mathcal{O},\varepsilon}^T()$, 6, 30
 $J_{\chi,\varepsilon}^T()$, 6, 30

 K_m , 135
 K_m^A , 148
 K_v , 4
 $K_{1,v}$, 8
 $K_{f,\varepsilon}$, 5
 $K_{f,\varepsilon}^T(h)$, 6
 $\mathbb{K}$, 29

 $L(\cdot, \omega)$, 3
 $L_c(\cdot, \omega)$, 4
 $L_d(\cdot, \omega)$, 4
 $L(\text{char.}, \text{repn.})$, 12, 58, 100, 113, 117
 $L(\text{repn.}, \text{char.})$, 58, 59

 $L^S(\cdot, \text{repn.})$, *see* Index: L -function, partial
 ${}^L H$, *see also* Index: L -group
 ${}^L H_1$, 17
 ${}^L H_2$, 19
 λ , 22–24
 $\lambda(\cdot)$, *see* Index: similitude character, 85
 $\lambda(\cdot, \psi)$, 89
 λ_1 , 92, 96
 λ_2 , 94, 98

 $(\mathbf{M}, \mathbf{L}, \tau, s)$, *see* Index: quadruple
 $M()$, 48
 $M_P(\cdot, \cdot)$, 30
 M_β , 11
 $\mathbf{M}_\alpha, \mathbf{M}_\beta, \dots$, etc., 32
 $\mathcal{M}_L^T(\cdot, \cdot)$, *see* Index: logarithmic derivative
 $m(\cdot)$, *see* Index: multiplicity formula

 $\mathbf{N}$, 23
 $\mathbf{N}_1$, 23
 $\mathbf{N}_2$, 24
 ν , 11, 57

 $O(\cdot, \cdot)$, *see* Index: orbital integral
 $O^T(\cdot, \cdot)$, *see* Index: orbital integral, weighted
 $O^{\text{us}}()$, 139
 $O_G^\varepsilon(\cdot, \cdot)$, *see* Index: orbital integral, twisted
 $\mathcal{O}_v$, 3, 29
 $\{\mathcal{O}\}_e$, 41
 $\text{Out}(\cdot, \cdot, \cdot)$, 42
 ord , 11
 ω , 3, 7

 P_0 , 11
 P_1 , 92
 P_β , 11
 $\text{PGSp}(2)$, 99
 $\mathbf{P}_0$, 29
 $\mathcal{P}()$, 29
 ϕ , 88
 $\phi()$, 148
 $\phi^A()$, 25
 $\phi^\cdot, \phi_\rho^\cdot, \phi^{\cdot\cdot}$, 128
 Π^+, Π^- , 57
 $\Pi^\times, \Pi^-$, 58
 $\pi(\text{char.})$, 11
 $\pi(\text{repn.})$, 12

- $\pi^+, \pi^-, 8, 35, 99$
 $\pi^\times, \pi^-, 99$
 $Q(,), 48$
 $\bar{Q}(,), 48$
 $q, 11$
 $R_{E/F}$, *see* Index: restriction of scalars
 $GL(1), 53$
 $GL(2), 70$
 $\text{repn.}(,), \text{repn.}(\times \varepsilon), \text{repn.}()A$, *see* Index: convolution operator
 $\rho, 3$
 $\rho(\varepsilon)$, *see* $\rho_\omega(\varepsilon)$
 ρ_ω , *see* ρ
 $\rho_\omega(\varepsilon), 5$
 $\rho_{\mathcal{H}}, 15$
 $SO(,),$ *see* Index: orbital integral, stable
 $SO_\delta(,),$ *see* Index: orbital integral, stable
 $ST_e^{**}(,), 42$
 $\text{St}_2, 12, 57, 100$
 $s, 15$
 $s_1, s_2, 16$
 $s_\alpha, s_\beta, \dots$, etc., 32
 $\sigma, 11, 16$
 $\sigma^*, 18, 20$
 ${}^\sigma \text{char.}, 31$
 ${}^\sigma \text{repn.}, 62, 87$
 $T_1, 92$
 $T_2, 94$
 $T_{\mathcal{K}}, 88$
 $T_e(,), 42$
 $\bar{T}_2, 94$
 $\otimes \otimes, 32$
 $\otimes \otimes_1, 92$
 $\times \rtimes, 11, 32$
 $\otimes_1, 86$
 $\otimes_2, 86$
 $\text{tl}^{\text{stable}}, 48$
 $\text{tr}\{\text{packet}\}(,), 48$
 $t(,),$ *see also* Index: Frobenius-Hecke class, 53
 $\text{char.} \otimes \text{repn.}, 87$
 $\text{char.} \rtimes \text{repn.}, 87$
 $\text{char.} \otimes \text{repn.}, 34$
 $\text{char.} \rtimes \text{repn.}, 34$
 $\text{repn.} \otimes \text{char.}, 33$
 $\text{repn.} \rtimes \text{char.}, 33$
 $\sim, 22$
 $(\boxtimes) \boxplus (\boxtimes), 9$
 $\otimes_1, 8$
 $\otimes_2, 8$
 $\rtimes, 11$
 $V, 3$
 $V_2^+, V_2^-, 87$
 $V_\pi, 5$
 $V_k, 64$
 $V^{\text{un}}(E/F), 46$
 $\mathcal{V}, 32$
 $v(,), 40$
 W , *see* Index: Weyl groups
 $W(A_0, H), 29$
 $W(,), 30$
 W_F , *see* Index: Weil group
 $W_{M \setminus G}, 90$
 $w, 6$
 $X^*(,), 18$
 $X_*(,), 18$
 $X_*(\mathbf{A}_M), 29$
 $\xi, 8, 15, 22, 27$
 $\xi_1, 17\text{--}19$
 $\xi_2, 19\text{--}21$
 ξ_H , *see also* ξ
 $\mathbf{Z}, 3$
 $\mathbf{Z}_0, 3, 25, 29, 85$
 $Z_0(H_1), Z_0(H_2), 86$
 $Z, 7$

Index

- absolutely semisimple, 128
- Arthur's trace formula, 6
- Arthur, James, 5, 6, 59
- atomic measure, 47
- automorphic
 - forms, 4
 - representation, 3
- bad places, 67
- base change, 9, 66, 67, 95, 121, 123–125
- central character
 - split, 86
- conjugacy, 22
 - F -conjugacy, 22
 - $\bar{F}$ -conjugacy, *see* conjugacy, stable
 - stable, 22
- conjugacy class
 - elliptic, 6
 - regular, 6
- continuous spectrum, 4
- contribution, 51
 - exclusive, 68
- convolution operator, 4, 7, 29–31, 86
- discrete spectrum, 4
- discriminant, 133
- dual group, 18, 20
- element
 - elliptic, 22, 128–133
 - G -regular, 23
 - parabolic, 40
 - regular, 22
 - semisimple, 22
 - singular, 41
 - type, 69, 103
- elliptic, *see* element, elliptic
- elliptic component, *see* elliptic function, *see* representation, elliptic
- elliptic function, 46
- endoscopic
 - data, 15–21
 - equivalent, 16
 - group, 6, 7, 15–21, 42, 86
- endoscopic group, 96
- ε -character, *see* Harish-Chandra character, 7
- ε -endoscopic group, *see* endoscopic, group
- ε -trace identity, 51
- F-H class, *see* Frobenius-Hecke class
- fine χ -expansion, *see also* trace formula, spectral side, 29–40
- Frobenius-Hecke class, 8, 45, 52–57
- function
 - elliptic, 41
 - regular, 14
 - type, 69
- Fundamental Lemma, 11, 27, 46, 127
- Galois action, 16
- global datum, 45–46
- H_1 , 96
- H_2 , 96
- Haar measure, 7, 86
- Harish-Chandra, 66
- Harish-Chandra character, 66
 - twisted, 66, 89
- Hecke algebra, 4, 27, 127
- Heisenberg parabolic subgroup, *see* parabolic subgroup, Heisenberg
- Howe, Piatetski-Shapiro, 9
- intertwining operator, 5, 10, 30, 37, 85, 87–88
- Iwasawa decomposition, 89
- Kazhdan decomposition, 128
- Kottwitz-Shelstad's trace formula, 6, 42–44
- L -
 - action, 15
 - embedding, 17, 19, 22, 27
 - group, 8, 15
 - homomorphism, 15
- L -function
 - partial, 64
- Langlands correspondence, 16
- Langlands quotient, 59, 113
- lattice, 23

- Lebesgue measure, 49
- Levi subgroup, 29
- lift, *see also* lifting, global, *see also* lifting, local, 8, 46
- lifting
 - global, 8, 123–125
 - local, 113–122
- logarithmic derivative, 30
- matching functions, 11, 26
 - weakly, 93, 94, 97, 98
- matrix coefficient, 41
- maximal compact subgroup, 29
- maximal torus, 15–21
 - elliptic, 22, 128–133
- module, *see* representation
- multiplicity formula, 36, 58, 104
- multiplicity one property, 4, 82
- norm, 7, 22
- norm correspondence, 11, 22–26, 128–133
- norm mapping, 18, 21, 35, 53
- $\mathcal{O}$ -expansion, *see* trace formula, geometric side
- orbital integral, 26, 92, 127
 - stable, 26, 42, 127
 - twisted, 89
 - weighted, 40
- packet, 7, 8, 57–66, 99
 - discrete spectrum, 7, 81
 - ε -invariant, 45, 59–60, 81, 99
 - global, 7, 36, 45, 123–125
 - S -part, 45
 - local, 7, 36, 99–122
 - multi-, 51, 60–66
 - quasi-, 7, 57–60, 81, 99, 123–125
 - Saito, Kurokawa, 58
 - stable, 8, 59, 81, 113–125
 - unstable, 8, 57–60, 81, 99–112, 123–125
- parabolic subgroup, 29
 - Heisenberg, 11, 87
 - standard, 30
- quadruple, 30
- regular, *see* element, regular
- representation
 - ε -elliptic, 66
 - ε -invariant, 5, 7, 83, 85
 - admissible, 6, 10
 - automorphic, 83
 - contragredient, 64
 - cuspidal, 106–112
 - discrete spectrum, 30, 83
 - discretely occurring, 31
 - elliptic, 66, 83
 - ε -invariant, 10
 - fully induced, *see* representation, induced
 - induced, 80, 86, 96, 124–125
 - monomial, 8, 31, 86, 109–112
 - normalizedly induced, 29, 87
 - one dimensional, 100
 - right regular, 3, 29
 - unramified, 4, 8
- restricted tensor product, 36, 45
- restriction of scalars, 7, 18, 123–125
- Riesz Representation Theorem, 50
- root datum, 15–21
- root system, 31
- Satake transform, 47
- semisimple, *see* element, semisimple
- σ -invariant, 31
- similitude character, 6
- similitude factor, 6, 85
- splitting, 15–21
- Stone-Weierstrass Theorem, 50
- symplectic group of similitudes, *see* List of Symbols: $\mathrm{GSp}(2)$
- Tamagawa measure, 30
- topologically unipotent, 128
- trace formula, *see also* Arthur's trace formula
 - continuous part, 31
 - discrete part, 31
 - ε -twisted, 5, 6
 - ε -twisted, 30
 - geometric side, 6, 42
 - elliptic part, 6
 - spectral side, *see also* fine χ -expansion, 6, 42
 - stable part, 36
- trace identity, *see* ε -trace identity

transfer factor, 26, 93, 94, 97, 98, 127,
138, 143, 147, 152

truncation, 6, 30

truncation parameter, *see* truncation

Weil group, 8

Weyl group, 23

Weyl integration formula, 91

Yoshida, 9